



planetmath.org

Math for the people, by the people.

optional process

Canonical name	OptionalProcess
Date of creation	2013-03-22 18:37:34
Last modified on	2013-03-22 18:37:34
Owner	gel (22282)
Last modified by	gel (22282)
Numerical id	5
Author	gel (22282)
Entry type	Definition
Classification	msc 60G07
Related topic	ProgressivelyMeasurableProcess
Related topic	PredictableProcess
Defines	optional

Suppose we are given a <http://planetmath.org/FiltrationOfSigmaAlgebrasfiltration>  $(\mathcal{F})_{t \in \mathbb{T}}$  on a measurable space  $(\Omega, \mathcal{F})$ . A stochastic process is said to be adapted if  $X_t$  is  $\mathcal{F}_t$ -measurable for every time  $t$  in the index set  $\mathbb{T}$ . For an arbitrary, uncountable, index set  $\mathbb{T} \subseteq \mathbb{R}$ , this property is too restrictive to be useful. Instead, we can impose measurability conditions on  $X$  considered as a map from  $\mathbb{T} \times \Omega$  to  $\mathbb{R}$ . For instance, we could require  $X$  to be progressively measurable, but that is still too weak a condition for many purposes. A stronger condition is for  $X$  to be *optional*. The index set  $\mathbb{T}$  is assumed to be a closed subset of  $\mathbb{R}$  in the following definition.

The class of optional processes forms the smallest set containing all adapted and right-continuous processes, and which is closed under taking limits of sequences of processes.

The  $\sigma$ -algebra,  $\mathcal{O}$ , on  $\mathbb{T} \times \Omega$  generated by the right-continuous and adapted processes is called the *optional*  $\sigma$ -algebra. Then, a process is optional if and only if it is  $\mathcal{O}$ -measurable.

Alternatively, the optional  $\sigma$ -algebra may be defined as

$$\mathcal{O} = \sigma(\{[T, \infty) : T \text{ is a stopping time}\}).$$

Here,  $[T, \infty)$  is a stochastic interval, consisting of the pairs  $(t, \omega) \in \mathbb{T} \times \Omega$  such that  $T(\omega) \leq t$ . In continuous-time, the equivalence of these two definitions for  $\mathcal{O}$  does require mild conditions on the filtration — it is enough for  $\mathcal{F}_t$  to be universally complete.

In the discrete-time case where the index set  $\mathbb{T}$  countable, then the definitions above imply that a process  $X_t$  is optional if and only if it is adapted.