

planetmath.org

Math for the people, by the people.

proof of Borel-Cantelli 2

Canonical name ProofOfBorelCantelli2
Date of creation 2013-03-22 14:29:35
Last modified on 2013-03-22 14:29:35

Owner kshum (5987) Last modified by kshum (5987)

Numerical id 4

Author kshum (5987)

Entry type Proof

Classification msc 60A99

Let E denote the set of samples that are in A_i infinitely often. We want to show that the complement of E has probability zero.

As in the proof of Borel-Cantelli 1, we know that

$$E^c = \bigcup_{k=1}^{\infty} \bigcap_{i=k}^{\infty} A_i^c$$

where the superscript c means set complement. But for each k,

$$P(\cap_{i=k} A_i^c) = \prod_{i=k}^{\infty} P(A_i^c)$$
$$= \prod_{i=k}^{\infty} (1 - P(A_i))$$

Here we use the assumption that the event A_i 's are independent. The inequality $1-a \leq e^{-a}$ and the assumption that the sum of $P(A_i)$ diverges together imply that

$$P(\cap_{i=k} A_i^c) \le \exp(-\sum_{i=k}^{\infty} P(A_i)) = 0$$

Therefore E^c is a union of countable number of events, each of them has probability zero. So $P(E^c) = 0$.