



planetmath.org

Math for the people, by the people.

Doob’s inequalities

Canonical name	DoobsInequalities
Date of creation	2013-03-22 18:39:52
Last modified on	2013-03-22 18:39:52
Owner	gel (22282)
Last modified by	gel (22282)
Numerical id	7
Author	gel (22282)
Entry type	Theorem
Classification	msc 60G46
Classification	msc 60G44
Classification	msc 60G42
Synonym	Doob’s inequality
Related topic	KolmogorovsMartingaleInequality
Defines	Doob’s maximal quadratic inequality

Doob's inequalities place bounds on the maximum value attained by a martingale in terms of the terminal value. We consider a process $(X_t)_{t \in \mathbb{T}}$ defined on the filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F})_{t \in \mathbb{T}}, \mathbb{P})$. The associated maximum process (X_t^*) is

$$X_t^* \equiv \sup_{s \leq t} |X_s|.$$

The notation $\|\cdot\|_p$ for the <http://planetmath.org/LpSpace> L^p -norm of a random variable will be used. In discrete-time or, more generally whenever the index set \mathbb{T} is countable, then Doob's inequalities are as follows.

Theorem 1 (Doob). *Let $(X_t)_{t \in \mathbb{T}}$ be a submartingale with countable index set \mathbb{T} . Then,*

$$\mathbb{P}\left(\sup_{s \leq t} X_s \geq K\right) \leq K^{-1} \mathbb{E}[(X_t)_+] \quad (1)$$

If X is either a martingale or nonnegative submartingale then,

$$\mathbb{P}(X_t^* \geq K) \leq K^{-1} \mathbb{E}[|X_t|], \quad (2)$$

$$\|X_t^*\|_p \leq \frac{p}{p-1} \|X_t\|_p. \quad (3)$$

for every $K > 0$ and $p > 1$.

In particular, (??) shows that the maximum of any L^p -bounded martingale is itself L^p -bounded and, martingales X^n converge to X in the L^p -norm if and only if $(X^n - X)^* \rightarrow 0$ in the L^p -norm. The special case where $p = 2$ gives

$$\mathbb{E}[(X_t^*)^2] \leq 4\mathbb{E}[X_t^2]$$

which is known as *Doob's maximal quadratic inequality*.

Similarly, (??) shows that any L^1 -bounded martingale is almost surely bounded and that convergence in the L^1 -norm implies ucp convergence. Inequality (??) is also known as Kolmogorov's submartingale inequality.

Doob's inequalities are often applied to continuous-time processes, where $\mathbb{T} = \mathbb{R}_+$. In this case, $X_t^* = \sup_{s \leq t} |X_s|$ is a supremum of uncountably many random variables, and need not be measurable. Instead, it is typically assumed that the processes are right-continuous, in which case, for any $t > 0$ the supremum may be restricted to the countable set

$$\mathbb{T}' = \{s \in \mathbb{R}_+ : s/t \in \mathbb{Q}\}.$$

Putting this into Theorem ?? gives the following continuous-time version of the inequalities.

Theorem 2 (Doob). *Let $(X_t)_{t \in \mathbb{R}_+}$ be a right-continuous submartingale. Then,*

$$\mathbb{P} \left(\sup_{s \leq t} X_s \geq K \right) \leq K^{-1} \mathbb{E}[X_t]$$

for every $K > 0$. If X is right-continuous and either a martingale or non-negative submartingale then,

$$\begin{aligned} \mathbb{P}(X_t^* \geq K) &\leq K^{-1} \mathbb{E}[|X_t|], \\ \|X_t^*\|_p &\leq \frac{p}{p-1} \|X_t\|_p. \end{aligned}$$

for every $K > 0$ and $p > 1$.