

## planetmath.org

Math for the people, by the people.

## gale

Canonical name Gale

Date of creation 2013-03-22 16:43:37 Last modified on 2013-03-22 16:43:37 Owner skubeedooo (5401) Last modified by skubeedooo (5401)

Numerical id 5

Author skubeedooo (5401)

Entry type Definition
Classification msc 60G46
Classification msc 60G44
Classification msc 60G42
Defines supergale
Defines gale

Defines supermartingale

Defines succeed

Defines succeed strongly

Defines success set

Defines strong success set

Let  $\nu$  be a probability measure on Cantor space C, and let  $s \in [0, \infty)$ .

1. A  $\nu$ -s-supergale is a function  $d:\{0,1\}^* \to [0,\infty)$  that satisfies the condition

$$d(w)\nu(w)^{s} \ge d(w0)\nu(w0)^{s} + d(w1)\nu(w1)^{s} \tag{1}$$

for all  $w \in \{0,1\}^*$ , the set of all finite strings of 0's and 1's (including e, the empty string).

- 2. A  $\nu$ -s-gale is a  $\nu$ -s-supergale that satisfies the condition with equality for all  $w \in \{0,1\}^*$ .
- 3. A  $\nu$ -supermartingale is a  $\nu$ -1-supergale.
- 4. A  $\nu$ -martingale is a  $\nu$ -1-gale.
- 5. An s-supergale is a  $\mu$ -s-supergale, where  $\mu$  is the uniform probability measure.
- 6. An s-gale is a  $\mu$ -s-gale.
- 7. A *supermartingale* is a 1-supergale.
- 8. A martingale is a 1-gale.

Put in another way, a martingale is a function  $d: \{0,1\}^* \to [0,\infty)$  such that, for all  $w \in \{0,1\}^*$ , d(w) = (d(w0) + d(w1))/2.

Let d be a  $\nu$ -s-supergale, where  $\nu$  is a probability measure on  $\mathbf{C}$  and  $s \in [0, \infty)$ . We say that d succeeds on a sequence  $S \in \mathbf{C}$  if

$$\limsup_{n \to \infty} d(S[0..n-1]) = \infty.$$

The success set of d is  $S^{\infty}[d] = \{S \in \mathbb{C} | d \text{ succeeds on } S\}$ . d succeeds on a language  $A \subseteq \{0,1\}^*$  if d succeeds on the characteristic sequence  $\chi_A$  of A. We say that d succeeds strongly on a sequence  $S \in \mathbb{C}$  if

$$\liminf_{n \to \infty} d(S[0..n-1]) = \infty.$$

The strong success set of d is  $S_{\text{str}}^{\infty}[d] = \{S \in \mathbf{C} | d \text{ succeeds strongly on } S\}.$ 

Intuitively, a supergale d is a betting strategy that bets on the next bit of a sequence when the previous bits are known. s is the parameter that tunes

the fairness of the betting. The smaller s is, the less fair the betting is. If d succeeds on a sequence, then the bonus we can get from applying d as the betting strategy on the sequence is unbounded. If d succeeds strongly on a sequence, then the bonus goes to infinity.