



planetmath.org

Math for the people, by the people.

type of a distribution function

Canonical name	TypeOfADistributionFunction
Date of creation	2013-03-22 16:25:48
Last modified on	2013-03-22 16:25:48
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	13
Author	CWoo (3771)
Entry type	Definition
Classification	msc 60E05
Classification	msc 62E10
Synonym	centering factor
Synonym	scale parameter
Synonym	location parameter
Defines	type
Defines	scale factor
Defines	location factor
Defines	standard distribution function
Defines	location family
Defines	scale family

Two distribution functions $F, G : \mathbb{R} \rightarrow [0, 1]$ are said to be of the same *type* if there exist $a, b \in \mathbb{R}$ such that $G(x) = F(ax + b)$. a is called the *scale parameter*, and b the *location parameter* or *centering parameter*. Let's write $F \stackrel{t}{=} G$ to denote that F and G are of the same type.

Remarks.

- Necessarily $a > 0$, for otherwise at least one of $G(-\infty) = 0$ or $G(\infty) = 1$ would be violated.
- If $G(x) = F(x + b)$, then the graph of G is *shifted* to the right from the graph of F by b units, if $b > 0$ and to the left if $b < 0$.
- If $G(x) = F(ax)$, then the graph of G is *stretched* from the graph of F by a units if $a > 1$, and *compressed* if $a < 1$.
- If X and Y are random variables whose distribution functions are of the same type, say, F and G respectively, and related by $G(x) = F(ax + b)$, then X and $aY + b$ are identically distributed, since

$$P(X \leq z) = F(z) = G\left(\frac{z - b}{a}\right) = P\left(Y \leq \frac{z - b}{a}\right) = P(aY + b \leq z).$$

When X and $aY + b$ are identically distributed, we write $X \stackrel{t}{=} Y$.

- Again, suppose X and Y correspond to F and G , two distribution functions of the same type related by $G(x) = F(ax + b)$. Then it is easy to see that $E[X] < \infty$ iff $E[Y] < \infty$. In fact, if the expectation exists for one, then $E[X] = aE[Y] + b$. Furthermore, $Var[X]$ is finite iff $Var[Y]$ is. And in this case, $Var[X] = a^2 Var[Y]$. In general, convergence of moments is a “typical” property.
- We can partition the set of distribution functions into disjoint subsets of functions belonging to the same types, since the binary relation $\stackrel{t}{=}$ is an equivalence relation.
- By the same token, we can classify all real random variables defined on a fixed probability space according to their distribution functions, so that if X and Y are of the same type τ iff their corresponding distribution functions F and G are of type τ .

- Given an equivalence class of distribution functions belonging to a certain type τ , such that a random variable Y of type τ exists with finite expectation and variance, then there is one distribution function F of type τ corresponding to a random variable X such that $E[X] = 0$ and $Var[X] = 1$. F is called the *standard distribution function* for type τ . For example, the standard (cumulative) normal distribution is the standard distribution function for the type consisting of all normal distribution functions.
- Within each type τ , we can further classify the distribution functions: if $G(x) = F(x + b)$, then we say that G and F belong to the same *location family* under τ ; and if $G(x) = F(ax)$, then we say that G and F belong to the same *scale family* (under τ).