



Math for the people, by the people.

martingale proof of Kolmogorov's strong law
for square integrable variables

Canonical name	MartingaleProofOfKolmogorovsStrongLawForSquareIntegrableVariables
Date of creation	2013-03-22 18:33:51
Last modified on	2013-03-22 18:33:51
Owner	gel (22282)
Last modified by	gel (22282)
Numerical id	4
Author	gel (22282)
Entry type	Proof
Classification	msc 60F15
Classification	msc 60G42
Related topic	MartingaleConvergenceTheorem
Related topic	KolmogorovsStrongLawOfLargeNumbers
Related topic	StrongLawOfLargeNumbers
Related topic	ProofOfKolmogorovsStrongLawForIIDRandomVariables

We apply the martingale convergence theorem to prove the following result.

Theorem. *Let X_1, X_2, \dots be independent random variables such that $\sum_n \text{Var}[X_n]/n^2 < \infty$. Then, setting*

$$S_n = \frac{1}{n} \sum_{k=1}^n (X_k - \mathbb{E}[X_k])$$

we have $S_n \rightarrow 0$ as $n \rightarrow \infty$, with probability one.

To prove this, we start by constructing a martingale,

$$M_n = \sum_{k=1}^n \frac{X_k - \mathbb{E}[X_k]}{k}.$$

If \mathcal{F}_n is the <http://planetmath.org/SigmaAlgebra> σ -algebra generated by X_1, \dots, X_n then

$$\mathbb{E}[M_{n+1} \mid \mathcal{F}_n] = M_n + \frac{\mathbb{E}[X_{n+1} \mid \mathcal{F}_n] - \mathbb{E}[X_{n+1}]}{n+1} = M_n.$$

Here, the independence of X_{n+1} and \mathcal{F}_n has been used to imply that $\mathbb{E}[X_{n+1} \mid \mathcal{F}_n] = \mathbb{E}[X_{n+1}]$. So, M is a martingale with respect to the filtration $(\mathcal{F}_n)_{n \in \mathbb{N}}$.

Also, by the independence of the X_n , the variance of M_n is

$$\text{Var}[M_n] = \sum_{k=1}^n \text{Var}[X_k/k] \leq \sum_{k=1}^{\infty} \frac{\text{Var}[X_k]}{k^2} < \infty.$$

So, the inequality $\mathbb{E}[|M_n|] \leq \sqrt{\mathbb{E}[M_n^2]} = \sqrt{\text{Var}[M_n]}$ shows that M is an L^1 -bounded martingale, and the martingale convergence theorem says that the limit $M_\infty = \lim_{n \rightarrow \infty} M_n$ exists and is finite, with probability one.

The strong law now follows from Kronecker's lemma, which states that for sequences of real numbers x_1, x_2, \dots and $0 < b_1, b_2, \dots$ such that b_n strictly increases to infinity and $\sum_n x_n/b_n$ converges to a finite limit, then $b_n^{-1} \sum_{k=1}^n x_k$ tends to 0 as $n \rightarrow \infty$. In our case, we take $x_n = X_n - \mathbb{E}[X_n]$ and $b_n = n$ to deduce that $n^{-1} \sum_{k=1}^n (X_k - \mathbb{E}[X_k])$ converges to zero with probability one.

References

- [1] David Williams, *Probability with martingales*, Cambridge Mathematical Textbooks, Cambridge University Press, 1991.
- [2] Olav Kallenberg, *Foundations of modern probability*, Second edition. Probability and its Applications. Springer-Verlag, 2002.