



Math for the people, by the people.

conditional independence

Canonical name	ConditionalIndependence
Date of creation	2013-03-22 16:25:09
Last modified on	2013-03-22 16:25:09
Owner	CWoo (3771)
Last modified by	CWoo (3771)
Numerical id	4
Author	CWoo (3771)
Entry type	Definition
Classification	msc 60A05
Defines	conditionally independent

Let (Ω, \mathcal{F}, P) be a probability space.

Conditional Independence Given an Event

Given an event $C \in \mathcal{F}$:

1. Two events A and B in \mathcal{F} are said to be *conditionally independent given C* if we have the following equality of conditional probabilities:

$$P(A \cap B|C) = P(A|C)P(B|C).$$

2. Two sub sigma algebras $\mathcal{F}_1, \mathcal{F}_2$ of \mathcal{F} are *conditionally independent given C* if any two events $A \in \mathcal{F}_1$ and $B \in \mathcal{F}_2$ are conditionally independent given C .
3. Two real random variables $X, Y : \Omega \rightarrow \mathbb{R}$ are *conditionally independent given event C* if \mathcal{F}_X and \mathcal{F}_Y , the sub sigma algebras generated by X and Y are conditionally independent given C .

Conditional Independence Given a Sigma Algebra

Given a sub sigma algebra \mathcal{G} of \mathcal{F} :

1. Two events A and B in \mathcal{F} are said to be *conditionally independent given \mathcal{G}* if we have the following equality of conditional probabilities (as random variables):

$$P(A \cap B|\mathcal{G}) = P(A|\mathcal{G})P(B|\mathcal{G}).$$

2. Two sub sigma algebras $\mathcal{F}_1, \mathcal{F}_2$ of \mathcal{F} are *conditionally independent given \mathcal{G}* if any two events $A \in \mathcal{F}_1$ and $B \in \mathcal{F}_2$ are conditionally independent given \mathcal{G} .
3. Two real random variables $X, Y : \Omega \rightarrow \mathbb{R}$ are *conditionally independent given event \mathcal{G}* if \mathcal{F}_X and \mathcal{F}_Y , the sub sigma algebras generated by X and Y are conditionally independent given \mathcal{G} .
4. Finally, we can define *conditional independence given a random variable*, say $Z : \Omega \rightarrow \mathbb{R}$ in each of the above three items by setting $\mathcal{G} = \mathcal{F}_Z$.