

Pontificia Universidad Católica del Perú Escuela de Posgrado  
Doctorado en Matemáticas

Variedades Complejas

TAREA 3

2020-II

Indicaciones Generales:

- La TAREA 3 puede ser subida a la plataforma Paideia o enviada al correo electrónico jcuadros@pucp.edu.pe.

1. **Toros de dimensión 1.**

Let  $\Lambda \subset \mathbb{C}$  be a lattice, and let  $X := \mathbb{C}/\Lambda$  be the associated complex torus.

- a) Show that  $X$  is diffeomorphic to  $S^1 \times S^1$ .
- b) Let  $\varphi : \mathbb{C}/\Lambda \rightarrow \mathbb{C}/\Lambda'$  be a biholomorphic map such that  $\varphi(0) = 0$ . Show that there exists a unique  $\alpha \in \mathbb{C}^* = \mathbb{C} - \{0\}$  such that  $\alpha\Lambda = \Lambda'$  and such that the diagram

$$\begin{array}{ccc} \mathbb{C} & \xrightarrow{z \mapsto \alpha z} & \mathbb{C} \\ \pi \downarrow & & \pi' \downarrow \\ \mathbb{C}/\Lambda & \xrightarrow{\varphi} & \mathbb{C}/\Lambda' \end{array}$$

commutes. (Hint: recall that the group of biholomorphic automorphisms of  $\mathbb{C}$  is  $\text{Aut}(\mathbb{C}) = \{z \mapsto \alpha z + \beta \mid \alpha \in \mathbb{C}^*, \beta \in \mathbb{C}\}$ .)

- c) Show that  $X$  is biholomorphic to a torus of the form  $X(\tau) := \mathbb{C}/(\mathbb{Z} + \mathbb{Z}\tau)$  where  $\tau \in \mathbb{C}$  such that  $\text{Im}(\tau) > 0$
- d) Let  $\mathbb{H} := \{\tau \in \mathbb{C} \mid \text{Im}(\tau) > 0\}$  be the Poincaré upper half plane. We define a group action

$$\text{SL}(2, \mathbb{Z}) \times \mathbb{H} \rightarrow \mathbb{H}, \left( \begin{pmatrix} a & b \\ c & d \end{pmatrix}, \tau \right) \mapsto \frac{a\tau + b}{c\tau + d}$$

Show that the biholomorphic equivalence classes of complex tori of dimension 1 have a natural bijection with  $\mathbb{H}/\text{SL}(2, \mathbb{Z})$ .

(The interesting aspect of the set  $\mathbb{H}/\text{SL}(2, \mathbb{Z})$  is that it has a natural complex structure and that  $\mathbb{H}/\text{SL}(2, \mathbb{Z})$  is biholomorphic to  $\mathbb{C}$ .)

## 2. Teorema de Chow.

Recall **Chow's theorem**: Let  $X$  be a compact complex manifold that admits an embedding  $X \hookrightarrow \mathbb{P}^N$  into some projective space. Then  $X$  is algebraic, i.e. defined by a finite number of homogeneous polynomials.

From the previous homework, recall that for  $V$  be a complex vector space of dimension  $n$ . For an integer  $0 < r < n$ , the **Grassmannian**  $G_r(V) := \{S \subset V \text{ subspace of dimension } r\}$  is a complex manifold. We define a map

$$\psi : G_k(V) \rightarrow \mathbb{P}(\bigwedge^k V)$$

as follows: let  $U \subset V$  be a subspace of dimension  $k$  and let  $u_1, \dots, u_k$  be a basis of  $U$ . The multivector

$$u_1 \wedge \dots \wedge u_k$$

gives a point in  $\mathbb{P}(\bigwedge^k V)$ .

- a) Show that this map is well-defined.
- b) Show that defines an embedding, the so-called Plücker embedding and conclude that  $G_k(V)$  is a projective manifold.
- c) From Chow' Theorem, the Grassmannian  $G_k(V)$  is algebraic. Exhibit this explicitly for  $G_2(V)$  with  $V := \mathbb{C}^4$ , with canonical basis  $e_1, \dots, e_4$ : since every 2-vector  $w \in \Lambda^2 \mathbb{C}^4$  has a unique decomposition

$$w = X_0 e_1 \wedge e_2 + X_1 e_1 \wedge e_3 + X_2 e_1 \wedge e_4 + X_3 e_2 \wedge e_3 + X_4 e_2 \wedge e_4 + X_5 e_3 \wedge e_4,$$

show that for the homogeneous coordinates  $[X_0 : \dots : X_5]$  on  $\mathbb{P}(\Lambda^2 \mathbb{C}^4)$ , the Plücker embedding of  $G_2(\mathbb{C}^4)$  in  $\mathbb{P}(\Lambda^2 \mathbb{C}^4) \simeq \mathbb{P}^5$  has the equation

$$X_0 X_5 - X_1 X_4 + X_2 X_3 = 0.$$

For more on Plücker embeddings, check <http://www.math.uchicago.edu/~may/VIGRE/VIGRE2007/REUPapers/FINALFULL/Hudec.pdf> for instance.