

AP[®] STATISTICS SCORE-5 BLUEPRINT SERIES —
VOLUME 1

AP[®] Statistics Formula & Inference Decision Guide

*The High-Yield Score-5 Cram Companion: Rapid
Formula Breakdowns, Inference Decision Trees, and
Fill-in-the-Blank FRQ Sentence Frames*

PR

2027 Exam Edition

Published for Independent Amazon KDP Distribution

© 2027 PR. All rights reserved.

No part of this publication may be reproduced, distributed, or transmitted in any form or by any means, including photocopying, recording, or other electronic or mechanical methods, without the prior written permission of the author and publisher, except in the case of brief quotations embodied in critical reviews and certain other noncommercial uses permitted by copyright law.

® Trademark Notice & Nominative Fair Use Disclaimer:

AP® and Advanced Placement® are trademarks registered by the College Board, which was not involved in the production of, and does not endorse, this product. TI-84® is a registered trademark of Texas Instruments. All trademarks, service marks, and trade names referred to in this book are the property of their respective owners and are used herein for nominative, educational, and descriptive purposes only under Fair Use rules.

Free Companion Online Portal & Resources:

Access downloadable PDF cheat sheets, interactive diagnostic drills, and live formula lookups:



Scan on Mobile to Access Interactive Web Companion

<https://narayanakdp.github.io/ap-stats-score5-blueprint/>

Published Independently via Amazon KDP

First Edition: 2027

ISBN / ASIN: Assigned by Amazon KDP

How to Use This Companion to Score a 5

Welcome to the **AP Statistics Formula & Inference Decision Guide!** This book was engineered specifically to solve the two biggest complaints students have with standard AP prep books:

1. **Textbook Bloat:** Standard books are 600+ pages of dense theory that take weeks to read.
2. **The FRQ Point Leak:** Students lose partial credit on Free-Response Questions not because their math is wrong, but because their verbal phrasing fails to satisfy strict AP rubric wording.

The 3-Step Score-5 System

- Step 1: Master the Formulas & Parameters (Chapters 1–9):** For every formula on the official College Board reference sheet, review the plain-English parameter translation, when to use it, and what conditions are required.
- Step 2: Navigate the Master Inference Flowchart (Chapter 10):** Before starting any hypothesis test or confidence interval problem, follow the decision tree to ensure you select the correct procedure without second-guessing.
- Step 3: Eliminate Grader Traps & Memorize Sentence Frames (Chapters 11–12):** Read through the Top 25 Exam Traps and adopt the exact fill-in-the-blank sentence frames for p -values, confidence levels, slope, and r^2 so that graders award you full points.

Pro Tip for the Final 14 Days Before Exam Day

Don't re-read entire textbooks. Spend 20 minutes each morning reviewing one unit's cheat sheet box and 10 minutes writing out the conditions checklist from memory.

Contents

How to Use This Companion to Score a 5	iii
1 Unit 1: Exploring One-Variable Data	1
1.1 Essential Concepts & Cheat Sheet	1
1.2 Outlier Identification Rules	1
1.3 Resistant vs. Non-Resistant Summary Statistics	2
2 Unit 2: Exploring Two-Variable Data	3
2.1 Essential Concepts & Cheat Sheet	3
3 Unit 3: Collecting Data	4
3.1 Sampling Methods & Experimental Design	4
4 Unit 4: Probability, Random Variables, & Distributions	5
5 Unit 5: Sampling Distributions	6
6 Unit 6: Inference for Proportions	7
7 Unit 7: Inference for Means	8
8 Unit 8: Chi-Square Tests	9
9 Unit 9: Inference for Slopes	10

10 Master Inference Decision Flowchart	11
10.1 Visual Inference Decision Tree	11
10.2 Inference Procedures Lookup Matrix	12
11 Top 25 "Never Forget" AP Exam Traps	13
12 Standard Interpretation Sentence Templates	15

Chapter 1

Unit 1: Exploring One-Variable Data

1.1 Essential Concepts & Cheat Sheet

When describing any quantitative single-variable distribution on the AP Exam, you **must** address four key characteristics: **SOCS** (Shape, Outliers, Center, Spread) always written **in context** of the problem.

Core Formulas: One-Variable Statistics

1. **Mean ($\bar{x}$):** $\bar{x} = \frac{\sum x_i}{n} = \frac{1}{n} \sum x_i$
2. **Sample Standard Deviation (s_x):** $s_x = \sqrt{\frac{1}{n-1} \sum (x_i - \bar{x})^2}$
3. **Standardized Score (z-score):** $z = \frac{x - \mu}{\sigma}$ or $z = \frac{x - \bar{x}}{s_x}$

1.2 Outlier Identification Rules

- **1.5 × IQR Rule:** $IQR = Q_3 - Q_1$. Outliers are values $< Q_1 - 1.5(IQR)$ or $> Q_3 + 1.5(IQR)$.
- **2-SD Rule (Normal data):** Outliers are values outside $\mu \pm 2\sigma$.

1.3 Resistant vs. Non-Resistant Summary Statistics

Measure	Resistant to Outliers / Skew?	When to Use on Exam
Median	Yes (Unaffected by extremes)	Skewed distributions
IQR	Yes (Middle 50% only)	Skewed distributions
Mean	No (Pulled toward tail/outliers)	Symmetric, unimodal distributions
Standard Deviation	No (Squared deviations inflate)	Symmetric, unimodal distributions

AP Exam Trap Alert: Comparing Distributions

When comparing two distributions on an FRQ:

1. You **must** use comparative words (e.g., "*Distribution A has a greater median than Distribution B*").
2. You **must** include context (e.g., grams of sugar, test scores, heights).

Chapter 2

Unit 2: Exploring Two-Variable Data

2.1 Essential Concepts & Cheat Sheet

Core Formulas: Linear Regression

1. **Least-Squares Regression Line (LSRL):** $\hat{y} = a + bx$ ($\hat{y}$ = predicted response)
2. **Slope (b):** $b = r \left(\frac{s_y}{s_x} \right)$
3. **y -Intercept (a):** $a = \bar{y} - b\bar{x}$ (The point $(\bar{x}, \bar{y})$ is always on the line).
4. **Residual (e):** Residual = Actual y – Predicted $\hat{y} = y - \hat{y}$
5. **Standard Error of Residuals (s):** $s = \sqrt{\frac{\sum (y_i - \hat{y}_i)^2}{n-2}}$

Score-5 Interpretation Sentence Templates

1. **Slope (b):** "For each additional 1 [unit of x], the predicted [response variable y] increases/decreases by [value of b] [units of y]."
2. **r^2 :** "Approximately [value of $r^2 \times 100\%$] of the variation in [response variable in context] is accounted for by the linear relationship with [explanatory variable in context]."

Chapter 3

Unit 3: Collecting Data

3.1 Sampling Methods & Experimental Design

Sampling Method	Key Procedure & Purpose
SRS	Every group of n individuals has equal chance of selection. Eliminates selection bias.
Stratified	Homogeneous groups (<i>strata</i>); sample some from <i>all</i> strata. Reduces variability.
Cluster	Heterogeneous groups (<i>clusters</i>); randomly select several clusters and sample <i>all</i> within them. Saves cost.
Systematic	Select every k -th individual from an ordered list.

AP Exam Trap Alert: Confounding vs Bias

Confounding occurs when an outside variable is associated with the explanatory variable AND affects the response variable. Always explain both connections on FRQs!

Chapter 4

Unit 4: Probability, Random Variables, & Distributions

Core Probability Formulas

General Addition Rule: $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

Conditional Probability: $P(A | B) = \frac{P(A \cap B)}{P(B)}$

Linear Combinations: $\mu_{X \pm Y} = \mu_X \pm \mu_Y$, $\sigma_{X \pm Y}^2 = \sigma_X^2 + \sigma_Y^2$ (**Always ADD variances!**)

Binomial Distribution $B(n, p)$: $P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$, $\mu = np$, $\sigma = \sqrt{np(1 - p)}$

Chapter 5

Unit 5: Sampling Distributions

Sampling Distributions Formulas

Sample Proportion ($\hat{p}$): $\mu_{\hat{p}} = p$, $\sigma_{\hat{p}} = \sqrt{\frac{p(1-p)}{n}}$ (Normal if $np \geq 10, n(1-p) \geq 10$)

Sample Mean ($\bar{x}$): $\mu_{\bar{x}} = \mu$, $\sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}}$ (Normal if $n \geq 30$ by CLT or pop normal)

Chapter 6

Unit 6: Inference for Proportions

Proportions Inference Formulas

1-Prop z -Interval: $\hat{p} \pm z^* \sqrt{\frac{\hat{p}(1-\hat{p})}{n}}$

1-Prop z -Test: $z = \frac{\hat{p} - p_0}{\sqrt{\frac{p_0(1-p_0)}{n}}}$

2-Prop z -Test (Pooled): $\hat{p}_c = \frac{x_1 + x_2}{n_1 + n_2}, \quad z = \frac{(\hat{p}_1 - \hat{p}_2) - 0}{\sqrt{\hat{p}_c(1-\hat{p}_c)\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$

Chapter 7

Unit 7: Inference for Means

Means Inference Formulas

1-Sample t -Interval: $\bar{x} \pm t^* \left(\frac{s}{\sqrt{n}} \right)$ (df = $n - 1$)

1-Sample t -Test: $t = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$ (df = $n - 1$)

2-Sample t -Test: $t = \frac{(\bar{x}_1 - \bar{x}_2) - 0}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$ (Pooled = NO)

Chapter 8

Unit 8: Chi-Square Tests

Chi-Square Formulas

$$\chi^2 = \sum \frac{(O-E)^2}{E}, \quad \text{Expected Count} = \frac{(\text{Row Total}) \times (\text{Column Total})}{\text{Table Total}}$$

Condition Alert

ALL Expected Counts MUST be ≥ 5 (never check observed counts).

Chapter 9

Unit 9: Inference for Slopes

Slope Inference


t -Interval for β : $b \pm t^* \cdot SE_b$ (df = $n - 2$)

t -Test for β : $t = \frac{b-0}{SE_b}$ (df = $n - 2$)

Chapter 10

Master Inference Decision Flowchart

10.1 Visual Inference Decision Tree



10.2 Inference Procedures Lookup Matrix

Data Type	Groups	Goal	Procedure Name	Calculator Command
Categorical	1 Sample	Estimate p	1-Prop z-Interval	1-PropZInt
Categorical	1 Sample	Test p	1-Prop z-Test	1-PropZTest
Categorical	2 Samples	Estimate $p_1 - p_2$	2-Prop z-Interval	2-PropZInt
Categorical	2 Samples	Test $p_1 = p_2$	2-Prop z-Test (Pooled)	2-PropZTest
Categorical	1 Sample (Fit)	Fit distribution	χ^2 GOF-Test	χ^2 GOF-Test
Categorical	2+ Samples	Homogeneity	χ^2 Test of Homogeneity	χ^2 -Test (Matrix)
Categorical	1 Sample (2 Var)	Association	χ^2 Test of Independence	χ^2 -Test (Matrix)
Quantitative	1 Sample	Estimate μ	1-Sample t-Interval	TInterval
Quantitative	1 Sample	Test μ	1-Sample t-Test	T-Test
Quantitative	Matched Pairs	Difference μ_d	Paired t-Test/Interval	T-Test on $L_1 - L_2$
Quantitative	2 Indep Samples	Compare $\mu_1 - \mu_2$	2-Sample t-Test/Interval	2-SampTTest
Quantitative	Linear Slope	Slope β	LinReg t-Test/Interval	LinRegTTest

Chapter 11

Top 25 "Never Forget" AP Exam Traps

1. **NEVER** say "Accept H_0 ": Always write "*Fail to reject H_0* ".
2. **NEVER** write "We prove H_a ": Write "*We have convincing statistical evidence that [context]*".
3. Hypotheses in parameters (μ, p, β) , **NEVER** statistics $(\bar{x}, \hat{p}, b)$.
4. Define parameters in context.
5. Use comparative words when comparing distributions (greater, lesser).
6. Include units on **ALL** quantitative statistics and interpretations.
7. Don't forget the hat ($\hat{y}$) on regression equations.
8. Correlation (r) measures **LINEAR** strength only.
9. Association is **NOT** Causation.
10. Blocking is for experiments; Stratification is for sampling.
11. Mutually exclusive events are **NEVER** independent.
12. Always **ADD** variances, **NEVER** add standard deviations.
13. CLT applies to sampling distribution of $\bar{x}$ ($n \geq 30$), not individual data.
14. Use p_0 for 1-Prop Z-Test Large Counts ($np_0 \geq 10, n(1 - p_0) \geq 10$).
15. Use $\hat{p}$ for 1-Prop Z-Interval Large Counts ($n\hat{p} \geq 10, n(1 - \hat{p}) \geq 10$).
16. Chi-square conditions require **EXPECTED** counts ≥ 5 , not observed counts.
17. Chi-square df: $df = (\text{rows} - 1)(\text{cols} - 1)$.

18. Never pool standard deviations for 2-sample t -tests (Pooled = NO).
19. Confidence Interval (plausible values) vs Confidence Level (long-run capture rate).
20. A high r^2 does NOT prove linearity (always check residual plot).
21. Voluntary response produces extreme bias.
22. State the conclusion in terms of H_a .
23. State the significance level (α).
24. Calculator Speak is NOT enough on FRQs (write formulas, df, p -value).
25. Context, Context, Context: Include problem context in every sentence!

Chapter 12

Standard Interpretation Sentence Templates

Template 1: Confidence Interval for a Parameter

"We are [C%] confident that the interval from [Lower Bound] to [Upper Bound] captures the true [proportion / mean] of [context with units]."

Template 2: Confidence Level (C%)

"If we take many random samples of size n and construct [C%] confidence intervals, about [C%] of the intervals will capture the true [parameter in context]."

Template 3: p -Value in Context

"Assuming that [null hypothesis H_0 in context] is true, there is a [p-value] probability of obtaining a sample statistic as extreme as or more extreme than observed, purely by random chance."

Template 4: Test Conclusion (Reject H_0)

"Because $p\text{-value } ([p]) < \alpha$ ($[0.05]$), we reject H_0 . There is convincing statistical evidence that [state H_a in context]."

Template 5: Test Conclusion (Fail to Reject H_0)

"Because $p\text{-value } ([p]) > \alpha$ ($[0.05]$), we fail to reject H_0 . There is NOT convincing statistical evidence that [state H_a in context]."