

Recitation 1

18-460/18-660 Optimization

Chaoyi Pan

Carnegie Mellon University

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

Gradient:

$$\nabla f(\mathbf{x}) = \begin{bmatrix} \partial_{x_1} f(\mathbf{x}) \\ \partial_{x_2} f(\mathbf{x}) \\ \vdots \\ \partial_{x_n} f(\mathbf{x}) \end{bmatrix}$$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

Gradient:

$$\nabla f(\mathbf{x}) = \begin{bmatrix} \partial_{x_1} f(\mathbf{x}) \\ \partial_{x_2} f(\mathbf{x}) \\ \vdots \\ \partial_{x_n} f(\mathbf{x}) \end{bmatrix}$$

Example. $f(\mathbf{x}) = 3x_1^2 + 5 \log x_2$. $\nabla f(\mathbf{x}) = \begin{bmatrix} \\ \end{bmatrix}$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

Gradient:

$$\nabla f(\mathbf{x}) = \begin{bmatrix} \partial_{x_1} f(\mathbf{x}) \\ \partial_{x_2} f(\mathbf{x}) \\ \vdots \\ \partial_{x_n} f(\mathbf{x}) \end{bmatrix}$$

Example. $f(\mathbf{x}) = 3x_1^2 + 5 \log x_2$. $\nabla f(\mathbf{x}) = \begin{bmatrix} 6x_1 \\ \end{bmatrix}$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

Gradient:

$$\nabla f(\mathbf{x}) = \begin{bmatrix} \partial_{x_1} f(\mathbf{x}) \\ \partial_{x_2} f(\mathbf{x}) \\ \vdots \\ \partial_{x_n} f(\mathbf{x}) \end{bmatrix}$$

Example. $f(\mathbf{x}) = 3x_1^2 + 5 \log x_2$. $\nabla f(\mathbf{x}) = \begin{bmatrix} 6x_1 \\ 5x_2^{-1} \end{bmatrix}$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

Hessian:

$$\nabla_f^2(\mathbf{x}) = \begin{bmatrix} \partial_{x_1 x_1}^2 f(\mathbf{x}) & \partial_{x_1 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_1 x_n}^2 f(\mathbf{x}) \\ \partial_{x_2 x_1}^2 f(\mathbf{x}) & \partial_{x_2 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_2 x_n}^2 f(\mathbf{x}) \\ \vdots & \vdots & \ddots & \vdots \\ \partial_{x_n x_1}^2 f(\mathbf{x}) & \partial_{x_n x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_n x_n}^2 f(\mathbf{x}) \end{bmatrix}$$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

Hessian:

$$\nabla_f^2(\mathbf{x}) = \begin{bmatrix} \partial_{x_1 x_1}^2 f(\mathbf{x}) & \partial_{x_1 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_1 x_n}^2 f(\mathbf{x}) \\ \partial_{x_2 x_1}^2 f(\mathbf{x}) & \partial_{x_2 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_2 x_n}^2 f(\mathbf{x}) \\ \vdots & \vdots & \ddots & \vdots \\ \partial_{x_n x_1}^2 f(\mathbf{x}) & \partial_{x_n x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_n x_n}^2 f(\mathbf{x}) \end{bmatrix}$$

Because $f \in \mathbf{C}^2$, $\partial_{x_i x_j}^2 f = \partial_{x_j x_i}^2 f$, and $\nabla_f^2(\mathbf{x})$ is symmetric.

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

Hessian:

$$\nabla_f^2(\mathbf{x}) = \begin{bmatrix} \partial_{x_1 x_1}^2 f(\mathbf{x}) & \partial_{x_1 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_1 x_n}^2 f(\mathbf{x}) \\ \partial_{x_2 x_1}^2 f(\mathbf{x}) & \partial_{x_2 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_2 x_n}^2 f(\mathbf{x}) \\ \vdots & \vdots & \ddots & \vdots \\ \partial_{x_n x_1}^2 f(\mathbf{x}) & \partial_{x_n x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_n x_n}^2 f(\mathbf{x}) \end{bmatrix}$$

Because $f \in \mathbf{C}^2$, $\partial_{x_i x_j}^2 f = \partial_{x_j x_i}^2 f$, and $\nabla_f^2(\mathbf{x})$ is symmetric.

Example. $f(\mathbf{x}) = x_1^3 + 5x_2 + 7x_1x_2$. $\nabla_f^2(\mathbf{x}) = \begin{bmatrix} & & \\ & & \\ & & \end{bmatrix}$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

Hessian:

$$\nabla_f^2(\mathbf{x}) = \begin{bmatrix} \partial_{x_1 x_1}^2 f(\mathbf{x}) & \partial_{x_1 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_1 x_n}^2 f(\mathbf{x}) \\ \partial_{x_2 x_1}^2 f(\mathbf{x}) & \partial_{x_2 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_2 x_n}^2 f(\mathbf{x}) \\ \vdots & \vdots & \ddots & \vdots \\ \partial_{x_n x_1}^2 f(\mathbf{x}) & \partial_{x_n x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_n x_n}^2 f(\mathbf{x}) \end{bmatrix}$$

Because $f \in \mathbf{C}^2$, $\partial_{x_i x_j}^2 f = \partial_{x_j x_i}^2 f$, and $\nabla_f^2(\mathbf{x})$ is symmetric.

Example. $f(\mathbf{x}) = x_1^3 + 5x_2 + 7x_1x_2$. $\nabla_f^2(\mathbf{x}) = \begin{bmatrix} 6x_1 & 7 \\ 7 & 0 \end{bmatrix}$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

Hessian:

$$\nabla_f^2(\mathbf{x}) = \begin{bmatrix} \partial_{x_1 x_1}^2 f(\mathbf{x}) & \partial_{x_1 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_1 x_n}^2 f(\mathbf{x}) \\ \partial_{x_2 x_1}^2 f(\mathbf{x}) & \partial_{x_2 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_2 x_n}^2 f(\mathbf{x}) \\ \vdots & \vdots & \ddots & \vdots \\ \partial_{x_n x_1}^2 f(\mathbf{x}) & \partial_{x_n x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_n x_n}^2 f(\mathbf{x}) \end{bmatrix}$$

Because $f \in \mathbf{C}^2$, $\partial_{x_i x_j}^2 f = \partial_{x_j x_i}^2 f$, and $\nabla_f^2(\mathbf{x})$ is symmetric.

Example. $f(\mathbf{x}) = x_1^3 + 5x_2 + 7x_1x_2$. $\nabla_f^2(\mathbf{x}) = \begin{bmatrix} 6x_1 & 7 \\ 7 & 0 \end{bmatrix}$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

Hessian:

$$\nabla_f^2(\mathbf{x}) = \begin{bmatrix} \partial_{x_1 x_1}^2 f(\mathbf{x}) & \partial_{x_1 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_1 x_n}^2 f(\mathbf{x}) \\ \partial_{x_2 x_1}^2 f(\mathbf{x}) & \partial_{x_2 x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_2 x_n}^2 f(\mathbf{x}) \\ \vdots & \vdots & \ddots & \vdots \\ \partial_{x_n x_1}^2 f(\mathbf{x}) & \partial_{x_n x_2}^2 f(\mathbf{x}) & \cdots & \partial_{x_n x_n}^2 f(\mathbf{x}) \end{bmatrix}$$

Because $f \in \mathbf{C}^2$, $\partial_{x_i x_j}^2 f = \partial_{x_j x_i}^2 f$, and $\nabla_f^2(\mathbf{x})$ is symmetric.

Example. $f(\mathbf{x}) = x_1^3 + 5x_2 + 7x_1x_2$. $\nabla_f^2(\mathbf{x}) = \begin{bmatrix} 6x_1 & 7 \\ 7 & 0 \end{bmatrix}$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_0^1 \nabla f(t\mathbf{y} + (1-t)\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt$$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_0^1 \nabla f(t\mathbf{y} + (1-t)\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt$$

Integration of *projection* of gradients along the line through \mathbf{x} and \mathbf{y} .

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_0^1 \nabla f(t\mathbf{y} + (1-t)\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt$$

Integration of *projection* of gradients along the line through \mathbf{x} and \mathbf{y} .

Proof. Let $\mathbf{z}(t) = t\mathbf{y} + (1-t)\mathbf{x}$ and $h(t) = f(\mathbf{z}(t))$.

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_0^1 \nabla f(t\mathbf{y} + (1-t)\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt$$

Integration of *projection* of gradients along the line through \mathbf{x} and \mathbf{y} .

Proof. Let $\mathbf{z}(t) = t\mathbf{y} + (1-t)\mathbf{x}$ and $h(t) = f(\mathbf{z}(t))$.

$\mathbf{z}(0) = \mathbf{x}$, $\mathbf{z}(1) = \mathbf{y}$.

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_0^1 \nabla f(t\mathbf{y} + (1-t)\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt$$

Integration of *projection* of gradients along the line through \mathbf{x} and \mathbf{y} .

Proof. Let $\mathbf{z}(t) = t\mathbf{y} + (1-t)\mathbf{x}$ and $h(t) = f(\mathbf{z}(t))$.

$$\mathbf{z}(0) = \mathbf{x}, \quad \mathbf{z}(1) = \mathbf{y}.$$

$$h'(t) = \nabla f(\mathbf{z}(t))^\top \begin{bmatrix} z'_1(t) \\ \vdots \\ z'_n(t) \end{bmatrix}$$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_0^1 \nabla f(t\mathbf{y} + (1-t)\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt$$

Integration of *projection* of gradients along the line through \mathbf{x} and \mathbf{y} .

Proof. Let $\mathbf{z}(t) = t\mathbf{y} + (1-t)\mathbf{x}$ and $h(t) = f(\mathbf{z}(t))$.

$$\mathbf{z}(0) = \mathbf{x}, \quad \mathbf{z}(1) = \mathbf{y}.$$

$$h'(t) = \nabla f(\mathbf{z}(t))^\top \begin{bmatrix} z'_1(t) \\ \vdots \\ z'_n(t) \end{bmatrix} = \nabla f(\mathbf{z}(t))^\top \begin{bmatrix} y_1 - x_1 \\ \vdots \\ y_n - x_n \end{bmatrix}$$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_0^1 \nabla f(t\mathbf{y} + (1-t)\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt$$

Integration of *projection* of gradients along the line through \mathbf{x} and \mathbf{y} .

Proof. Let $\mathbf{z}(t) = t\mathbf{y} + (1-t)\mathbf{x}$ and $h(t) = f(\mathbf{z}(t))$.

$$\mathbf{z}(0) = \mathbf{x}, \quad \mathbf{z}(1) = \mathbf{y}.$$

$$h'(t) = \nabla f(\mathbf{z}(t))^\top \begin{bmatrix} z'_1(t) \\ \vdots \\ z'_n(t) \end{bmatrix} = \nabla f(\mathbf{z}(t))^\top \begin{bmatrix} y_1 - x_1 \\ \vdots \\ y_n - x_n \end{bmatrix} = \nabla f(\mathbf{z}(t))^\top (\mathbf{y} - \mathbf{x}).$$

Recap from Calculus

$f : \mathbb{R}^n \rightarrow \mathbb{R}$ is continuously differentiable.

$$f(\mathbf{y}) - f(\mathbf{x}) = \int_0^1 \nabla f(t\mathbf{y} + (1-t)\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt$$

Integration of *projection* of gradients along the line through \mathbf{x} and \mathbf{y} .

Proof. Let $\mathbf{z}(t) = t\mathbf{y} + (1-t)\mathbf{x}$ and $h(t) = f(\mathbf{z}(t))$.

$$\mathbf{z}(0) = \mathbf{x}, \quad \mathbf{z}(1) = \mathbf{y}.$$

$$h'(t) = \nabla f(\mathbf{z}(t))^\top \begin{bmatrix} z_1'(t) \\ \vdots \\ z_n'(t) \end{bmatrix} = \nabla f(\mathbf{z}(t))^\top \begin{bmatrix} y_1 - x_1 \\ \vdots \\ y_n - x_n \end{bmatrix} = \nabla f(\mathbf{z}(t))^\top (\mathbf{y} - \mathbf{x}).$$

$$f(\mathbf{y}) - f(\mathbf{x}) = h(1) - h(0) = \int_0^1 h'(t) dt = \int_0^1 \nabla f(t\mathbf{y} + (1-t)\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) dt.$$

Recap from Calculus

$f : \mathbb{R} \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

For any $x, y \in \mathbb{R}$, there exists $t \in [0, 1]$ such that

$$f(y) = f(x) + f'(x)(y - x) + \frac{1}{2}f''(ty + (1 - t)x)(y - x)^2.$$

Recap from Calculus

$f : \mathbb{R} \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

For any $x, y \in \mathbb{R}$, there exists $t \in [0, 1]$ such that

$$f(y) = f(x) + f'(x)(y - x) + \frac{1}{2}f''(ty + (1 - t)x)(y - x)^2.$$

What if f is multi-variate?

Recap from Calculus

$f : \mathbb{R} \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

For any $x, y \in \mathbb{R}$, there exists $t \in [0, 1]$ such that

$$f(y) = f(x) + f'(x)(y - x) + \frac{1}{2}f''(ty + (1 - t)x)(y - x)^2.$$

What if f is multi-variate?

Hint: Define $h(s) = f(s\mathbf{y} + (1 - s)\mathbf{x})$, and do Taylor's expansion of $h(1)$ on $s = 0$.

Recap from Calculus

$f : \mathbb{R} \rightarrow \mathbb{R}$ is *twice* continuously differentiable.

For any $x, y \in \mathbb{R}$, there exists $t \in [0, 1]$ such that

$$f(y) = f(x) + f'(x)(y - x) + \frac{1}{2}f''(ty + (1 - t)x)(y - x)^2.$$

What if f is multi-variate?

Hint: Define $h(s) = f(s\mathbf{y} + (1 - s)\mathbf{x})$, and do Taylor's expansion of $h(1)$ on $s = 0$.

$$f(\mathbf{y}) = f(\mathbf{x}) + \nabla f(\mathbf{x})^\top (\mathbf{y} - \mathbf{x}) + \frac{1}{2}(\mathbf{y} - \mathbf{x})^\top \nabla_f^2(t\mathbf{y} + (1 - t)\mathbf{x})(\mathbf{y} - \mathbf{x})$$

Cauchy-Schwarz Inequality

$$\mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

$$\mathbf{x}^T \mathbf{y} \leq \|\mathbf{x}\|_2 \|\mathbf{y}\|_2.$$

Cauchy-Schwarz Inequality

$$\mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

$$\mathbf{x}^T \mathbf{y} \leq \|\mathbf{x}\|_2 \|\mathbf{y}\|_2.$$

Proof. Consider (univariate) quadratic function

$$h(t) = \frac{1}{2}(t\mathbf{x} + \mathbf{y})^T(t\mathbf{x} + \mathbf{y}) = \frac{1}{2}\|\mathbf{x}\|_2^2 t^2 + \mathbf{x}^T \mathbf{y} t + \frac{1}{2}\|\mathbf{y}\|_2^2.$$

Cauchy-Schwarz Inequality

$$\mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

$$\mathbf{x}^T \mathbf{y} \leq \|\mathbf{x}\|_2 \|\mathbf{y}\|_2.$$

Proof. Consider (univariate) quadratic function

$$h(t) = \frac{1}{2}(t\mathbf{x} + \mathbf{y})^T(t\mathbf{x} + \mathbf{y}) = \frac{1}{2}\|\mathbf{x}\|_2^2 t^2 + \mathbf{x}^T \mathbf{y} t + \frac{1}{2}\|\mathbf{y}\|_2^2.$$

$$h(t) \geq 0, \quad \forall t \in \mathbb{R}.$$

Cauchy-Schwarz Inequality

$$\mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

$$\mathbf{x}^T \mathbf{y} \leq \|\mathbf{x}\|_2 \|\mathbf{y}\|_2.$$

Proof. Consider (univariate) quadratic function

$$h(t) = \frac{1}{2}(t\mathbf{x} + \mathbf{y})^T(t\mathbf{x} + \mathbf{y}) = \frac{1}{2}\|\mathbf{x}\|_2^2 t^2 + \mathbf{x}^T \mathbf{y} t + \frac{1}{2}\|\mathbf{y}\|_2^2.$$

$$h(t) \geq 0, \quad \forall t \in \mathbb{R}.$$

Reminder: quadratic function $ax^2 + bx + c$ has less than 2 distinct real roots $\Leftrightarrow \Delta = b^2 - 4ac \leq 0$.

Cauchy-Schwarz Inequality

$$\mathbf{x}, \mathbf{y} \in \mathbb{R}^n.$$

$$\mathbf{x}^T \mathbf{y} \leq \|\mathbf{x}\|_2 \|\mathbf{y}\|_2.$$

Proof. Consider (univariate) quadratic function

$$h(t) = \frac{1}{2}(t\mathbf{x} + \mathbf{y})^T(t\mathbf{x} + \mathbf{y}) = \frac{1}{2}\|\mathbf{x}\|_2^2 t^2 + \mathbf{x}^T \mathbf{y} t + \frac{1}{2}\|\mathbf{y}\|_2^2.$$

$$h(t) \geq 0, \quad \forall t \in \mathbb{R}.$$

Reminder: quadratic function $ax^2 + bx + c$ has less than 2 distinct real roots $\Leftrightarrow \Delta = b^2 - 4ac \leq 0$.

$$0 \geq \Delta = (\mathbf{x}^T \mathbf{y})^2 - \|\mathbf{x}\|_2^2 \|\mathbf{y}\|_2^2.$$

Lipschitz Functions

$f : \mathcal{D} \rightarrow \mathbb{R}$ is L -Lipschitz continuous if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$|f(\mathbf{y}) - f(\mathbf{x})| \leq L \|\mathbf{y} - \mathbf{x}\|_2.$$

Lipschitz Functions

$f : \mathcal{D} \rightarrow \mathbb{R}$ is L -Lipschitz continuous if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$|f(\mathbf{y}) - f(\mathbf{x})| \leq L\|\mathbf{y} - \mathbf{x}\|_2.$$

A differentiable $f : \mathcal{D} \rightarrow \mathbb{R}$ has L -Lipschitz continuous gradients if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \leq L\|\mathbf{y} - \mathbf{x}\|_2.$$

Lipschitz Functions

$f : \mathcal{D} \rightarrow \mathbb{R}$ is L -Lipschitz continuous if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$|f(\mathbf{y}) - f(\mathbf{x})| \leq L \|\mathbf{y} - \mathbf{x}\|_2.$$

A differentiable $f : \mathcal{D} \rightarrow \mathbb{R}$ has L -Lipschitz continuous gradients if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \leq L \|\mathbf{y} - \mathbf{x}\|_2.$$

Gradient doesn't have unbounded difference between two close points.

Lipschitz Functions

$f : \mathcal{D} \rightarrow \mathbb{R}$ is L -Lipschitz continuous if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$|f(\mathbf{y}) - f(\mathbf{x})| \leq L \|\mathbf{y} - \mathbf{x}\|_2.$$

A differentiable $f : \mathcal{D} \rightarrow \mathbb{R}$ has L -Lipschitz continuous gradients if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \leq L \|\mathbf{y} - \mathbf{x}\|_2.$$

Gradient doesn't have unbounded difference between two close points.

Note. Differentiable $f : \mathcal{D} \rightarrow \mathbb{R}$ is m -strongly convex \implies for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \geq m \|\mathbf{y} - \mathbf{x}\|_2.$$

Lipschitz Functions

$f : \mathcal{D} \rightarrow \mathbb{R}$ is L -Lipschitz continuous if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$|f(\mathbf{y}) - f(\mathbf{x})| \leq L \|\mathbf{y} - \mathbf{x}\|_2.$$

A differentiable $f : \mathcal{D} \rightarrow \mathbb{R}$ has L -Lipschitz continuous gradients if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \leq L \|\mathbf{y} - \mathbf{x}\|_2.$$

Gradient doesn't have unbounded difference between two close points.

Note. Differentiable $f : \mathcal{D} \rightarrow \mathbb{R}$ is m -strongly convex \implies for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \geq m \|\mathbf{y} - \mathbf{x}\|_2.$$

Proof.

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \stackrel{(1)}{\geq} \left(\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}) \right)^T \left(\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}) \right) = \left(\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}) \right)^T \left(\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}) \right)$$

Lipschitz Functions

$f : \mathcal{D} \rightarrow \mathbb{R}$ is L -Lipschitz continuous if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$|f(\mathbf{y}) - f(\mathbf{x})| \leq L \|\mathbf{y} - \mathbf{x}\|_2.$$

A differentiable $f : \mathcal{D} \rightarrow \mathbb{R}$ has L -Lipschitz continuous gradients if for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \leq L \|\mathbf{y} - \mathbf{x}\|_2.$$

Gradient doesn't have unbounded difference between two close points.

Note. Differentiable $f : \mathcal{D} \rightarrow \mathbb{R}$ is m -strongly convex \implies for every $\mathbf{x}, \mathbf{y} \in \mathcal{D}$,

$$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \geq m \|\mathbf{y} - \mathbf{x}\|_2.$$

Proof.

$\|\nabla f(\mathbf{y}) - \nabla f(\mathbf{x})\|_2 \geq m \|\mathbf{y} - \mathbf{x}\|_2$ (1) $\nabla f(\mathbf{y}) - \nabla f(\mathbf{x}) = \int_0^1 \nabla^2 f(\mathbf{x} + t(\mathbf{y} - \mathbf{x})) dt$ (2)

Eigenvalues and Eigenvectors

Let $A \in \mathbb{R}^{n \times n}$. A nonzero vector v is an eigenvector with eigenvalue $\lambda \in \mathbb{C}$ if

$$Av = \lambda v.$$

Eigenvalues and Eigenvectors

Let $A \in \mathbb{R}^{n \times n}$. A nonzero vector v is an eigenvector with eigenvalue $\lambda \in \mathbb{C}$ if

$$Av = \lambda v.$$

If A is real symmetric ($A = A^T$), then

- ▶ All eigenvalues are real.
- ▶ There exists an orthonormal basis of eigenvectors.
- ▶ A admits the eigendecomposition $A = Q\Lambda Q^T$ with Q orthogonal, $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$.

Rayleigh Quotient and Inequalities (Symmetric A)

For $A = A^T$ and any nonzero $x \in \mathbb{R}^n$, the Rayleigh quotient satisfies

$$\lambda_{\min}(A) \leq \frac{x^T A x}{x^T x} \leq \lambda_{\max}(A).$$

Rayleigh Quotient and Inequalities (Symmetric A)

For $A = A^T$ and any nonzero $x \in \mathbb{R}^n$, the Rayleigh quotient satisfies

$$\lambda_{\min}(A) \leq \frac{x^T A x}{x^T x} \leq \lambda_{\max}(A).$$

Equivalent statements:

- ▶ $A \succeq 0$ (positive semidefinite, PSD) $\Leftrightarrow x^T A x \geq 0$ for all x .
- ▶ $A \preceq B \Leftrightarrow B - A \succeq 0$ (matrix Loewner order).

Positive Semidefinite (PSD) and Notation $A \succeq B$

- ▶ $A \succeq 0$ (PSD) $\Leftrightarrow x^T A x \geq 0$ for all $x \in \mathbb{R}^n$.
- ▶ $A \succ 0$ (PD) $\Leftrightarrow x^T A x > 0$ for all nonzero x .
- ▶ $A \succeq B$ means $A - B \succeq 0$.

Positive Semidefinite (PSD) and Notation $A \succeq B$

- ▶ $A \succeq 0$ (PSD) $\Leftrightarrow x^T A x \geq 0$ for all $x \in \mathbb{R}^n$.
- ▶ $A \succ 0$ (PD) $\Leftrightarrow x^T A x > 0$ for all nonzero x .
- ▶ $A \succeq B$ means $A - B \succeq 0$.

For symmetric A :

$$A \succeq 0 \Leftrightarrow \lambda_i(A) \geq 0 \ \forall i \Leftrightarrow A = Q \Lambda Q^T, \ \Lambda \succeq 0.$$

Singular Value Decomposition (SVD)

For any $A \in \mathbb{R}^{m \times n}$,

$$A = U\Sigma V^T,$$

where $U \in \mathbb{R}^{m \times m}$ and $V \in \mathbb{R}^{n \times n}$ are orthogonal, and $\Sigma \in \mathbb{R}^{m \times n}$ is diagonal with nonnegative entries $\sigma_1 \geq \cdots \geq \sigma_r > 0$ (singular values).

Singular Value Decomposition (SVD)

For any $A \in \mathbb{R}^{m \times n}$,

$$A = U \Sigma V^T,$$

where $U \in \mathbb{R}^{m \times m}$ and $V \in \mathbb{R}^{n \times n}$ are orthogonal, and $\Sigma \in \mathbb{R}^{m \times n}$ is diagonal with nonnegative entries $\sigma_1 \geq \dots \geq \sigma_r > 0$ (singular values). Useful facts:

- ▶ $\sigma_i(A) = \sqrt{\lambda_i(A^T A)}$.
- ▶ $\|A\|_2 = \sigma_{\max}(A)$, $\|A\|_F^2 = \sum_i \sigma_i^2$.
- ▶ Best rank- k approximation (Eckart–Young):
 $A_k = U_{:,1:k} \Sigma_{1:k,1:k} V_{:,1:k}^T$ minimizes $\|A - X\|_F$ over $\text{rank}(X) \leq k$.

Common Inequalities

- ▶ Cauchy–Schwarz: $|x^T y| \leq \|x\|_2 \|y\|_2$.
- ▶ For PSD A : $|x^T A y| \leq \sqrt{x^T A x} \sqrt{y^T A y}$.
- ▶ Submultiplicativity: $\|Ax\|_2 \leq \|A\|_2 \|x\|_2$.
- ▶ For symmetric A : $\lambda_{\min}(A) \|x\|_2^2 \leq x^T A x \leq \lambda_{\max}(A) \|x\|_2^2$.

Use Hessian Matrix: Example 1

Prove that $f(x, y) = xy + a(x^2 + y^2)$ is convex if and only if $a \geq 1/2$.

Use Hessian Matrix: Example 1

Prove that $f(x, y) = xy + a(x^2 + y^2)$ is convex if and only if $a \geq 1/2$.

$$\nabla_f^2(x, y) = \begin{bmatrix} 2a & 1 \\ 1 & 2a \end{bmatrix}.$$

Use Hessian Matrix: Example 1

Prove that $f(x, y) = xy + a(x^2 + y^2)$ is convex if and only if $a \geq 1/2$.

$$\nabla_f^2(x, y) = \begin{bmatrix} 2a & 1 \\ 1 & 2a \end{bmatrix}.$$

Let λ be its eigenvalue.

Use Hessian Matrix: Example 1

Prove that $f(x, y) = xy + a(x^2 + y^2)$ is convex if and only if $a \geq 1/2$.

$$\nabla_f^2(x, y) = \begin{bmatrix} 2a & 1 \\ 1 & 2a \end{bmatrix}.$$

Let λ be its eigenvalue.

$$0 = \begin{vmatrix} 2a - \lambda & 1 \\ 1 & 2a - \lambda \end{vmatrix} = (2a - \lambda)^2 - 1.$$

Use Hessian Matrix: Example 1

Prove that $f(x, y) = xy + a(x^2 + y^2)$ is convex if and only if $a \geq 1/2$.

$$\nabla_f^2(x, y) = \begin{bmatrix} 2a & 1 \\ 1 & 2a \end{bmatrix}.$$

Let λ be its eigenvalue.

$$0 = \begin{vmatrix} 2a - \lambda & 1 \\ 1 & 2a - \lambda \end{vmatrix} = (2a - \lambda)^2 - 1.$$

$$\lambda = 2a - 1, 2a + 1.$$

Use Hessian Matrix: Example 1

Prove that $f(x, y) = xy + a(x^2 + y^2)$ is convex if and only if $a \geq 1/2$.

$$\nabla_f^2(x, y) = \begin{bmatrix} 2a & 1 \\ 1 & 2a \end{bmatrix}.$$

Let λ be its eigenvalue.

$$0 = \begin{vmatrix} 2a - \lambda & 1 \\ 1 & 2a - \lambda \end{vmatrix} = (2a - \lambda)^2 - 1.$$

$$\lambda = 2a - 1, 2a + 1.$$

Both λ 's are non-negative iff $a \geq 1/2$.

Use Hessian Matrix: Example 2

Prove that $f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

Use Hessian Matrix: Example 2

Prove that $f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

$$\nabla f(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} \\ \frac{-2e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} \end{bmatrix}^T.$$

Use Hessian Matrix: Example 2

Prove that $f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

$$\nabla f(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} & -\frac{2e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} \end{bmatrix}^T.$$

Use Hessian Matrix: Example 2

Prove that $f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

$$\nabla f(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} & -\frac{2e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} \end{bmatrix}^T.$$

$$\nabla_f^2(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} & & \\ & & \end{bmatrix}$$

Use Hessian Matrix: Example 2

Prove that $f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

$$\nabla f(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} & -\frac{2e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} \end{bmatrix}^T.$$

$$\nabla_f^2(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} & -\frac{2e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} \\ -\frac{2e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} & \frac{4e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} \end{bmatrix}$$

Use Hessian Matrix: Example 2

Prove that $f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

$$\nabla f(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} & -\frac{2e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} \end{bmatrix}^T.$$

$$\nabla_f^2(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} & -\frac{2e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} \\ -\frac{2e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} & \frac{4e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} \end{bmatrix}$$

Use Hessian Matrix: Example 2

Prove that $f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

$$\nabla f(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} & -\frac{2e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} \end{bmatrix}^T.$$

$$\nabla_f^2(\beta) = \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} & -\frac{2e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} \\ -\frac{2e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} & \frac{4e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} \end{bmatrix} \propto +$$
$$\begin{bmatrix} 1 & -2 \\ -2 & 4 \end{bmatrix}.$$

Use Hessian Matrix: Example 2

Prove that $f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

$$\begin{aligned}\nabla f(\beta) &= \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} & -\frac{2e^{\beta_1 - 2\beta_2}}{1 + e^{\beta_1 - 2\beta_2}} \end{bmatrix}^T. \\ \nabla_f^2(\beta) &= \begin{bmatrix} \frac{e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} & -\frac{2e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} \\ -\frac{2e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} & \frac{4e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} \end{bmatrix} \propto_+ \\ &\begin{bmatrix} 1 & -2 \\ -2 & 4 \end{bmatrix}.\end{aligned}$$

The eigenvalues are 0 and $\frac{5e^{\beta_1 - 2\beta_2}}{(1 + e^{\beta_1 - 2\beta_2})^2} \geq 0$.

$\nabla_f^2(\beta) \succeq 0$. f is convex.

Use Hessian Matrix: Example 2

$f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

Extension. Is $f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta})$ always convex for any constant vector \mathbf{c} ?

Use Hessian Matrix: Example 2

$f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

Extension. Is $f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta})$ always convex for any constant vector \mathbf{c} ?

$$\nabla_f^2(\beta) = \frac{e^{\mathbf{c}^\top \beta}}{(1 + e^{\mathbf{c}^\top \beta})^2} \begin{bmatrix} c_1^2 & c_1 c_2 & \cdots & c_1 c_n \\ c_1 c_2 & c_2^2 & \cdots & c_2 c_n \\ \vdots & \vdots & \ddots & \vdots \\ c_1 c_n & c_2 c_n & \cdots & c_n^2 \end{bmatrix} \propto_+ \mathbf{c} \mathbf{c}^\top.$$

Use Hessian Matrix: Example 2

$f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

Extension. Is $f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta})$ always convex for any constant vector \mathbf{c} ?

$$\nabla_f^2(\beta) = \frac{e^{\mathbf{c}^\top \beta}}{(1 + e^{\mathbf{c}^\top \beta})^2} \begin{bmatrix} c_1^2 & c_1 c_2 & \cdots & c_1 c_n \\ c_1 c_2 & c_2^2 & \cdots & c_2 c_n \\ \vdots & \vdots & \ddots & \vdots \\ c_1 c_n & c_2 c_n & \cdots & c_n^2 \end{bmatrix} \propto_+ \mathbf{c} \mathbf{c}^\top.$$

Why is $\nabla_f^2(\beta)$ always positive semi-definite?

Use Hessian Matrix: Example 2

$f(\beta_1, \beta_2) = \log(1 + e^{\beta_1 - 2\beta_2})$ is convex over \mathbb{R}^2 .

Extension. Is $f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta})$ always convex for any constant vector \mathbf{c} ?

$$\nabla_f^2(\beta) = \frac{e^{\mathbf{c}^\top \beta}}{(1 + e^{\mathbf{c}^\top \beta})^2} \begin{bmatrix} c_1^2 & c_1 c_2 & \cdots & c_1 c_n \\ c_1 c_2 & c_2^2 & \cdots & c_2 c_n \\ \vdots & \vdots & \ddots & \vdots \\ c_1 c_n & c_2 c_n & \cdots & c_n^2 \end{bmatrix} \propto_+ \mathbf{c} \mathbf{c}^\top.$$

Why is $\nabla_f^2(\beta)$ always positive semi-definite?

Because for every $\mathbf{x}, \beta \in \mathbb{R}^n$,

$$\mathbf{x}^\top \nabla_f^2(\beta) \mathbf{x} = \frac{e^{\mathbf{c}^\top \beta}}{(1 + e^{\mathbf{c}^\top \beta})^2} \mathbf{x}^\top \mathbf{c} \mathbf{c}^\top \mathbf{x} = \frac{e^{\mathbf{c}^\top \beta}}{(1 + e^{\mathbf{c}^\top \beta})^2} (\mathbf{x}^\top \mathbf{c})^2 \geq 0.$$

Use Hessian Matrix: Example 2

$f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta})$ is always convex over \mathbb{R}^n .

Extension II. Is $f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta}) - \mathbf{m}^\top \beta + \text{Const}$ always convex for any constant vector \mathbf{c}, \mathbf{m} ?

Use Hessian Matrix: Example 2

$f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta})$ is always convex over \mathbb{R}^n .

Extension II. Is $f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta}) - \mathbf{m}^\top \beta + \text{Const}$ always convex for any constant vector \mathbf{c}, \mathbf{m} ?

Application: Logistic Regression. $X_i \in \mathbb{R}^n, y_i \in \{0, 1\}$ are given data.

Use Hessian Matrix: Example 2

$f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta})$ is always convex over \mathbb{R}^n .

Extension II. Is $f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta}) - \mathbf{m}^\top \beta + \text{Const}$ always convex for any constant vector \mathbf{c}, \mathbf{m} ?

Application: Logistic Regression. $X_i \in \mathbb{R}^n, y_i \in \{0, 1\}$ are given data.

We want a good β for predicting y with $\hat{y} \in [0, 1]$ given another X .

Use Hessian Matrix: Example 2

$f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta})$ is always convex over \mathbb{R}^n .

Extension II. Is $f(\beta) = \log(1 + e^{\mathbf{c}^\top \beta}) - \mathbf{m}^\top \beta + \text{Const}$ always convex for any constant vector \mathbf{c}, \mathbf{m} ?

Application: Logistic Regression. $X_i \in \mathbb{R}^n, y_i \in \{0, 1\}$ are given data.

We want a good β for predicting y with $\hat{y} \in [0, 1]$ given another X .

With a good β^* , we predict that

$$\hat{y} = \frac{1}{1 + e^{-X^\top \beta^*}}.$$

Use Hessian Matrix: Example 2

$f(\beta) = \log(1 + e^{c^T \beta})$ is always convex over \mathbb{R}^n .

Extension II. Is $f(\beta) = \log(1 + e^{c^T \beta}) - \mathbf{m}^T \beta + \text{Const}$ always convex for any constant vector \mathbf{c}, \mathbf{m} ?

Application: Logistic Regression. $X_i \in \mathbb{R}^n, y_i \in \{0, 1\}$ are given data.

We want a good β for predicting y with $\hat{y} \in [0, 1]$ given another X .

With a good β^* , we predict that

$$\hat{y} = \frac{1}{1 + e^{-X^T \beta^*}}.$$

Maximum Likelihood Estimation (MLE) gives that