
Three dimensional coordinates into two dimensional coordinates transformation.
Overworked english edition. Almost completely rewritten chapters from top to bottom.

Three dimensional coordinates into two dimensional coordinates transformation

Edward Gerhold

September 2, 2015

Version 0.4.0-under-construction
old file: gerhold-3dto2d-0.3.99-with-mistakes.pdf
german draft: gerhold-3dto2d-de.pdf

Remark. This is a development version. With proper mistakes. The coordinate system itself is easy, useful and a working tool.

Contents

1	Introduction	4
2	Designing a 2x3 coordinate system	4
2.1	Indizes and notations in this document	4
2.2	Axis angles	4
2.3	Axis units	4
2.4	Axis vectors	4
2.5	Time to show the operation - sums of three scaled 2-D position vectors.	4
2.6	Some topology	4
3	Transformation tools	4
3.1	In place of a 3x3 basis	4
3.2	The functional $f(x)$	4
3.2.1	Natural $\langle f, x \rangle$	4
3.2.2	Examples for $f \circ g$	4
3.2.3	Derivative	4
3.2.4	Integral of the derivative	4
3.2.5	Integrals of $f(x)$	4
3.3	The matrix	4
3.3.1	Perfect 2x3 coordinate system	4
3.3.2	Examples of 2x3 coordinate systems	4
3.3.3	∇f^T Jacobi matrix of $f(x)$	4
3.3.4	Singular value decomposition	4
3.4	Projecting z onto a vector	4
4	Transformation behaviour	4
4.1	Linear transformation	4
4.2	Linear map, bounds on the operator	4
4.3	Sequences and Convergence	4
5	Computer Implementation	4
5.1	Generic Computer Code	4
5.2	EcmaScript 6 Example Code	4
6	Basis vs. Linear Combination	4
6.1	Orthogonality	4
6.1.1	Orthogonal Projection	4
6.2	Gram-Schmidt applied onto 2x3	4

7 Corollary	4
7.1 Transforming four dimensions with 2x4	4
7.2 3-D curve of a complex sequence, family surfaces	4
7.3 My plane in front of the drawn side of the space	4
7.4 $\ker(A)$, $f=0$ points are on the line into and through the origin \perp to the planes origin . . .	4
7.5 Covered points on a line normal to the front of the drawn	4
8 Summary	4
A Vector spaces	4
A.1 Basic tools	4
B To be continued	4
B.1 Measuring projected objects	4

1 Introduction

2 Designing a 2x3 coordinate system

2.1 Indices and notations in this document

2.2 Axis angles

2.3 Axis units

2.4 Axis vectors

2.5 Time to show the operation - sums of three scaled 2-D position vectors.

2.6 Some topology

3 Transformation tools

3.1 In place of a 3x3 basis

3.2 The functional $f(x)$

3.2.1 Natural $\langle f, x \rangle$

3.2.2 Examples for $f \circ g$

3.2.3 Derivative

3.2.4 Integral of the derivative

3.2.5 Integrals of $f(x)$

3.3 The matrix

3.3.1 Perfect 2x3 coordinate system

3.3.2 Examples of 2x3 coordinate systems

3.3.3 ∇f^T Jacobi matrix of $f(x)$

3.3.4 Singular value decomposition

3.4 Projecting z onto a vector

4 Transformation behaviour

4.1 Linear transformation

4.2 Linear map, bounds on the operator

4.3 Sequences and Convergence

5 Computer Implementation

5.1 Generic Computer Code

5.2 EcmaScript 6 Example Code

6 Basis vs. Linear Combination

6.1 Orthogonality

6.1.1 Orthogonal Projection

6.2 Gram-Schmidt applied onto 2x3

7 Corollary

7.1 Transforming four dimensions with 2x4

7.2 3-D curve of a complex sequence, family surfaces

7.3 My plane in front of the drawn side of the space