

COMP-2310 Formula Sheet

Midterm Test 2

University of Windsor
Edward Nafornta

November 7, 2022

Chapter 3

Definition (Principle of Extension). Two sets A and B are **equal**, denoted by $A = B$, iff $(\forall x)(x \in A \Leftrightarrow x \in B)$

Lemma (3.2.1). Let A, B, C be sets.

(i) $A = A$

(ii) If $(A = B)$, then $B = A$

(iii) If $(A = B)$ and $(B = C)$, then $A = C$

Lemma (3.2.2). $A \neq B$ iff $(\exists x)(x \in A \wedge x \notin B) \vee (\exists x)(x \notin A \wedge x \in B)$

Corollary (3.2.2.1). Let A and B be sets.

$$(\exists x)(x \in A \wedge x \notin B) \Rightarrow A \neq B$$

Definition (Subset). Let A, B be two sets. A is a **subset** of B or B **includes** A , denoted by $A \subseteq B$, iff $(\forall x)(x \in A \Rightarrow x \in B)$.

A is a **proper subset** of B , denoted by $A \subset B$, iff $A \subseteq B$ and $A \neq B$.

Lemma (3.2.3). Let A, B, C be sets.

(i) $A \subseteq A$

(ii) If $A \subseteq B$ and $B \subseteq A$, then $A = B$

(iii) If $A \subseteq B$ and $B \subseteq C$, then $A \subseteq C$

Lemma (3.2.4). Let A, B be sets. If $A = B$, then $(A \subseteq B) \wedge (B \subseteq A)$

Lemma (3.2.5). $A \not\subseteq B \Leftrightarrow (\exists x)(x \in A \wedge x \notin B)$

Lemma (3.2.6). If $A \subset B$, then $(\exists x)(x \in B \wedge x \notin A)$

Definition (Principle of Specification). For every set A and every formula $S(x)$, there exists a set B whose elements are exactly those elements of A for which $S(x)$ is true. The set B is denoted by: $\{x|x \in A \wedge S(x)\}$ or $\{x \in A|S(x)\}$

Definition (Power Set). Let A be a set. The **power set** of A , denoted by $\mathcal{P}(A)$, is the set $\{X|X \subseteq A\}$

Chapter 4

Definition (Union). Let A, B be two sets. The **union** of A and B is the set $A \cup B = \{x | x \in A \vee x \in B\}$

Theorem (4.1.1). Let A, B be sets.

- (i) $A \cup \emptyset = A$
- (ii) $A \cup A = A$
- (iii) $A \cup B = B \cup A$
- (iv) $(A \cup B) \cup C = A \cup (B \cup C)$
- (v) $A \subseteq B$ iff $A \cup B = B$

Definition (Intersection). Let A, B be two sets. The **intersection** of A and B is the set $A \cap B = \{x | x \in A \wedge x \in B\}$

Theorem (4.2.2). Let A, B be sets.

- (i) $A \cap \emptyset = \emptyset$
- (ii) $A \cap A = A$
- (iii) $A \cap B = B \cap A$
- (iv) $(A \cap B) \cap C = A \cap (B \cap C)$
- (v) $A \subseteq B$ iff $A \cap B = A$

Corollary (4.2.2.1). Let A, B be any two sets. Then, $A \cap B \subseteq A \subseteq A \cup B$

Theorem (4.2.3). Let A, B, C be sets.

- (i) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
- (ii) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

Definition (Disjoint). Two sets A and B are **disjoint** if $A \cap B = \emptyset$

Definition (Relative Complement). Let A, B be two sets. The **relative complement** of B in A is the set $A - B = \{x | x \in A \wedge x \notin B\}$

Theorem (4.3.4). *Let A, B, C be sets.*

$$(i) \ A \subseteq B \text{ iff } A - B = \emptyset$$

$$(ii) \ A - (A - B) = A \cap B$$

$$(iii) \ A \cap (B - C) = (A \cap B) - (A \cap C)$$

Definition (Absolute Complement). *The **(relative) Universal set**, denoted by \mathbf{U} , is a set which contains every object (thereby includes every set) in our discussion. Let A be any set such that $A \subseteq \mathbf{U}$. The **absolute complement** of A is the set $\overline{A} = \mathbf{U} - A$*

[4.3.5]

Lemma. *Let $A \subseteq \mathbf{U}$. Then, $x \in \overline{A} \Leftrightarrow x \notin A$*

Theorem (4.3.6 DeMorgan's Theorem). *Let A, B be any two subsets of \mathbf{U}*

$$(i) \ \overline{A \cup B} = \overline{A} \cap \overline{B}$$

$$(ii) \ \overline{A \cap B} = \overline{A} \cup \overline{B}$$

Theorem (4.3.7). *Let A, B be any two subsets of \mathbf{U} .*

$$(i) \ \overline{\overline{A}} = A$$

$$(ii) \ \overline{\emptyset} = \mathbf{U} \text{ and } \overline{\mathbf{U}} = \emptyset$$

$$(iii) \ A \cap \overline{A} = \emptyset \text{ and } A \cup \overline{A} = \mathbf{U}$$

$$(iv) \ A \subseteq B \text{ iff } \overline{B} \subseteq \overline{A}$$

$$(v) \ A - B = A \cap \overline{B}$$

Definition (Ordered Pair). *The **ordered pair** of a and b is the set*

$$(a, b) = \{\{a\}, \{a, b\}\} = \{x | x = \{a\} \vee x = \{a, b\}\}$$

Lemma (4.4.8). *$(a, b) = (x, y)$ implies that $a = x$ and $b = y$*

Lemma (4.4.9). *$a = c \wedge b = d \Rightarrow (a, b) = (c, d)$*

Definition (Cartesian Product). *Let A, B be two sets. The **Cartesian Product** of A and B is the set $A \times B = \{x | (\exists a)(\exists b)(a \in A \wedge b \in B \wedge x = (a, b))\} = \{(a, b) | a \in A \wedge b \in B\}$*

Lemma (4.5.10). *Let A , B , X and Y be sets.*

$$(i) \quad (A \cup B) \times X = (A \times X) \cup (B \times X)$$

$$(ii) \quad (A \cap B) \times X = (A \times X) \cap (B \times X)$$

$$(iii) \quad (A - B) \times X = (A \times X) - (B \times X)$$

$$(iv) \quad (A \cap B) \times (X \cap Y) = (A \times X) \cap (B \times Y)$$

$$(v) \quad (A = \emptyset \vee B = \emptyset) \Leftrightarrow A \times B = \emptyset$$

Definition (Union of a Collection of Sets). *Let \mathcal{C} be a collection of sets. The **union of \mathcal{C}** is the set:*

$$\bigcup_{X \in \mathcal{C}} X = \{x | (\exists X)(X \in \mathcal{C} \wedge x \in X)\} = \{x | x \in X \text{ for some } X \in \mathcal{C}\}$$

Definition (Intersection of a Collection of Sets). *Let \mathcal{C} be a collection of sets such that $\mathcal{C} \neq \emptyset$. The **intersection of \mathcal{C}** is the set:*

$$\bigcap_{X \in \mathcal{C}} X = \{x | (\forall X)(X \in \mathcal{C} \Rightarrow x \in X)\} = \{x | x \in X \text{ for all } X \in \mathcal{C}\}$$

Lemma. $A: (X \cap Y) \cup X = X$

Chapter 5

Definition (Relation). A **relation** is a set of ordered pairs. Specifically, a set R is a relation if $(\forall x)(x \in R \Rightarrow (\exists a)(\exists b)x = (a, b))$

Definition (Domain & Range of Relations). Let R be a relation. The **domain** of R is the set $\text{Dom}(R) = \{x | (\exists y)(x, y) \in R\}$ and the **range** of R is the set $\text{Ran}(R) = \{y | (\exists x)(x, y) \in R\}$

Definition (Cartesian Product of Relations). Let $R \subseteq X \times Y$, we say that R is a **relation from X to Y** . In particular, when $X = Y$, (ie. $R \subseteq X \times X$), we say that R is a **relation in X** .

Lemma (5.1.1). Let R be a relation from X to Y . Then $\text{Dom}(R) \subseteq X$ and $\text{Ran}(R) \subseteq Y$

Definition (Properties of Relation). Let R be a relation in a set X

R is **reflexive** if $(\forall x \in X)(x, x) \in R$;

R is **irreflexive** if $(\forall x \in X)(x, x) \notin R$;

R is **symmetric** if $(x, y) \in R \Rightarrow (y, x) \in R$;

R is **antisymmetric** if $(x, y) \in R \wedge (y, x) \in R \Rightarrow x = y$;

R is **asymmetric** if $\forall x, y \in X, ((x, y) \in R \wedge (y, x) \in R)$, or equivalently,
 $\forall x, y \in X, (x, y) \in R \Rightarrow (y, x) \notin R$;

R is **transitive** if $(x, y) \in R \wedge (y, z) \in R \Rightarrow (x, z) \in R$.

Lemma (5.2.1). Let R be a relation in X . If $(\exists x \in X)(x, x) \in R$, then R is not asymmetric.

Definition (Equivalence Relation). A relation R in a set X is an **equivalence relation** if R is reflexive, symmetric and transitive.

Definition (Equivalence Class). Let R be an equivalence relation in a set X . For each $x \in X$, the **equivalence class of x w.r.t. R** is the set

$[x] / R = \{y | y \in X \wedge (x, y) \in R\} = \{y \in X | (x, y) \in R\}$ or equivalently,

$[x] / R = \{y | (x, y) \in R\}$

Definition (Collection of Equivalence Classes). Let R be an equivalence relation in a set X . **The collection of equivalence classes w.r.t. R** is the set: $[X] / R = \{S | (\exists x)(x \in X \wedge S = [x] / R)\} = \{[x] / R | x \in X\}$

Definition (Partition of a Collection of Sets).
continue on page 122 definition at the very top!