

# Math-1730 Formula Sheet

Edward Nafornta

Evan Petrimoulx

April 24, 2022

## **1   Indefinite Integrals:**

**Constant Integration:**

$$\int a \, dx = x + C$$

$$\int a f(x) \, dx = f(x) + C$$

**Distribution:**

$$\int f(x) \pm g(x) \, dx = \int f(x) \, dx \pm \int g(x) \, dx$$

**Exponentials:**

$$\int x^n \, dx = \frac{x^n + 1}{n + 1} + C, \quad \text{if } n \neq -1$$

$$\int e^x \, dx = e^x + C$$

$$\int a^x \, dx = \frac{a^x}{\ln a} + C$$

$$\int \frac{1}{x} \, dx = \ln |x| + C$$

### Trigonometric:

$$\int \sin x \, dx = -\cos x + C$$

$$\int \sec^2 x \, dx = \tan x + C$$

$$\int \sec x \tan x \, dx = \sec x + C$$

$$\int \tan x \, dx = \ln |\sec x| + C$$

$$\int \sec x \, dx = \ln |\sec x + \tan x| + C$$

$$\int \frac{1}{x^2 + 1} \, dx = \arctan x + C$$

$$\int \cos x \, dx = \sin x + C$$

$$\int \csc^2 x \, dx = -\cot x + C$$

$$\int \csc x \cot x \, dx = -\csc x + C$$

$$\int \cot x \, dx = \ln |\sin x| + C$$

$$\int \csc x \, dx = \ln |\csc x - \cot x| + C$$

$$\int \frac{1}{\sqrt{1-x^2}} = \arcsin x + C$$

Generalization:

$$\int \frac{1}{x^2 + a^2} \, dx = \frac{1}{a} \arctan \frac{x}{a} + C, \quad \text{if } a \neq 0$$

Generalization:

$$\int \frac{1}{\sqrt{a^2 - x^2}} \, dx = \arcsin \frac{x}{a} + C, \quad \text{if } a \neq 0$$

## Integration By Parts:

Formula:

$$\int f(x)g'(x) \, dx = f(x)g(x) - \int f'(x)g(x) \, dx \quad (1)$$

Examples:

$$\begin{aligned} \int x \cos x \, dx &= \int x(\sin x)' \, dx \\ &= x(\sin x) - \int (x)' \sin x \, dx \\ &= x(\sin x) - \int \sin x \, dx \\ &= x \sin x + \cos x + C \end{aligned}$$

$$\begin{aligned}
\int 2^x x \, dx &= \int x \left( \frac{2^x}{\ln 2} \right)' dx \\
&= x \frac{2^x}{\ln 2} - \int (x)' \left( \frac{2^x}{\ln 2} \right) dx \\
&= x \frac{2^x}{\ln 2} - \int \frac{2^x}{\ln 2} dx \\
&= x \frac{2^x}{\ln 2} - \frac{1}{\ln 2} \int 2^x dx \\
&= x \frac{2^x}{\ln 2} - \frac{1}{\ln 2} \cdot \frac{2^x}{\ln 2} + C \\
&= x \frac{2^x}{\ln 2} - \frac{2^x}{\ln^2(2)} + C
\end{aligned}$$

## U-Substitution:

Formula:

$$u = g(x) \text{ then,} \tag{2}$$

$$\int f(g(x))g'(x) dx = \int f(u) du \tag{3}$$

When using U-Substitution, normally pick the most complex part of the problem then calculate "du=g'(x)" and algebraically solve for 'dx', or until everything is in terms of 'u' and 'du'.

Examples:

$$\int x^2 \cos(x^3) dx$$

$$\text{let } u = x^3 \text{ then, } du = 3x^2 dx$$

$$\int x^2 \cos(x^3) dx \quad \text{Sub : } \cos(x^3) = \cos(u)$$

$$\int \cos(x^3)x^2 dx = \int \cos(u) \cdot \left(\frac{1}{3} du\right)$$

$$= \int \frac{1}{3} \cos(u) du$$

$$= \frac{1}{3} \int \cos(u) dx$$

$$= \frac{1}{3} \cdot \sin(u) + C$$

$$= \frac{\sin(x^3)}{3} + C$$

# Trigonometric Integrals:

## Formulas:

### 1. Integrals of the form:

$$\int \sin^m(x) \cos^n(x) dx$$

### Three Cases:

1. If  $n$  is odd, save a factor of  $\cos(x)$  and use  $\cos^2(x) = 1 - \sin^2(x)$  to get everything in terms of  $\sin(x)$ .
2. If  $m$  is odd, save one factor of  $\sin(x)$  and use  $\sin^2(x) = 1 - \cos^2(x)$  to get everything in terms of  $\cos(x)$ .
3. If  $m, n$  are both even use half-angle formulas:  $\sin^2(x) = \frac{1 - \cos(2x)}{2}$  :  $\cos^2(x) = \frac{1 + \cos(2x)}{2}$ .

### 2. Integrals of the form:

$$\int \tan^m(x) \sec^n(x) dx$$

### Two Cases:

1. If  $n$  is even, save a factor of  $\sec^2(x)$  and use  $\sec^2(x) = 1 + \tan^2(x)$  to get everything in terms of  $\tan(x)$ .
2. If  $m$  is odd, save a factor of  $\sec(x) \tan(x)$  and use  $\tan^2(x) = \sec^2(x) - 1$  to get everything in terms of  $\sec(x)$ .

### 3. Integrals of the form:

1.  $\sin(mx) \cos(nx) dx$   
Use identity:  $\sin(A) \cos(B) = \frac{1}{2}(\sin(A - B) + \sin(A + B))$
2.  $\sin(mx) \sin(nx) dx$   
Use identity:  $\frac{1}{2}(\cos(A - B) - \cos(A + B))$
3.  $\cos(mx) \cos(nx) dx$   
Use identity:  $\frac{1}{2}(\cos(A - B) + \cos(A + B))$

## Trigonometric Substitution:

Idea of Trigonometric Substitution:

Use the 'reverse' substitution rule in the form  $x = ab(c)$ , where 'a' is a suitable constant, and 'b(c)' is a suitable trigonometric function, to boil down the integral to some sort of trigonometric integral.

If You See	Try Subbing	This Identity will Help
$\sqrt{a^2 - x^2}$	$x = a \sin(\theta)$ $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$	$1 - \sin^2(\theta) = \cos^2(\theta)$
$\sqrt{a^2 + x^2}$	$x = a \tan(\theta)$ $-\frac{\pi}{2} < \theta < \frac{\pi}{2}$	$1 + \tan^2(\theta) = \sec^2(\theta)$
$\sqrt{x^2 - a^2}$	$x = a \sec(\theta)$ $0 \leq \theta < \frac{\pi}{2}$ or $\pi \leq \theta < \frac{3\pi}{2}$	$\sec^2(\theta) - 1 = \tan^2(\theta)$

### Examples:


1.  $\int x\sqrt{16 - x^2} dx$

Let  $x = 4 \sin(\theta)$ , where  $-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}$ , and  $dx = 4 \cos(\theta) d\theta$ .

$$\begin{aligned}
 \sqrt{16 - x^2} &= \sqrt{16 - (4 \sin(\theta))^2} \\
 &= \sqrt{16 - 16 \sin^2(\theta)} \\
 &= \sqrt{16(1 - \sin^2(\theta))} \\
 &= \sqrt{16(\cos^2(\theta))} \\
 &= |4 \cos(\theta)| \\
 &= 4 \cos(\theta) \text{ because } 4 \cos(\theta) \geq 0 \text{ for } -\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2}.
 \end{aligned}$$

$$\begin{aligned}
 \int x\sqrt{16 - x^2} dx &= \int (4 \sin(\theta) \cdot (4 \cos(\theta)) \cdot (4 \cos(\theta)) d\theta \\
 &= 64 \int \sin(\theta) \cos^2(\theta) d\theta
 \end{aligned}$$

$$\begin{aligned}
 \text{Let } u &= \cos(\theta), \quad du = -\sin(\theta) d\theta. \\
 &= -64 \int u^2 du \\
 &= -64 \cdot \frac{u^3}{3} + C \\
 &= -\frac{64}{3} \cos^3(\theta) + C
 \end{aligned}$$



By Pythagorean Theorem:  $\cos(\theta) = \frac{\sqrt{16 - x^2}}{4}$

$$\int x\sqrt{16 - x^2} dx = -\frac{64}{3} \cdot \frac{\sqrt{16 - x^2}^3}{4^3} + C$$

$$\int x\sqrt{16 - x^2} dx = -\frac{\sqrt{16 - x^2}^3}{3} + C$$

## Partial Fraction Decomposition:

Case 1:  $Q(x)$  is a product of distinct linear factors

$$Q(x) = (a_1x + b_1)(a_2x + b_2) \dots (a_kx + b_k)$$

The partial fraction decomposition of  $f(x)$  is  $\frac{P(x)}{Q(x)} = \frac{A_1}{a_1x+b_1} + \frac{A_2}{a_2x+b_2} + \dots + \frac{A_k}{a_kx+b_k}$ .

Case 2:  $Q(x)$  is a product of linear factors, some of which are repeated. If  $(a_ix + b_i)^r$  (where  $r \neq 1$ ) occurs in the factorization of  $Q(x)$  then instead of  $\frac{A_i}{a_ix+b_i}$  term, we will have the sum:  $(\frac{B_1}{a_ix+b_i} + \frac{B_2}{a_ix+b_i}^2 + \dots + \frac{B_i}{a_ix+b_i}^i) \in \mathbb{R}$ .

Case 3: If  $Q(x)$  has a non-repeated quadratic factor  $ax^2 + bx + c$ , then there is a  $\frac{Ax+B}{ax^2+bx+c}$  term.

Case 4: More generally, if  $Q(x)$  has a repeated quadratic factor  $ax^2 + bx + c$ , then you will see:  $\frac{A_1x+B_1}{ax^2+bx+c} + \frac{A_2x+B_2}{ax^2+bx+c}^2 + \dots + \frac{A_rx+B_r}{ax^2+bx+c}^r$

### Examples:

$$\int \frac{x}{x^2 + 4x + 3} dx$$

Factor  $x^2 + 4x + 3 = (x + 3)(x + 1)$

Then the form of the partial fraction decomposition is:

$$\frac{x}{(x+1)(x+3)} = \frac{A}{x+1} + \frac{B}{x+3}$$

Reduce the fractions:  $\frac{x}{(x+1)(x+3)} = \frac{A}{x+1} + \frac{B}{x+3} \mid \cdot (x+1)(x+3)$

$x = A(x+3) + B(x+1)$ , is true  $\forall x \in \mathbb{R}$ .

Let  $x = -3$

$$-3 = A(-3+3) + B(-3+1)$$

$$-3 = -2B$$

$$B = \frac{3}{2}$$

Let  $x = -1$

$$-1 = A(-1+3) + B(-1+1)$$

$$-1 = 2A$$

$$A = -\frac{1}{2}$$

$$\begin{aligned} \int \frac{x}{x^2+4x+3} dx &= \int \left( \frac{-\frac{1}{2}}{x+1} + \frac{\frac{3}{2}}{x+3} \right) dx \\ &= -\frac{1}{2} \ln |x+1| + \frac{3}{2} \ln |x+3| + C \end{aligned}$$

## 2 Definite Integrals:

**Formula:**

$$\int_a^b f(x) dx = F(b) - F(a) \quad (4)$$

**Examples:**


$$\int_0^1 x^2 dx = \left. \frac{x^3}{3} \right|_0^1 = \frac{1^3}{3} - \frac{0^3}{3} = \frac{1}{3}$$

In Essence: Compute the indefinite integral first, then apply the definite integral formula.

### Properties of Definite Integrals

1.  $\int_a^a f(x) dx = 0$
2.  $\int_b^a f(x) dx = -\int_a^b f(x) dx$
3.  $\int_a^b f(x) dx + \int_b^c f(x) dx = \int_a^c f(x) dx$

### Improper Integrals



The area below this graph can be denoted as  $A(t) = \int_1^t \frac{1}{x^2} dx$ , and the solution would be:

$$\int_1^t \frac{1}{x^2} dx = \left. -\frac{1}{x} \right|_1^t = -\frac{1}{t} - \left(-\frac{1}{1}\right) = 1 - \frac{1}{t}$$

Therefore, to calculate this improper integral, we must take the limit and see where it approaches.

$$A = \lim_{t \rightarrow \infty} A(t) = \lim_{t \rightarrow \infty} \left(1 - \frac{1}{t}\right) = 1$$



### First Type of Improper Integrals:

- 1: If  $\int_a^t f(x) dx$  is defined for all  $t \geq a$ , define  
 $\int_a^\infty f(x) dx = \lim_{t \rightarrow \infty} \int_a^t f(x) dx$ , if the limit exists, the integral is convergent, otherwise, it's divergent.
- 2: If  $\int_t^b f(x) dx$  is defined for all  $t \leq b$ , define  
 $\int_{-\infty}^b f(x) dx = \lim_{t \rightarrow -\infty} \int_t^b f(x) dx$ .
- 3: If  $\int_a^\infty f(x) dx$  and  $\int_{-\infty}^a f(x) dx$  converge, let  
 $\int_{-\infty}^\infty f(x) dx = \int_a^\infty f(x) dx + \int_{-\infty}^a f(x) dx$

### Examples:

$$1. \int_1^\infty \frac{1}{x} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x} dx = \lim_{t \rightarrow \infty} (\ln |x| \Big|_1^t) = \lim_{t \rightarrow \infty} \ln(t) - \ln(1) = \lim_{t \rightarrow \infty} \ln(t) = +\infty$$

Since  $\int_1^\infty \frac{1}{x} dx$  is not finite (meaning doesn't resolve to an approachable number), therefore  $\int_1^\infty \frac{1}{x} dx$  diverges.

### Second Type of Improper Integrals:

- 1: If  $f(x)$  is continuous on  $[a, b)$  and discontinuous at  $b$ , define  
 $\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx$ .
- 2: If  $f(x)$  is continuous on  $(a, b]$  and discontinuous at  $a$ , define  
 $\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx$ .
- 3: If  $f(x)$  has a discontinuity at  $c \in (a, b)$  and  $\int_a^c f(x) dx$  and  $\int_c^b f(x) dx$  converges, define  
 $\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$ .

### Examples:

1.  $\int_0^1 \frac{1}{\sqrt{x}} dx$ , is an improper integral since it's discontinuous at  $x=0$ .


$$\int_0^1 \frac{1}{\sqrt{x}} dx = \lim_{t \rightarrow 0^+} \int_t^1 \frac{1}{\sqrt{x}} dx = \lim_{t \rightarrow 0^+} \left. \frac{x^{\frac{1}{2}}}{\frac{1}{2}} \right|_t^1 = \lim_{t \rightarrow 0^+} \left( \frac{1^{\frac{1}{2}}}{\frac{1}{2}} - \frac{t^{\frac{1}{2}}}{\frac{1}{2}} \right) = \frac{1}{\frac{1}{2}} = 2$$

Since  $\int_0^1 \frac{1}{\sqrt{x}} dx$  is finite (meaning it does resolve to an approachable number), therefore  $\int_0^1 \frac{1}{\sqrt{x}} dx$  converges.

## Areas Between Curves:

If  $f(x) \geq g(x) \forall x \in [a, b]$  then the area bounded by the curves;  $y=f(x)$ ,  $y=g(x)$ ,  $x=a$ , and  $x=b$  is  $A = \int_a^b (f(x) - g(x)) dx$ .

### Examples:



$$A = \int_{-1}^1 (f(x) - g(x)) dx$$

Let  $f(x)$  be the bigger function;  $f(x) = x^2 + 1$ , then  $g(x) = x$

$$A = \int_{-1}^1 (x^2 + 1 - x) dx = \left. \frac{x^3}{3} + x - \frac{x^2}{2} \right|_{-1}^1 = \frac{1}{3} + 1 - \frac{1}{2} - \left( -\frac{1}{3} - 1 + \frac{1}{2} \right) = \frac{2}{3} + 2 - 1 = \frac{5}{3}$$

### 3 Volumes Via Areas of Cross Sections:

If a solid lies between the planes  $x = a$  and  $x = b$  and  $A(x)$  is the area of the  $x$ -cross-section for  $a \leq x \leq b$ , then the volume of a solid is:

$$V = \int_a^b A(x) dx \quad (5)$$

#### Example - The Volume of a Sphere of Radius R:

$$R^2 = |x|^2 + r^2$$

Where  $R$  is the radius of the sphere,  $r$  is the length of the cross section, and  $|x|$  is the  $x$ -distance to the slice / cross section.

$$r = \sqrt{R^2 - x^2}$$

$$\begin{aligned} V &= \int_{-R}^R A(x) dx = \int_{-R}^R \pi(R^2 - x^2) dx \\ \pi R^2 x - \pi \frac{x^3}{3} \Big|_{-R}^R &= \left( \pi R^3 - \pi \frac{R^3}{3} \right) - \left( -\pi R^3 + \pi \frac{\pi R^3}{3} \right) \\ \frac{2\pi}{3} R^3 - \left( -\frac{2\pi}{3} R^3 \right) &= \frac{4\pi R^3}{3} \end{aligned}$$

#### Solids of Revolution

There are different formats for solids of revolution:

- Cylinders
- Cones
- Tori (sing. torus)

Each of which have two different methods on computing them:

- Disk/Washer Method
- Cylindrical Shells Method

#### Disk/Washer

- If the cross-section is a disk, then its area is

$$A(x) = \pi(\text{radius})^2 \quad (6)$$

- If the cross-section is a washer (annulus), then its area is

$$A(x) = \pi[(\text{outer radius})^2 - (\text{inner radius})^2] \quad (7)$$

## Cylindrical Shells

Let  $b > a \geq 0$ ,  $f(x) \geq 0$  on  $[a, b]$  and  $R$ , the region under the graph of  $y = f(x)$  between  $x = a$  and  $x = b$ . The volume of the solid of revolution is obtained by rotating  $R$  around the  $y$ -axis.

$$V = \int_a^b 2\pi x f(x) dx \quad (8)$$

It is important to note the Surface Area of a cylindrical shell:


$$2\pi hr$$

Idea: Slice the solid of revolution into cylindrical shells. You want to make the cylinder for every  $dx$  until you can approximate the area.

$$V = \int_a^b (\text{surface area of cylindrical shell}) dx$$

$$V = \int_a^b 2\pi (\text{height}) (\text{radius}) dx = \int_a^b 2\pi x f(x) dx$$

**Example:**



## 4 Physical Applications

### Work

If a constant force,  $F$ , acts on an object. The work done to move the object a distance,  $d$ , is:

$$W = F \cdot d \quad (9)$$

If the object moves from,  $x = a$ , to  $x = b$ , under the action of a variable force,  $F(x)$ , where  $(x)$  is the position, the work is:

$$W = \int_a^b F(x) dx \quad (10)$$

### Example

When a particle is  $x$  meters from the origin, a force of  $F(x) = 3x^2$  Newtons acts on it. How much work is done moving it from  $x = 2$ , to  $x = 4$ ?


$$W = \int_2^4 F(x) dx = \int_2^4 3x^2 dx = x^3 \Big|_2^4 = 4^3 - 2^3 = 56J$$

### Hooke's Law

The force required to maintain a spring stretched  $x$ -units beyond its natural length is  $F = k\Delta x$  where  $k$  is the spring constant.

### Example

A force of 60N is required to hold a spring stretched 5cm beyond its natural length. How much work is done in stretching the spring 5cm beyond its natural length (Starting from its original, natural length). From 5cm to 10cm beyond its natural length?



$$F = k\Delta x$$

$$60 = k(0.05m)$$

$$1200 = k$$

$$W = \int_0^{0.05} 1200x dx = 1200 \int_0^{0.05} x dx = \frac{1200x^2}{2} \Big|_0^{0.05} = 600(0.05)^2 = 1.5J$$

$$W = \int_0^{0.1} 1200x dx = 600x^2 \Big|_{0.05}^{0.1} = 600((0.1)^2 - (0.05)^2) = 4.5J$$

### Average Values of Functions:

#### Definition

The average value of a function on the interval  $[a, b]$  is:

$$f_{ave} = \frac{1}{b-a} \int_a^b f(x) dx$$

## Mean Value Theorem For Integrals

If  $f$  is continuous on  $[a, b]$ , then there exists some  $c \in [a, b]$  such that:

$$f(c) = f_{ave} = \frac{1}{b-a} \int_a^b f(x) dx$$

The fundamental theorem of calculus implies that the mean value theorem for Integrals and the mean value theorem for Derivatives are equivalent.

## Centers of Mass

The center of mass of  $n$  point masses  $m_1, m_2, m_3, \dots, m_n$  at the points  $(x_1, y_1), (x_2, y_2), (x_3, y_3) \dots (x_n, y_n)$  is  $(\bar{x}, \bar{y})$  where:

$$\bar{x} = \frac{\sum_i^n m_i x_i}{\sum_i^n m_i}$$

and

$$\bar{y} = \frac{\sum_i^n m_i y_i}{\sum_i^n m_i}$$

Let  $f(x) \geq g(x)$  and  $R$  be the region bounded by the graphs  $y = f(x)$  and  $y = g(x)$  between  $x = a$  and  $x = b$ . Then the center of mass (sometimes called the centroid) of  $R$  is the point  $(\bar{x}, \bar{y})$  where;

$$\bar{x} = \frac{1}{A} \int_a^b x(f(x) - g(x)) dx$$
$$\bar{y} = \frac{1}{2A} \int_a^b (f(x)^2 - g(x)^2) dx$$

And "A" is the area of  $R$

## Surface of Revolution

Translation: Surface Area of a Solid of Revolution

If the curve,  $y = f(x)$ ,  $a \leq x \leq b$  (with  $f'(x)$  continuous), is rotated around the x-axis, we obtain a surface of revolution whose surface area is:

$$A = \int_a^b 2\pi f(x) \sqrt{1 + f'(x)^2} dx \quad (11)$$

In Leibnitz notation:  $A = \int_a^b 2\pi f(x) \sqrt{1 + f'(x)^2} dx$

If  $x = g(y)$ ,  $c \leq y \leq d$ , then:  $A = \int_c^d 2\pi y \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$

If rotating around y-axis:  $A = \int_a^b 2\pi x \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx = \int_c^d 2\pi x \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$

## 5 Sequences

In this course we will only consider infinite sequences

*Notation:*

$$a_n = \frac{n}{2n-1}, \quad n \geq 3 \text{ or } \left\{ \frac{n}{2n-1} \right\}_{n=3}^{\infty}$$

There are various ways to describe a sequence:

- By formula ( $a_n = 2^n$  for  $n \geq 0$ )
- By a recurrence relation ( $a_0 = 1$  and  $a_n = 2a_{n-1}$  for  $n \geq 1$ )

### Limits of Sequences (Informal Definition)

The sequence of  $a_1, a_2, \dots, a_n$  has a Limit  $L$  if we can make  $a_n$  as close as we want to  $L$  by making  $n$  sufficiently large.

$$\lim_{n \rightarrow \infty} a_n = L \quad \text{or} \quad a_n \rightarrow L$$

The sequence  $a_n$  has limit  $L$  or the sequence  $a_n$  tends to / approaches / converges to  $L$ . If  $a_n$  has a finite limit, then it is called convergent. If it does not, it is referred to as divergent.

### Limits of Sequences (Rigorous)

The sequence  $a_1, a_2, a_3, \dots, a_n$  has limit  $L$  if any  $\epsilon > 0$ . There exists a positive integer  $N$  such that  $|a_n - L| < \epsilon \forall n \geq N$ .

There is a notation of infinite limits  $\lim_{n \rightarrow \infty} a_n = +\infty$  (respectively  $-\infty$ ) if we can make  $a_n$  as large (respectively as small) as we want by making  $n$  sufficiently large. The sequence  $a_1, a_2, \dots$  has limit  $+\infty$  if for any number  $M$ , there exists a positive integer  $N$  such that  $a_n > M \forall n \geq N$ .

### Limit Laws For Sequences

If  $a_n$  and  $b_n$  are convergent sequences, then:

$$\lim_{n \rightarrow \infty} (a_n + b_n) = \lim_{n \rightarrow \infty} a_n + \lim_{n \rightarrow \infty} b_n$$

$$\lim_{n \rightarrow \infty} (a_n - b_n) = \lim_{n \rightarrow \infty} a_n - \lim_{n \rightarrow \infty} b_n$$

$$\lim_{n \rightarrow \infty} ca_n = c \lim_{n \rightarrow \infty} a_n$$

$$\lim_{n \rightarrow \infty} (a_n b_n) = \lim_{n \rightarrow \infty} a_n \cdot \lim_{n \rightarrow \infty} b_n$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \frac{\lim_{n \rightarrow \infty} a_n}{\lim_{n \rightarrow \infty} b_n} \quad \text{if } \lim_{n \rightarrow \infty} b_n \neq 0$$

## Squeeze Theorem for Sequences

If  $a_n \leq b_n \leq c_n \forall n$  or  $n \geq n_0 \exists n_0$

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = L \implies \lim_{n \rightarrow \infty} b_n = L$$

If  $\lim_{n \rightarrow \infty} a_n = L$  and  $f$  is continuous at  $L$ , then:

$$\lim_{n \rightarrow \infty} f(a_n) = f(L)$$

If  $f$  is a function such that  $\lim_{x \rightarrow \infty} f(x) = L$  and  $a_n = f(n)$  then:

$$\lim_{n \rightarrow \infty} a_n = L$$

A sequence is not convergent if and only if:

- It has a subsequence with limit  $\pm\infty$
- It has two convergent subsequences with different limits.
- Examples:  $(-1)^n$ ,  $n$ ,  $(-1)^n n$

### DEFINITION

A sequence  $\{a_n\}$  is increasing if  $a_n < a_{n+1} \forall n$ . It is decreasing if  $a_n > a_{n+1} \forall n$ . A sequence which is either increasing or decreasing is called a monotonic.

A sequence  $\{a_n\}$  is bounded above if there exists a number  $M$  such that  $a_n \leq M \forall n$ . It is bounded below if there exists a number  $m$  such that  $a_n \geq m \forall n$ . It is bounded if it is bounded both above and below.

## 6 Series

An (infinite) series is an infinite summation:

$$a_1 + a_2 + a_3 + \cdots + a_n = \sum_{n=1}^{\infty} a_n$$

Infinite summations don't need to have a value, but some do.

### Definition

Let  $\sum_{n=1}^{\infty} a_n$  be a series. The  $n$ -th partial sum is:

$$S_n = a_1 + a_2 + a_3 + \cdots + a_n = \sum_{i=1}^n a_i \quad (12)$$

If the sequence  $S_n$  is convergent, then  $\sum_{n=1}^{\infty} a_n = S$ . The number  $S$  is the sum of the series. However if the sequence is divergent, then so is the series.

Note: If  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$

However, if  $\lim_{n \rightarrow \infty} a_n = 0 \neq \sum_{n=1}^{\infty} a_n$  is convergent.



## Test for Divergence

If  $a_n$  is divergent or  $\lim_{n \rightarrow \infty} a_n \neq 0$  then  $\sum_{n=1}^{\infty} a_n$  is divergent.

Typically, it is not too hard to determine whether a series is convergent or divergent, but most of the time it is extremely hard (often impossible) to determine precisely the sum of the series if it's convergent.

Some exceptions are:

## Telescoping Sums

Idea: Find  $b_n$  such that  $a_n = b_{n+1} - b_n$  or some variation on this idea (eg.  $a_n = b_{n+k} - b_n$  where  $k$  is a constant, or anything that leads up to a massive simplification when summing up).

**Example:**

$$\sum_{n=1}^{\infty} \frac{1}{n(n+1)} = \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \frac{1}{3 \cdot 4} + \dots$$

Solution: The  $n$ -th partial sum is,

$$S_n = \frac{1}{1 \cdot 2} + \frac{1}{2 \cdot 3} + \dots + \frac{1}{n(n+1)} = \sum_{i=1}^n \frac{1}{i(i+1)}$$

$$\begin{aligned} \sum_{n=1}^n \frac{1+i-i}{i(i+1)} &= \sum_{i=1}^n \frac{(i+1)}{i(i+1)} - \frac{i}{i(i+1)} \\ &= \sum_{i=1}^n \left( \frac{1}{i} - \frac{1}{i+1} \right) \\ &= 1 - \frac{1}{n+1} \end{aligned}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} S_n &= \lim_{n \rightarrow \infty} \left( 1 - \frac{1}{n+1} \right) = 1 \\ \sum_{n=1}^{\infty} \frac{1}{n(n+1)} &= 1 \end{aligned}$$

## Geometric Series

A geometric series is a series of the form  $a + ar + ar^2 + \dots = \sum_{n=1}^{\infty} ar^{n-1} = \sum_{n=0}^{\infty} ar^n$  where  $a$  and  $r$  are non-zero constants.

Fact: If  $|r| < 1$ , then the geometric series converges and its sum is  $\sum_{n=1}^{\infty} ar^{n-1} = \frac{a}{1-r}$

If  $|r| \geq 1$ , the geometric series diverges.

### Geometric Series Calculation:

$$S_n = a + ar + ar^2 + ar^3 + \dots + a^{n-1} \Big| \cdot r \quad (1)$$

$$rS_n = ar + ar^2 + ar^3 + \dots + a^{n-1} + a^n \quad (2)$$

Equation (1) - Equation (2)

$$\begin{aligned} S_n - rS_n &= a - ar^n \rightarrow (1-r)S_n = a(1-r^n) \\ &\rightarrow S_n = \frac{a(1-r^n)}{(1-r)} \end{aligned}$$

### Harmonic Series:

$$\sum_{n=1}^{\infty} \frac{1}{n} = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \frac{1}{5} + \dots$$

### Famous Series

$$\sum_{n=1}^{\infty} \frac{1}{2^n} = \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots = 1 \quad (\text{Zeno's Paradox})$$

$$\sum_{n=1}^{\infty} \frac{1}{n} = \frac{1}{1} + \frac{1}{2} + \frac{1}{3} + \dots = \zeta(1) \quad (\text{Zeta Function})$$

$$\sum_{n=0}^{\infty} \frac{1}{n!} = 1 + 1 + \frac{1}{2} + \frac{1}{6} + \frac{1}{24} \dots = e \quad (\text{Eulers Constant})$$

$$\sum_{n=0}^{\infty} \frac{(-1)^n}{n!} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \dots = \ln(2) \quad (\text{Proof of } \ln(2) \text{ being irrational})$$

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = 1 + \frac{1}{4} + \frac{1}{9} + \frac{1}{16} + \dots = \frac{\pi^2}{6} = \zeta(2) \quad (\text{Second Zeta Function})$$

$$\sum_{n=1}^{\infty} \frac{1}{n^4} = 1 + \frac{1}{16} + \frac{1}{81} + \frac{1}{256} + \dots = \frac{\pi^4}{96} = \zeta(4) \quad (\text{Rienmann's Zeta Function})$$

$$\sum_{n=1}^{\infty} \frac{1}{n^3} = 1 + \frac{1}{8} + \frac{1}{27} + \frac{1}{64} + \dots = \zeta(3) \quad (\text{Apéry's Construct})$$

The Harmonic series is divergent (It's the first example of a series which diverges in spite of the fact that the terms converge to 0)

## 7 Series Tests

These tests are used to compute whether a series is convergent or divergent.

### 7.1 Integral Test

Let  $f$  be continuous, positive, decreasing function on  $[1, \infty]$  and let  $a_n = f(n)$  then

- If  $\int_1^\infty f(x) dx$  is convergent,  $\sum_{n=1}^\infty a_n$  is convergent.
- If  $\int_1^\infty f(x) dx$  is divergent,  $\sum_{n=1}^\infty a_n$  is divergent.

Note: There is no need to start at  $n = 1$ , you can use  $\int_N^\infty f(x) dx$  where  $N$  is a number as large as you need.

### 7.2 P-Test

If  $p \in \mathbb{R}$ , then  $\sum_{n=1}^\infty \frac{1}{n^p}$  converges if  $p > 1$ , diverges if  $p \leq 1$

*NOTE*

Although the p-test by itself is quite limited, it will turn out to be very powerful if used in combination with the comparison tests.

The Comparison Test and the Limit Comparison Test allow us to determine whether a series is convergent or divergent by comparing it with a simpler series - most of the time a p-series or a geometric series.

### 7.3 Comparison Test

Let  $\sum_{n=1}^\infty a_n$  and  $\sum_{n=1}^\infty b_n$  be series with positive terms:

- If  $a_n \leq b_n$  for all  $n$ , and  $\sum_{n=1}^\infty b_n$  converges, then  $\sum_{n=1}^\infty a_n$  converges as well.
- If  $a_n \geq b_n$  for all  $n$ , and  $\sum_{n=1}^\infty b_n$  diverges, then  $\sum_{n=1}^\infty a_n$  diverges as well.

Note: "For all  $n$ " may be replaced with "For all sufficiently large  $n$ " which means "There exists  $N$  such that [the property holds] for all  $n \geq N$ ".

### 7.4 Limit Comparison Test

Let  $\sum_{n=1}^\infty a_n$  and  $\sum_{n=1}^\infty b_n$  be series with positive terms, and assume that the  $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$  where  $c$  is a finite number and  $c > 0$ . Then  $\sum_{n=1}^\infty a_n$  and  $\sum_{n=1}^\infty b_n$  have the same behavior. (If one converges, so does the other, and conversely, if one diverges, so does the other)

## 7.5 Alternating Series Test

A series is called alternating if its terms alternate between positive and negative. Assuming the first term is positive, we can write:

$$\sum_{n=1}^{\infty} a_n = \sum_{n=1}^{\infty} (-1)^{n-1} b_n$$

If the first term is negative, we can change the  $n - 1$  to  $n$ .

Given an alternating series as above, if:

- $b_n$  is eventually decreasing and;
- $\lim_{n \rightarrow \infty} b_n = 0$

Then the series is divergent.

*USEFULL FACT*

If  $a_n = \pm b_n$ , then  $\lim_{n \rightarrow \infty} a_n = 0$  if and only if  $\lim_{n \rightarrow \infty} b_n = 0$ .

*DEFINITION*

Let  $\sum_{n=1}^{\infty} a_n$  be a series.

- If  $\sum_{n=1}^{\infty} |a_n|$  is convergent, then  $\sum_{n=1}^{\infty} a_n$  is said to be absolutely convergent.
- If the series is convergent, but not absolutely convergent, it is said to be conditionally convergent.

## 7.6 Ratio Test

Let  $\sum_{n=1}^{\infty} a_n$  be a series:

- If  $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = L < 1$ , then the series is convergent (and in particular, absolutely convergent).
- If  $\lim_{n \rightarrow \infty} \frac{|a_{n+1}|}{|a_n|} = L > 1$  (or  $+\infty$ ), then the series is divergent.

## 7.7 Root Test

Let  $\sum_{n=1}^{\infty} a_n$  be a series:

- If  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L < 1$ , then the series is convergent (and in particular, absolutely convergent).
- If  $\lim_{n \rightarrow \infty} \sqrt[n]{|a_n|} = L > 1$  (or  $+\infty$ ), then the series is divergent.

In many cases, the limits of the root // ratio test are equal to 1, and then the tests are inconclusive.

## 8 Taylor and Maclaurin Series

The formula for a Taylor series Expansion is:

$$f(a) = \sum_{n=0}^{\infty} \frac{(f^n(a)(x-a)^n)}{n!}$$

Where  $f^n$  represents the derivative of the original function, some number of times (n).

It's purpose is to approximate a function using a series to see how it behaves. Another useful expansion is called the Maclaurin series. It is similar in nature to a Taylor Series, but the main difference is that the Maclaurin Series approximates the function at  $a = 0$ . instead of any arbitrary value of  $a$  like in a Taylor Expand. Think of it as just a more specific case of the Taylor Expansion. The formula for a Maclaurin Expansion is shown below:

$$f(0) = \sum_{n=0}^{\infty} \frac{f^n(0)(x)^n}{n!}$$

Both of these are used to identify many important proofs in science and math. Both series representations are often power series representations for  $f$ . However, there are a few issues with Taylor Series:

- Convergence of the right hand side.
- Even if the RHS converges, is it equal to  $f(x)$ ? Unfortunately, this is not always exactly true.

*EXAMPLE:*

Maclaurin Series of  $e^x$ :

$$f(a) = \sum_{n=0}^{\infty} \frac{(f^n(a)(x-a)^n)}{n!} = f(0) = \sum_{n=0}^{\infty} \frac{(f^n(0)(x-0)^n)}{n!}$$

$$f(0) = e^0 + e^0(x) + \frac{e^0(x)^2}{2} + \frac{e^0(x)^3}{6} + \frac{e^0(x)^4}{24} + \dots$$

$$f(0) = 1 + x + \frac{x^2}{2} + \frac{x^3}{6} + \frac{x^4}{24} + \dots$$

$$f(0) = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

We can do the same thing here for the Taylor Expansion of  $e^x$  about 1:

$$f(a) = \sum_{n=0}^{\infty} \frac{(f^n(a)(x-a)^n)}{n!} = f(1) = \sum_{n=0}^{\infty} \frac{(f^n(1)(x-1)^n)}{n!}$$

$$f(1) = e^1 + e^1(x) + \frac{e^1(x)^2}{2} + \frac{e^1(x)^3}{6} + \frac{e^1(x)^4}{24} + \dots$$

$$f(1) = e + e(x-1) + \frac{e(x-1)^2}{2} + \frac{e(x-1)^3}{6} + \frac{e(x-1)^4}{24} + \dots$$

$$f(1) = \sum_{n=0}^{\infty} \frac{e(x-1)^n}{n!}$$

## 8.1 Famous Examples of Taylor Series

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + \cdots = \sum_{n=0}^{\infty} x^n \quad (\text{if } -1 < x < 1)$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \cdots = \sum_{n=0}^{\infty} \frac{x^n}{n!} \quad (\text{for all } x)$$


$$\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \frac{x^9}{9!} - \cdots = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{(2n+1)!} \quad (\text{for all } x)$$

$$\cos(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \cdots = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!} \quad (\text{for all } x)$$

$$\ln(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \cdots = -\sum_{n=1}^{\infty} \frac{x^n}{n} \quad (\text{for } -1 < x < 1)$$

$$\tan^{-1}(x) = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \cdots = \sum_{n=0}^{\infty} \frac{(-1)^n x^{2n+1}}{2n+1} \quad (\text{for } -1 < x < 1)$$

$$(1+x)^\alpha = \sum_{n=0}^{\infty} \frac{\alpha(\alpha-1)\cdots(\alpha-n+1)}{n!} \cdot x^n \quad (\text{if } -1 < x < 1)$$



Taylor Series approximation of the graph  $\cos(x)$

— End of Document —