

Probability and Random Processes

Gude Pravarsh EE22BTECH11023*

Q) In an examination, 20 questions of true-false type are asked. Suppose a student tosses a fair coin to determine his answer to each question. If the coin falls heads, he answers 'true'; if it falls tails, he answers 'false'. Find the probability that he answers at least 12 questions correctly.

Solution: Let the random variable X denotes true(1) or false(0) for a question with p being the probability that true is answered. Then,

$$X \sim \text{Ber}(p) \quad (1)$$

Suppose $X_i, 1 \leq i \leq n$ represent each of the n draws. Define Y as

$$Y = \sum_{i=1}^n X_i \quad (2)$$

Then, since the X_i are iid, the pmf of Y is given by

$$Y \sim \text{Bin}(n, p) \quad (3)$$

The cdf of Y is given by

$$F_Y(k) = \Pr(Y \leq k) \quad (4)$$

$$= \begin{cases} 0 & k < 0 \\ \sum_{i=1}^k \binom{n}{i} p^i (1-p)^{n-i} & 1 \leq k \leq n \\ 1 & k \geq n \end{cases} \quad (5)$$

In this case,

$$p = \frac{1}{2}, \quad n = 20 \quad (6)$$

1) We require $\Pr(Y \geq 12)$. Since $n = 20$,

$$\Pr(Y \geq 12) = 1 - \Pr(Y < 12) \quad (7)$$

$$= F_Y(20) - F_Y(11) \quad (8)$$

$$= \sum_{k=12}^{20} p_Y(k) \quad (9)$$

$$= \sum_{k=12}^{20} \binom{n}{k} p^k (1-p)^{n-k} \quad (10)$$

$$= 0.2517 \quad (11)$$