

High Resolution Layer Croplands - Product User Manual (PUM)

Copernicus Land Monitoring Service



Author: **European Environment Agency (EEA)**

Date: **2025-03-28**

Version: **1**

Content

1. Executive summary.....	2
2. Background of the document.....	2
2.1 Scope.....	2
2.2 Content and structure.....	3
3. Lineage of HRL Tree Cover and Forests, Grasslands, and Croplands.....	3
4. Review of User Requirements.....	5
5. Product structure - What are the High Resolution Layers?.....	7
6. Product application areas and examples of use cases.....	9
6.1 Use case: C3/C4 crops specific parametrization for biomass monitoring.....	9
6.2 Use case: Land cover specific monitoring.....	10
7. Product description.....	10
7.1 Thematic characteristics of the HRL Croplands layers.....	14
7.2 Download content, file naming convention and file format(s).....	17
7.3 Projection and spatial coverage.....	19
7.4 Spatial resolution.....	20
7.5 Temporal information.....	20
7.6 Product characteristics and class codes.....	21
8. Production quality assessment.....	25
8.1 Layers to be verified.....	25
8.2 Sampling Design.....	26
8.3 Response Design.....	29
8.4 Statistical Analysis.....	31
8.5 Verification Results.....	32
8.5.1 Crop Group level.....	32
8.5.2 CTY2018 - EU27 (Weighted).....	32
8.5.3 CTY2018 - EU27 (Unweighted).....	33
8.5.4 CTY2021 - EU27 (Weighted).....	33
8.5.5 CTY2021 - EU27 (Unweighted).....	33
8.5.6 CTY2018 - EEA38 (Weighted).....	34
8.5.7 CTY2018 - EEA38 (Unweighted).....	34
8.5.8 CTY2021 - EEA38 (Weighted).....	35
8.5.9 CTY2021 - EEA38 (Unweighted).....	35
8.5.10 Aggregation level 1.....	36
9. Terms of use and product technical support.....	36
9.1 Terms of use.....	36

9.2 Citation.....	37
9.3 Product technical support.....	37
10. List of Abbreviations & Acronyms.....	38
11. References.....	40
12. Annex I – Colour tables for HRL Croplands.....	41
13. Document history:.....	59

Contact:

European Environment Agency (EEA)
Kongens Nytorv 6
1050 Copenhagen K
Denmark
<https://land.copernicus.eu/>

1. Executive summary

Copernicus is the European Union's Earth Observation Programme. It offers information services based on satellite Earth observation and in situ (non-space) data. These information services are freely and openly accessible to its users through six thematic Copernicus services (Atmosphere Monitoring, Marine Environment Monitoring, Land Monitoring, Climate Change, Emergency Management and Security).

The **Copernicus Land Monitoring Service (CLMS)** provides geographical information on land cover and its changes, land use, vegetation state, water cycle and earth surface energy variables to a broad range of users in Europe and across the world in the field of environmental terrestrial applications.

CLMS is jointly implemented by the **European Environment Agency (EEA)** and the European Commission's DG **Joint Research Centre (JRC)**.

The **High-Resolution Layer (HRL)** Vegetated land cover characteristics are a set of harmonised yearly maps dedicated to the thematic themes **Tree Cover & Forests, Grasslands and Croplands**. Those include a rich suite of raster products mapping the yearly status of those land cover types at a spatial resolution of 10 meters and change layers at 3-yearly interval and 20-meter resolution. HRL vegetated land cover characteristics extends the time-series of the existing HRL's Tree Cover & Forests and Grasslands and complements the CLMS portfolio with new layer dedicated to the mapping of crop types and agricultural practices such as mowing, harvest and cover crops.

2. Background of the document

2.1 Scope

This Product User Manual is the primary document that users are recommended to read before using the product. It provides a description of the product characteristics, production methodologies and workflows, and information about the product quality of the annual provision of **HRL Croplands**.

Furthermore, it gives information on the terms of use and product technical support. More detailed information on the methodologies and processing workflows that were used to produce the products can be found in the Algorithm Theoretical Baseline Document (ATBD) [7].

2.2 Content and structure

In more detail, the document is structured as follows:

- Chapter 3 provides an overview of the lineage of the products in relation to previous productions;
- Chapter 4 contains a review of user requirements;
- Chapter 5 provides an overview of what is included in the **High Resolution Layers** vegetated land cover characteristics and how the comprised products relate to each other;
- Chapter 6 presents potential application areas and example use cases;
- Chapter 7 provides a description of the products including the nomenclature and class definitions, file naming, spatial resolution format(s), etc.;
- Chapter 8 summarizes the quality assessment, validation procedure and the results;
- Chapter 9 provides information about product access and use conditions as well as the technical product support.

3. Lineage of HRL Tree Cover and Forests, Grasslands, and Croplands

High Resolution Layers (HRL) on Tree Cover & Forests had already been established in the **Copernicus Land Monitoring Service (CLMS)** portfolio since the reference years 2012 producing initially a **Dominant Leaf Type**, a **Tree Cover Density**, and a **Forest Type** map at a spatial resolution of 20 meters (Figure 3-1). Change layers and reference datasets were included with the reference year 2015. At the same time the accuracy targets were raised towards at least 90% user's and producer's accuracy. A further important step followed with the first production for the reference years 2018 (further referred to as **Historic HRL Forest 2018**) where the spatial resolution of the status layers was raised to 10 meters, the implementation of the change layers was partially reconsidered and target accuracies for the change layers were defined. In addition, new aggregated layer depicting the density of coniferous and broadleaved tree cover was introduced. With the **HRL Tree Cover & Forests** starting from the reference year 2018 the product specifications have been kept largely in line with the definitions used for the Historic HRL Forest 2018 whereas major changes concerned in particular the move to a yearly update cycle for the status layers and changes to some confidence layers (not shown in Figure 3-1). The new HRL Tree Cover & Forests therefore replace and extend the Historic HRL Forest 2018. This does not include an update of the change layer 2015 – 2018; the new status layers for 2018 are therefore not consistent with the original change layers 2015 – 2018.

HRL's on **Grasslands** had already been established in the Copernicus Land Monitoring Service (CLMS) portfolio since the reference years 2015 producing initially a status layer on the absence / presence of



grassland (Figure 3-1) with a target Overall Accuracy of 85%. With the reference year 2018 the spatial resolution of the status layers was raised to 10 meters and a change layer with a target Overall Accuracy of 80% was introduced. With the **HRL Grasslands** starting from the reference year 2018 the product specifications have been kept largely in line with the definitions used for the **Historic HRL Grassland 2018** whereas major changes concerned in particular the move to a yearly update cycle for the status layers, an additional yearly **Herbaceous Cover layer (HER)** (including also temporary grassland in the reference year), new layers on the count and timing of Grassland Mowing (Minimum Mapping Unit (MMU) of 0.25 ha) and changes to some confidence layers (not shown in detail in Figure 3-1). The new HRL Grasslands therefore replaces and extends the Historic HRL Grassland 2018. This does not include an update of the change layer 2015 – 2018; the new status layers for 2018 are therefore not consistent with the original change layers 2015 – 2018.

The **HRL Croplands** is a new set of layers dedicated to agriculture and comprises several yearly layers mapping crop types (10-metre spatial resolution) and agricultural practices such as harvest, fallow land and secondary crops (10-metre spatial resolution, Minimum Mapping Unit of 0.25 ha).




Figure 3-1 Evolution of HRL Forest and Grassland towards the three product groups HRL Tree Cover & Forests, HRL Grasslands, HRL Croplands.

4. Review of User Requirements

In the frame of the **Horizon 2020 (H2020)** project **ECOLaSS** a survey [5] of key stakeholders has been performed in order to evaluate the user requirements towards the evolution of existing and future **Copernicus** products. This survey made also use of the results from the Nextspace User Study [6] and revealed that **HRL** users like European institutions, service industry, research and academia, national agencies, regional administrations, NGS or private users would in general appreciate:

- High thematic quality/meaningful and application-oriented product definitions;
- Sufficient spatial and timely resolution concerning both, status layer and change layer;

- Short update cycles;
- Change monitoring;
- Free and open access;
- High technical quality;
- Standardized and comparable nomenclature;
- Transparent and scientific workflows and state-of-the-art methodology;
- Detailed documentation of these workflow and the respective methodology;
- Consistency of the Pan-European products enabling synergistic use of all products;
- Streamlining the pan-European product with global ones;
- Availability of historic data and compatibility of time series;
- Open access to the original Copernicus Sentinel data;
- Sophisticated product presentation and visualisation possibilities in an online viewer on the Copernicus platform;
- IPCC conformity.

While many of these requirements had already been satisfied with previous HRL reference years some could only be implemented within the current update:

- A long-standing thematic gap in the European **CLMS** portfolio concerning the monitoring of agriculture has been addressed through new products on crop types and agricultural activities. This also improves the separation between grassland and cropland and the IPCC conformity;
- Yearly update cycle for status layer;
- Grassland use intensity (or the dynamics of intensification/ extensification) is partially addressed through a new product on Grassland mowing.

Further requirements that remain to be considered for future updates are for example:

- More fine-grained differentiation among species-rich (extensively used) and separation from species-poor (intensively used) and managed grassland;
- Tree-species compositions and shifts between extensive and intensive management;
- Increased timeliness of availability of the products: The mid-term goal is a product provision at latest 12 months after the end of the reference year.

5. Product structure - What are the High Resolution Layers?

The **High Resolution Layers (HRL)** vegetated land cover characteristics portfolio consists of **Tree Cover & Forests**, **Grasslands** and **Croplands** products (Figure 5-1), which together cover most of what is defined as the Biotic component of the **EAGLE** Land Cover Components¹. More specifically, the mapping is focused on surfaces with a vegetation cover above 30%; an exception to this is tree cover where the objective is to map tree cover with a continuous range of 1-100% **Tree Cover Density (TDC)**, i.e. also below 30%, as far as detectable from 10-metre resolution satellite imagery. This definition is also in line with the Sparsely Vegetated class in the **Corine Land Cover (CLC)+ Backbone Raster**² and considers that detection / classification of vegetation below this threshold is typically more error-prone. The definition also aims at largely avoiding overlaps with the non-vegetated land cover characteristics such as **HRL Imperviousness**, which is focused on areas with less than 10% vegetation cover during any time of the year, for a reference period of 3 year.

Some overlaps between the three product groups are allowed by definition, for example areas with tree crops (i.e. olive, fruit and nut trees) which are included in both the Tree Cover & Forests and the Croplands products. Furthermore, specific vegetations types are not included in the HRLVLCC portfolio; this concerns areas dominated by natural shrubs (i.e. shrubs that are not under agricultural use) and associations of lichens and mosses.




Figure 5-1: Products within the HRL vegetated land cover characteristics.

Given several interdependencies and potential overlaps among the **Grasslands**, **Croplands** and **Tree Cover & Forests** products, the overall workflow starts with the classification of **Base Vegetation Layer (BVL)**. A high-level description is provided for the overall workflow in Figure 5-2. The yearly BVL classification initially targets the separation of five basic land cover classes being:

¹ https://land.copernicus.eu/en/eagle?tab=technical_implementation

² <https://land.copernicus.eu/en/technical-library/product-user-manual-clc-backbone-2021>

1. herbaceous vegetation;
2. cropland;
3. tree cover;
4. tree crops (i.e. nomenclature overlap between broadleaved tree cover and permanent crops in the Croplands product);
5. background class (including bare and sparsely vegetated areas and non-agricultural shrubs);

In a subsequent post-processing step two further classes are derived to delineate the:

6. potential overlap herbaceous – cropland (i.e. pixels which are classified as cropland and herbaceous at least once in the time-series);
7. The second derived class is derived from the intersection of all areas classified as tree cover and a preliminary version of the Tree Cover Density to delineate areas with low Tree Cover Density and hence allowed overlaps of herbaceous and tree cover.

The derived yearly BVL is considered for the downstream productions of Grasslands, Croplands and Tree Cover & Forests products as follows:

- For the production of the Grasslands layers: all areas classified as herbaceous, overlap herbaceous – tree cover, or overlap herbaceous – Cropland are considered as the potential maximum extent for the **Herbaceous Cover (HER)** layer. In addition, the BVL classification probabilities for the herbaceous class are used as the main input for the derivation of the **HER** layer.
- For the Croplands layers: the areas delineated as cropland, overlap herbaceous –cropland, or Tree Crops are considered as the maximum extent for the **CTY** classification and further refinement.
- For the **Tree Cover & Forests** layers: the areas classified as tree cover, overlap herbaceous – tree cover, tree crops and the respective probabilities are used directly to derive the respective change layers and yearly **DLT** and **TCD** status layers.

Within the areas identified as overlap herbaceous – cropland, a further harmonization step is carried out downstream. To this end the **CTY** classification initially includes a class for fodder crops which are transferred to the **HER** layer if occurring in the designated overlap class.




Figure 5-2: High-level overview of the relationship between the Base Vegetation Layer and the subsequent production of Grasslands, Croplands and Tree Cover & Forests products

6. Product application areas and examples of use cases

The **HRL Tree Cover & Forests, Grasslands and Croplands** set of products is designed for use by a broad user community as basis for environmental and regional analysis and for supporting political decision-making, such as the **Common Agricultural Policy (CAP)**, **LULUCF (Land Use, Land Use Change and Forestry)** regulation, the **Nature Restoration Regulation (NRR)**, or the proposed **European Forest Monitoring Law (FML)**. With the new products the **EEA** will ensure continuity and further densification of the well-established HRL Tree Cover & Forests and Grasslands product time series. Those include a rich suite of raster products at a 3-yearly interval mapping the status of those land cover types with a spatial resolution of 10-meter and change layers at 20-meter spatial resolution.

As an example, the following sections provide short information on (potential) use cases at national level, for which the **Copernicus HRL Croplands** product represent a fundamental input.

6.1 Use case: C3/C4 crops specific parametrization for biomass monitoring.

The **HRL Crop Types (CTY)** layer offers comprehensive information on various crop types, allowing for further differentiation between C3 and C4 crops (if class not mixed-up). Maize, being the most common C4 crop cultivated in Europe could be easily derived from this dataset. By providing detailed information on crop types and their specific locations, the **CTY** layer enables the use of tailored parameters for biomass estimation at C3/C4 crop locations on an annual basis. This information is particularly relevant for on-going work within the **EvoLand³** project, where the enhancement of biomass mapping for both crop and grassland areas is envisaged. In this respect, the reliability of the biomass mapping could

³ <https://www.evo-land.eu/method/biomass-mapping/>

benefit from the availability of this detailed crop type information. Moreover, the distinction between C3 and C4 crops is useful for better estimating autotrophic respiration, which in turn allows for more accurate conversions of gross primary productivity to net primary productivity. Given that C3 and C4 crops have distinct photosynthetic pathways and respiration rates, this differentiation supports the use of more precise parameters in productivity models, contributing to enhanced biomass mapping across crop landscapes.

6.2 Use case: Land cover specific monitoring

Detailed and dynamic information on the state of the land as provided by the **HRL Croplands** layers allows to analyse regional trends in the area occupied by these land covers, which could be relevant information for authorities and policy makers. Furthermore, applications which require information on the land cover status can benefit from the HRL Croplands. For example, in case of biomass mapping often land cover specific parametrization is applied. Using **HRL Croplands**, allows to do this in a much more detailed and dynamic way. In case of the Evoland project, it is intended to use these layers to apply specific parametrization over crop and grassland locations.

7. Product description

The **HRL Croplands** layers are generally provided in 100km Lambert Azimuthal Equal Area (LAEA) projection tiles as shown in Figure 7-1. The five French Oversea Territories are provided in UTM with the layout of the respective territory. The layers are available as Cloud-Optimized GeoTIFFs (COG) per reference year and 100km LAEA tile aligned with the **EEA reference grid**. Each raster file is accompanied by a Persistent Auxiliary Metadata (PAM) XML and an INSPIRE XML.




Figure 7-1: LAEA tile layout incl. distinction between tiles to cover EU27 and EEA38.

The **HRL Croplands** information aims at filling the data gap needed for detailed agricultural monitoring and enables repeatable and harmonised continental scale assessments within the **CLMS** portfolio. The **HRL Croplands** layers entail the mapping the extent of croplands and crop types, delineation of parcel-complexes, mapping agricultural practices on large scale and with wall-to-wall coverage and high spatial details.




Figure 7-2 HRL Croplands product overview

The primary layer is the yearly **Crop Types (CTY)** layer at 10m, which maps the specific crop grown in the main growing season for all pixels identified as arable or permanent crops by the **BVL**. Maximum uniformity, comparability, and interpretability of the final product is guaranteed across different regions in Europe by focusing on the main growing season. Besides the **CTY** layer a supportive **Crop Types Confidence Layer (CTYCL)** layer is provided to match the **CTY** layer. **CTYCL** indicates the degree of confidence of the provided crop type label. The probability value from the winning class as determined by the model is taken as the confidence level.

Several further status **HRL Cropping Patterns (CP)** layers, covering 5 thematic domains, give information on the agricultural practices within the classified cropland area for the following domains:

- Main Crops;
- Bare Soil;
- Secondary Crops;
- Fallow Land;
- Cropping Seasons;

All the **CP** layers are only applied over annual arable land, excluding permanent crops. Permanent crops do not have specific seasonal cycles, which is the primary focus of the cropping pattern layers. The workflow used to define the different agricultural practices is applied at field level scale, to ensure homogeneity in the seasonal signal and reduce signal noise. Afterwards, all the different layers are rasterized and aligned on the same spatial grid as the **CTY** layer. Therefore, only for those locations where initially a field could be delineated and where it coincides with a classified annual cropland, the Cropping Pattern layers are generated (excluding crop rotation layer, see last paragraph of section). Since the MMU is applied to the originally delineated parcel geometry used for defining CP information, further alignment with annual cropland may result in smaller spatial patterns due to the clipping of the original field boundaries. However, the CP information itself remains based on an MMU of at least 0.25 ha, as it is still derived from the original parcel geometry.

The **Main Crops (MC)** layers focus on the characteristics of the main growing season which usually takes place during (part of) the summer season. These layers provide information on the **emergence (CPMCE)** and **harvest (CPMCH)** of the main growing season, as well as the **duration of the season (CPMCD)**. Each of these layers include a **Confidence Layer (CL)** that indicates the level of uncertainty in days. Information on the characteristics of the growing season and its variations over time give valuable information on how growing seasons are impacted by (extreme) weather conditions and climate change at regional and national levels.

The **Bare Soil (BS)** group of layers include information on the duration of the bare soil period. Presence of a period of **Bare Soil Before (CPBSB)** the emergence of the main crop season and **Bare Soil After (CPBSA)** the harvest of the main crop season are evaluated. Both layers also feature a CL that indicates the level of uncertainty in the duration of the bare soil period in days. This uncertainty is based on the uncertainty of the emergence and/or harvest events that define the bare soil period. The bare soil period is only defined for the duration within the reference year. Therefore, emergence events occurring after the calendar year that marks the end of the bare soil period following the main season will not be considered in the uncertainty calculation. Similarly, for the bare soil period before the main season, if the preceding harvest event took place in the previous calendar year, it will not be included in the uncertainty calculation. Information of bare soil periods are relevant for policy makers to assess the susceptibility of soils to leaching and erosion.

The **Secondary Crops (SC)** group of layers provides information on agricultural parcels where an off-season secondary growing season occurs. Typically, this secondary season involves planting a cover crop in the field. Secondary season presence is only evaluated if a main season is detected. The secondary season is categorized into four **Secondary Crop Types (CPSCT)** based on the emergence date and length of the season: short/long summer and short/long winter secondary seasons. In addition, to this

classification, information on the **Secondary Crop Emergence (CPSCE)** and **Secondary Crop Duration (CPSCD)** of the season is included. Similarly, as for the **Main Crops (MC)** layer, uncertainty information is provided for both the duration and emergence of the secondary crops. The **SC** layers offer valuable information for assessing policy implementation to prevent soil degradation and monitoring cropping changes during off-season over time at regional and national level.

The **Fallow Land (FL)** layers give information on **Fallow Land Presence (CPFLP)** and **Fallow Land Duration (CPFLD)**. Fallow land presence is defined as agricultural land where, temporarily, no agricultural activity takes place. Fallow land duration indicates the number of days a parcel is left fallow for the five-year period i.e., 2017-2021. Both the **FLP** layer and the **FLD** layer have an accompanying **Confidence Layer (CL)**.

Finally, the **Cropping Seasons** layers provide information on the number of detected growing seasons in the **Cropping Seasons Yearly (CPCSY)** layer and information on main crop rotation in the **Cropping Seasons Types over 3 years (CPCST)** layer. The **CPCSY** layer is based on the **CTY** and the **CPSC** layers to count the number of growing seasons in a specific year. **CPCST** is based on the crop diversity count and contains the number of unique crop types grown in the main agricultural season over a 3-year period. Since, **CPCST** is based on information of previous year's crop types, it is only available for the 2017-2019, 2018-2020, and 2019 till 2021 time periods. Furthermore, it does not rely on the delineation of parcel complexes compared to the other **CP** layers as it is a purely **CTY** based layer. Information on crop rotation provides valuable insights to monitor the application of monoculture or rotation cycle in specific regions.

7.1 Thematic characteristics of the HRL Croplands layers

The **HRL Croplands** layers are only provided for those locations where the **BVL** designates arable or permanent cropland (i.e., **BVL** classes cropland, tree crops and overlap herbaceous and crops). For those locations the pixel-based crop type algorithm classifies according to the crop type grown in the **main growing season**. A main crop season is defined as the most crucial period for cultivating crops of economic value. Initially, the algorithm also classifies the grassland and fodders crops that are present in the applied **BVL** mask (overlap herbaceous and crops), but those locations are transferred to the **Herbaceous Cover (HER)** layer (see 5.2). In Table 7-1, the final mapped crop types and the aggregation levels of the hierarchical nomenclature are given. The crop type aggregation level is used within the **Crop Types (CTY)** layer. For those cases where the classification algorithm remains undecided due to low confidence values, a separate category is added, subdivided in unclassified arable and permanent crop. This output is further spatially cleaned and harmonised over the years to enhance temporal and spatial consistency of the product. As a result of these post-processing steps, it might happen that there is some minimal overlap between the **HRL Croplands** and **Grassland** layers that are both located within the arable or permanent crops classes defined by **BVL** [7]. This potential overlap can be present in the published layers and solely occurs at borders between cropland and temporary grassland.

Code	Land Cover	Crop Group	Crop type
1110	Arable Crops	Cereals	Wheat
1120			Barley
1130			Maize
1140			Rice
1150			Other cereals
1210		Dry pulses & Vegetables	Fresh Vegetables
1220			Dry pulses
1310		Root/tuber crops	Potatoes
1320			Sugar Beet
1410		Non-permanent industrial crops	Sunflower
1420			Soybeans
1430			Rapeseed
1440			Flax, cotton and hemp
2100	Permanent Crops	Permanent Crops	Grapes
2200			Olives
2310			Fruits
2320			Nuts
3100	Arable Crops	Unclassified arable crop	Unclassified arable crop
3200	Permanent Crops	Unclassified permanent crop	Unclassified permanent crop

Table 7-1: HRL Crop Types nomenclature and aggregation levels

The **emergence** and **harvest** detection are the main input for the different **HRL Cropping Patterns (CP)** layers that provide information on the agricultural practices. The following definitions of emergence and harvest are used to detect these events from satellite imagery:

- Emergence is defined as the moment when the first leaves of the crop unfold
- Harvest is defined as the moment when all standing biomass is removed.

A **main crop season** is defined as the most crucial period for cultivating crops of economic value. Expert rules, developed by analysing seasonal boundaries in Europe and using USDA crop calendar data, are used to classify seasons as main or secondary [7]. Region specific seasonal boundaries are selected to reflect the seasonal shifts across Europe. Additionally, rules for prioritization are established to resolve conflicts in cases where multiple main seasons are possible. The complete procedure for selecting the growing season is outlined in the **ATBD**; while the detection is only performed for annual crops it is fully independent from the crop type label of the annual crop defined in the **CTY** layer [7]. However, due to the complexity of agricultural patterns, there may be challenges in accurately distinguishing the main season from the secondary season. This is particularly true in cases where a very late onset of the main growing season follows on a season with similar behaviour as winter cereals. Since the cultivation of winter cereals is also considered, the main season onset may occur before the reference year.

Additionally, in rare instances, the harvest of the main season crop may occur shortly after the end of the year. In some situations, it is impossible to generate a **Cropping Pattern** product for the **main crop** layer based on the defined seasonal boundaries (i.e., season length should fall between 40 and 366 days, period of main growing season) and the detection of emergence and harvest events. In such cases, the corresponding pixels are flagged with dedicated codes (see Table 7-7).

The **Secondary Crop (SC)** layers follow a similar methodology to the main season classification, but with distinct seasonal boundaries based on the emergence and harvest detected for the secondary crop. The secondary season can occur both preceding and following the main growing season, although preference is given to labelling a secondary season that follows the main season when multiple instances arise. If there are more than one seasonal signal after the main season within the same calendar year, the **SC** product may not align with the actual cover crop sown, but rather with a short-term crop planted in a double cropping system. Each secondary crop season is subdivided into four categories, short/long summer, and short/long winter secondary seasons. A seasonal length shorter than 100 days is considered as a short growing variant, otherwise it is a long variant. If the secondary crop emergence is before September and takes place after the end of the main season, it is a summer variant. In all other cases, it is a winter variant.

The **Bare Soil** layers indicate the period before and after the main growing season when no crop cultivation activities occur. This period is determined by the detection of emergence and harvest events within that year in a particular field. Since a specific bare soil detection algorithm is not employed, there may be instances where the field is not completely bare during this period, possibly due to weed growth or the presence of residual crop remnants. Additionally, activities such as fertilization or ploughing of the field can fall within the bare soil period.

The **Fallow Land** layers consider only delineated parcels that are classified as annual cropland by the **CTY** layer, other parcels are omitted for the fallow land evaluation. For the determination of fallow land presence thematic rules, based on the expected behaviour of fallow land, are used. The rules consider the average **Crop Types Confidence Layer (CTYCL)**, the number and sequence of detected harvests and emergences in combination with the maximum **fAPAR** value. Harvest dates are used to determine the **fallow land duration**. Therefore, the accuracy of the **fallow land layers** depends on the accuracy of the **CTYCL**, the detected harvests and emergence, and the **fAPAR** timeseries.

The **Cropping Seasons** layers provide insights into the annual crop rotation strategy and the number of growing seasons at specific locations. Note however that a maximum limit of two growing seasons is considered. If more than two seasons are identified, they will not be considered. Permanent crops are ignored in counting the crop diversity.

An overview of how cropping pattern layers(excluding Fallow Land and Cropping Seasonslayers) can be generated at the field level based on the identified emergence and harvest events within a six-month period before and after the calendar year is given in Figure 7-3. The emergence and harvest events are used to classify the season as either main or secondary based on predefined seasonal boundaries. In the example shown in Figure 7-3, the main season corresponds to a winter cereal, resulting in no bare soil period before the main season for the reference year. Instead, a bare soil period is indicated between the main and secondary seasons, defined by the harvest of the main season and the emergence of the secondary season. The secondary season is classified as a long winter variant. The uncertainty surrounding **fAPAR** values around the emergence and harvest events is utilized to generate the respective **Confidence Layers**.




Figure 7-3: Example how the different cropping patterns layers for a barley field located in Belgium are generated for the reference year 2019. The season delineation is based on the predicted emergence (black line) and harvest dates (red line). The green timeseries curve is the daily fAPAR value (with uncertainty). BS after stands for the length of the bare soil period after the main season.

7.2 Download content, file naming convention and file format(s)

Table 7-2: Download content, file naming convention and file format(s) for the HRL Croplands layers

Name of layer	Acronym	Abbreviation
Crop types	CTY	CTY_S2017_R10mCTY_S2018_R10mCTY_S2019_R10mCTY_S2020_R10mCTY_S2021_R10m
Crop types confidence	CTYCL	CTYCL_S2017_R10mCTYCL_S2018_R10mCTYCL_S2019_R10mCTYCL_S2020_R10mCTYCL_S2021_R10m
Main Crop Harvest	CPMCH	CPMCH_S2017_R10mCPMCH_S2018_R10mCPMCH_S2019_R10mCPMCH_S2020_R10mCPMCH_S2021_R10m
Main Crop Harvest Confidence Layer	CPMCHCL	CPMCHCL_S2017_R10mCPMCHCL_S2018_R10mCPMCHCL_S2019_R10mCPMCHCL_S2020_R10mCPMCHCL_S2021_R10m
Main Crop Emergence	CPMCE	CPMCE_S2017_R10mCPMCE_S2018_R10mCPMCE_S2019_R10mCPMCE_S2020_R10mCPMCE_S2021_R10m
Main Crop Emergence Confidence Layer	CPMCECL	CPMCECL_S2017_R10mCPMCECL_S2018_R10mCPMCECL_S2019_R10mCPMCECL_S2020_R10mCPMCECL_S2021_R10m
Main Crop Duration	CPMCD	CPMCD_S2017_R10mCPMCD_S2018_R10mCPMCD_S2019_R10mCPMCD_S2020_R10mCPMCD_S2021_R10m
Main Crop Duration Confidence Layer	CPMCDCL	CPMCDCL_S2017_R10mCPMCDCL_S2018_R10mCPMCDCL_S2019_R10mCPMCDCL_S2020_R10mCPMCDCL_S2021_R10m
Bare Soil Before	CPBSB	CPBSB_S2017_R10mCPBSB_S2018_R10mCPBSB_S2019_R10mCPBSB_S2020_R10mCPBSB_S2021_R10m
Bare Soil Before Confidence Layer	CPBSBCL	CPBSBCL_S2017_R10mCPBSBCL_S2018_R10mCPBSBCL_S2019_R10mCPBSBCL_S2020_R10mCPBSBCL_S2021_R10m
Bare Soil After	CPBSA	CPBSA_S2017_R10mCPBSA_S2018_R10mCPBSA_S2019_R10mCPBSA_S2020_R10mCPBSA_S2021_R10m

Name of layer	Acronym	Abbreviation
Bare Soil After Confidence Layer	CPBSACL	CPBSACL_S2017_R10mCPBSACL_S2018_R10mCPBSACL_S2019_R10mCPBSACL_S2020_R10mCPBSACL_S2021_R10m
Secondary Crops Type	CPSCT	CPSCT_S2017_R10mCPSCT_S2018_R10mCPSCT_S2019_R10mCPSCT_S2020_R10mCPSCT_S2021_R10m
Secondary Crops Emergence	CPSCE	CPSCE_S2017_R10mCPSCE_S2018_R10mCPSCE_S2019_R10mCPSCE_S2020_R10mCPSCE_S2021_R10m
Secondary Crops Duration	CPSCD	CPSCD_S2017_R10mCPSCD_S2018_R10mCPSCD_S2019_R10mCPSCD_S2020_R10mCPSCD_S2021_R10m
Secondary Crops Duration Confidence Layer	CPSCDCL	CPSCDCL_S2017_R10mCPSCDCL_S2018_R10mCPSCDCL_S2019_R10mCPSCDCL_S2020_R10mCPSCDCL_S2021_R10m
Fallow Land Presence	CPFLP	CPFLP_S2017_R10mCPFLP_S2018_R10mCPFLP_S2019_R10mCPFLP_S2020_R10mCPFLP_S2021_R10m
Fallow Land Presence Confidence Layer	CPFLPCL	CPFLPCL_S2017_R10mCPFLPCL_S2018_R10mCPFLPCL_S2019_R10mCPFLPCL_S2020_R10mCPFLPCL_S2021_R10m
Fallow Land Duration	CPFLD	CPFLD_P2017-2021_R10m
Fallow Land Duration Confidence Layer	CPFLDCL	CPFLDCL_P2017-2021_R10m
Cropping Seasons Yearly	CPCSY	CPCSY_S2017_R10mCPCSY_S2018_R10mCPCSY_S2019_R10mCPCSY_S2020_R10mCPCSY_S2021_R10m
Cropping Seasons Types over 3 years	CPCST	CPCST_P2017-2019_R10mCPCST_P2018-2020_R10mCPCST_P2019-2021_R10m

Delivery format (for Crop types & Crop types confidence):
→ Tiles of Cloud-Optimized GeoTIFF aligned with the 100km LAEA grid with embedded colormaps, as well as separate colour legends in the formats .qml, .sld and *.lyr.

Delivery format (for all rows except Crop types & Crop types confidence): → Tiles of Cloud-Optimized GeoTIFF aligned with the 100km LAEA grid. Separate colour legends in the formats .qml, .sld and *.lyr.

Metadata (applies to all rows):
→ XML metadata files according to INSPIRE metadata standards and GDAL-style Permanent Auxiliary Metadata (PAM) *.aux.xml including statistics and Raster Attribute Table.

7.3 Projection and spatial coverage

Table 7-3: Projection and spatial coverage for the HRL Croplands layers

Name of layer	Acronym
Crop types	CTY
Crop types confidence	CTYCL
Main Crop Harvest	CPMCH
Main Crop Harvest Confidence Layer	CPMCHCL
Main Crop Emergence	CPMCE
Main Crop Emergence Confidence Layer	CPMCECL
Main Crop Duration	CPMCD
Main Crop Duration Confidence Layer	CPMCDCL
Bare Soil Before	CPBSB

Name of layer	Acronym
Bare Soil Before Confidence Layer	CPBSBCL
Bare Soil After	CPBSA
Bare Soil After Confidence Layer	CPBSACL
Secondary Crops Type	CPSCT
Secondary Crops Emergence	CPSCE
Secondary Crops Duration	CPSCD
Secondary Crops Duration Confidence Layer	CPSCDCL
Fallow Land Presence	CPFLP
Fallow Land Presence Confidence Layer	CPFLPCL
Fallow Land Duration	CPFLD
Fallow Land Duration Confidence Layer	CPFLDCL
Cropping Seasons Yearly	CPCSY
Cropping Seasons Types over 3 years	CPCST

Spatial coverage (applies to all rows): → 5.751.002 km² (covering the full EEA-38)

Coordinate reference system (WKT) (applies to all rows): → PROJCS["ETRS89-extended / LAEA Europe", GEOGCS["ETRS89", DATUM["European_Terrestrial_Reference_System_1989", SPHEROID["GRS 1980",6378137,298.257222101, AUTHORITY["EPSG","7019"]], AUTHORITY["EPSG","6258"]], PRIMEM["Greenwich",0, AUTHORITY["EPSG","8901"]], UNIT["degree",0.0174532925199433, AUTHORITY["EPSG","9122"]], AUTHORITY["EPSG","4258"]], PROJECTION["Lambert_Azimuthal_Equal_Area"], PARAMETER["latitude_of_center",52], PARAMETER["longitude_of_center",10], PARAMETER["false_easting",4321000], PARAMETER["false_northing",3210000], UNIT["metre",1, AUTHORITY["EPSG","9001"]], AXIS["Northing",NORTH], AXIS["Easting",EAST], AUTHORITY["EPSG","3035"]] Except for French DOMs where the following CRS are used: YT: EPSG 32738 RE: EPSG 32740 MQ: EPSG 32620 GP: EPSG 32620 GF: EPSG 32620

7.4 Spatial resolution

Table 7-4: Spatial resolution for the HRL Croplands layers

Name of layer	Acronym
Crop types	CTY
Crop types confidence	CTYCL
Main Crop Harvest	CPMCH
Main Crop Harvest Confidence Layer	CPMCHCL
Main Crop Emergence	CPMCE
Main Crop Emergence Confidence Layer	CPMCECL
Main Crop Duration	CPMCD
Main Crop Duration Confidence Layer	CPMCDCL
Bare Soil Before	CPBSB
Bare Soil Before Confidence Layer	CPBSBCL
Bare Soil After	CPBSA

Name of layer	Acronym
Bare Soil After Confidence Layer	CPBSACL
Secondary Crops Type	CPSCT
Secondary Crops Emergence	CPSCE
Secondary Crops Duration	CPSCD
Secondary Crops Duration Confidence Layer	CPSCDCL
Fallow Land Presence	CPFLP
Fallow Land Presence Confidence Layer	CPFLPCL
Fallow Land Duration	CPFLD
Fallow Land Duration Confidence Layer	CPFLDCL
Cropping Seasons Yearly	CPCSY
Cropping Seasons Types over 3 years	CPCST

Pixel size (applies to all rows): → 10m

MMU (applies to all rows): → 0.25ha*

- For CP layers, smaller patterns may appear due to alignment with annual cropland. However, the CP information is derived from the original parcel geometries (before alignment), which adhere to the MMU requirements.

7.5 Temporal information

Table 7-5: Temporal information for the HRL Croplands layers

Name of layer	Acronym	Reference year
Crop types	CTY	20172018201920202021
Crop types confidence	CTYCL	20172018201920202021
Main Crop Harvest	CPMCH	20172018201920202021
Main Crop Harvest Confidence Layer	CPMCHCL	20172018201920202021
Main Crop Emergence	CPMCE	20172018201920202021
Main Crop Emergence Confidence Layer	CPMCECL	20172018201920202021
Main Crop Duration	CPMCD	20172018201920202021
Main Crop Duration Confidence Layer	CPMCDCL	20172018201920202021
Bare Soil Before	CPBSB	20172018201920202021
Bare Soil Before Confidence Layer	CPBSBCL	20172018201920202021
Bare Soil After	CPBSA	20172018201920202021
Bare Soil After Confidence Layer	CPBSACL	20172018201920202021
Secondary Crops Type	CPSCT	20172018201920202021
Secondary Crops Emergence	CPSCE	20172018201920202021
Secondary Crops Duration	CPSCD	20172018201920202021
Secondary Crops Duration Confidence Layer	CPSCDCL	20172018201920202021
Fallow Land Presence	CPFLP	20172018201920202021
Fallow Land Presence Confidence Layer	CPFLPCL	20172018201920202021
Fallow Land Duration	CPFLD	2017-2021

Name of layer	Acronym	Reference year
Fallow Land Duration Confidence Layer	CPFLDCL	2017-2021
Cropping Seasons Yearly	CPCSY	20172018201920202021
Cropping Seasons Types over 3 years	CPCST	2017-20192018-20202019-2021

7.6 Product characteristics and class codes

Table 7-6: Overview of the used quality flags within the HRL Croplands layers

Quality flag	Definition
0	Flags locations where the respective Croplands product could not be produced (used in all except CPFLPCL & CTYCL). Below an overview of the meaning per layer category: CTY : No cropland CPCST : Over the three years period this location is not consistently classified as an annual crop. CPFLP , CPFLD , CPFLDCL : No fallow land location. All other CP layers (excluding CPFLPCL) : No annual cropland
253	CPFLPCL : No fallow land CTYCL : No cropland.
255	Outside area in case of CTYCL .
65526	Fallow land occurrence and hence only fallow land related layers will be generated. The other layers will be flagged with at locations of fallow land (used for all cropping pattern layers, excluding CPCST , CPFLD , CPFLDCL , CPFLPCL , CPFLP).
65527	No delineated field geometry overlaps with an annual cropland location (from CTY) and hence no Cropping Pattern (CP) layers could be defined (used for all CP layers, excluding CPCST , CPFLD , CPFLDCL , CPFLPCL , CPFLP).
65528	No CPCST could be defined for this location. This occurs when the CTY has labelled this location in the reference year as a permanent or grassland or fodder crop, whereas in the previous year(s) it was classified as an annual crop (used only for CPCST layer).
65529	It is not possible to derive this specific bare soil period (before or after) since it does not occur within the calendar boundaries of the reference year (used for all Bare Soil related layers).
65530	No secondary crop growing season could be delineated beyond the main growing season delineated for that location (used for all Secondary Crop layers).
65531	There are not enough satellite observations to utilize the emergence and harvest detection algorithm, which is crucial for deriving the cropping pattern layers. Insufficient observations occur when there is a temporal gap of more than 60 days in Sentinel-1 observations or when no Sentinel-2 observations are available after cloud masking, which may occur near bright surfaces (used for all Cropping Patterns , except Fallow Land related layers and CPCST).
65532	No season delineation is possible based on the output of the emergence and harvest detection algorithms. This can happen under the following situations:• No harvest event or emergence event has been detected.• No harvest event does occur after the emergence event.• All the linked season(s) are too long (>366 days) or too short (<40 days).• The start of the linked season(s) is after the start of the reference year.• The end of the linked season(s) is before the start of the reference year. Used for all Cropping Pattern layers, except Fallow Land related layers and CPCST .
65533	If the identified growing season(s) has not been flagged by the conditions in flag 65532, and the derived seasonality period extends beyond the defined timeframe for the main growing season, then it is not feasible to produce any Cropping Pattern layer in this situation.Used for all Cropping Pattern layers, except Fallow Land related layers and CPCST .
65534	This flag is utilized when none of the aforementioned flags are already in effect, indicating that no confidence can be calculated for the specific Cropping Patterns layer. The confidence assessment for the layers relies on the uncertainty of the satellite time series at the identified emergence and harvest date(s). If the uncertainty is deemed significant or impossible to calculate, this flag will be applied. If this situation happens to derive the confidence level on the harvest date(s), it will impact only those layers that utilized these specific harvest date(s), and similarly for the emergence date(s).In case of the CPFLDCL layer, this flag could also occur when the Fallow Land Duration over the years always entails the full calendar year. As a result, it is not possible to

Quality flag	Definition
	calculate a confidence without information on the confidence at the start or end point defined by a harvest event. Used for all Cropping Patterns Confidence Layers .
65535	Outside area

Table 7-7: Summary of the applicable quality flags of the HRL Croplands layers

Name of layer	Acronym	Classified feature	Class coding	Applicable quality flags
Crop types	CTY	Categorical label on crop type.	0: No cropland1110: Wheat1120: Barley1130: Maize1140: Rice1150: Other cereals1210: Fresh Vegetables1220: Dry Pulses1310: Potatoes1320: Sugar Beet1410: Sunflower1420: Soybeans1430: Rapeseed1440: Flax cotton and hemp2100: Grapes2200: Olives2310: Fruits2320: Nuts3100: Unclassified annual crop3200: Unclassified permanent crop65535: outside area	[0, 65535]
Crop types confidence	CTYCL	Probability as obtained from the confidence of the selected crop type class.	0-100: Probability expressed as a percentage.	[253,255]
Main Crop Harvest	CPMCH	The harvest date of the main (annual) crop seasons expressed in DOY (days of the year).	YYDOY where YY = last 2 digits of the year (e.g., 19 for 2019) and DOY is the day of the year (1-366).	[0, 65526, 65527, 65531, 65532, 65533, 65535]
Main Crop Harvest Confidence Layer	CPMCHCL	The confidence layer for the harvest date indicating the uncertainty (days) on the detected harvest date of the main (annual) crop.	Uncertainty in days (1-40)	[0, 65526, 65527, 65531, 65532, 65533, 65534, 65535]
Main Crop Emergence	CPMCE	The emergence date of the main (annual) crop expressed in DOY (day of year).	YYDOY where YY = last 2 digits of the year (e.g., 19 for 2019) and DOY is the day of the	[0, 65526, 65527, 65531, 65532, 65533, 65535]

Name of layer	Acronym	Classified feature	Class coding	Applicable quality flags
			year (1-366).	
Main Crop Emergence Confidence Layer	CPMCECL	The uncertainty of the emergence date of the main (annual) crop expressed in days.	Uncertainty in days (1-40)	[0, 65526, 65527, 65531, 65532, 65533, 65534, 65535]
Main Crop Duration	CPMCD	The duration (in days) on the season duration of the main (annual) crop.	Days (40-366)	[0, 65526, 65527, 65531, 65532, 65533, 65535]
Main Crop Duration Confidence Layer	CPMCDCL	The uncertainty (in days) on the season duration of the main (annual) crop.	Uncertainty in days (1-80)	[0, 65526, 65527, 65531, 65532, 65533, 65534, 65535]
Bare Soil Before	CPBSB	The bare soil period (in days) before the emergence of the main annual crop.	Days (1-295)	[0, 65526, 65527, 65529, 65531, 65532, 65533, 65535]
Bare Soil Before Confidence Layer	CPBSBCL	The uncertainty of the bare soil period (in days) before the emergence of the main annual crop.	Uncertainty in days (1-80)	[0, 65526, 65527, 65529, 65531, 65532, 65533, 65534, 65535]
Bare Soil After	CPBSA	The bare soil period (in days) after the harvest of the main annual crop.	Days (1-275)	[0, 65526, 65527, 65529, 65531, 65532, 65533, 65535]
Bare Soil After Confidence Layer	CPBSACL	The uncertainty of the bare soil period (in days) after the harvest of the main annual crop.	Uncertainty in days (1-80)	[0, 65526, 65527, 65529, 65531, 65532, 65533, 65534, 65535]
Secondary Crops Type	CPSCT	Indicates the presence of a secondary (cover crop) season and differentiates it into four possible categories.	1: Short Summer 2: Long Summer 3: Short Winter 4: Long Winter	[0, 65526, 65527, 65530, 65531, 65532, 65533, 65535]
Secondary Crops Emergence	CPSCE	The emergence date of the secondary crop in DOY.	YYDOY where YY = last 2 digits of the year (e.g., 19 for 2019) and DOY is the day of the year (1-366).	[0, 65526, 65527, 65530, 65531, 65532, 65533, 65535]
Secondary Crops Duration	CPSCD	The duration (in days) of the secondary crop season.	Days (40-366)	[0, 65526, 65527, 65530, 65531, 65532, 65533, 65535]
Secondary Crops Duration Confidence Layer	CPSCDCL	The uncertainty (in days) of the duration of the secondary of a secondary crop season.	Uncertainty in days (1-80)	[0, 65526, 65527, 65530, 65531, 65532, 65533, 65534, 65535]

Name of layer	Acronym	Classified feature	Class coding	Applicable quality flags
Fallow Land Presence	CPFLP	Yearly fallow land classification indicating if arable land has been left fallow in the respective calendar year.	1: Fallow land	[0, 65535]
Fallow Land Presence Confidence Layer	CPFLPCL	Uncertainty of the fallow presence classification	0-100: Probability expressed as percentage	[253, 65535]
Fallow Land Duration	CPFLD	The total fallow land duration (in days) over the period 2017-2021.	Days (1-1826)	[0,65535]
Fallow Land Duration Confidence Layer	CPFLDCL	The uncertainty (in days) of the total fallow land duration over the period 2017-2021.	Uncertainty in days (1-200)	[0, 65534, 65535]
Cropping Seasons Yearly	CPCSY	Number of growing seasons detected within 1 year.	[1-2]	[0, 65526, 65527,65531, 65532, 65533 65535]
Cropping Seasons Types over 3 years	CPCST	Indicating the number of different crop types grown in a 3-year period (excl. secondary crops)	[1-3]	[0, 65528, 65535]

8. Production quality assessment

The aim of this chapter is to inform about the procedures for internal validation and accuracy assessment for the status and change layer with full **EEA38** coverage. While the different layers have their own quality requirements, all have in common an assessment of the thematic quality which relies on comparing mapped information within the layers with reference data at selected locations.

This procedure contains 3 steps that will be described in the following sections:

- Sampling design
- Response design
- Statistical Analysis

The internal validation of the **HRL Croplands** layers follows scientifically accepted and operationally proven validation design, applied in previous productions of various HRL's of reference years 2012, 2015 and 2018. According to the product specifications, results will be presented in the form of **Overall Accuracies (OA)**, **Producer's** and **User's Accuracies** or **F-Score measure**.

8.1 Layers to be verified

While the full portfolio of the **HRL Croplands** product includes numerous layers and reference years, only a subset of them is concerned by the internal verification exercise. The focus of the assessment has been set on the primary layer being the **Crop Types** layer and the reference years 2018 and 2021. The rationale for this selection has been mainly the availability of essential reference datasets such as **LUCAS** 2018 [2], the VHR IMAGE coverages of 2018 and 2021 imagery, and publicly available GSAA datasets. An overview of the verified layers and their target accuracies is given in Table 8-1.

Table 8-1: Layers to be verified and target accuracies

Layers	Target accuracy
Crop Types 10m 2018	F-score > 85% at crop group level (6 classes)F-score > 80% at aggregation level 2 (11 classes)
Crop Types10m 2021	F-score > 85% at crop group level (6 classes)F-score > 80% at aggregation level 2 (11 classes)

8.2 Sampling Design

Two sampling approaches are considered to ensure an adequate accuracy assessment of the layers at different hierarchical nomenclature levels.

The **first sampling approach** is dedicated to assess the accuracies of the **Crop Types** layer at crop group level. At pan-European level, this corresponds to a non-stratified, systematic and random sampling approach based on the 2 km by 2 km LUCAS grid. For the assessment of status layers, 10 000 samples (Primary Sampling Units) were randomly selected over the extended LUCAS grid (Figure 8-1).



Figure 8-1: Approach I sample distribution for verification at Crop Group Level

The **second sampling approach** is dedicated to the accuracy assessment of the **Crop Types** layer at aggregation level 1 which contains 11 classes (Table 8-2). To ensure a correct and precise identification of crop parcels during the verification exercise, the use of LUCAS and GSAA data as reference data is mandatory. However, some limitations exist about those datasets:

- LUCAS survey information exists for the year 2018 but not for the year 2021.
- GSAA data is not openly available for all countries and for all reference years.

To circumvent those limitations, the approach II proposes to:

- For the year 2018, take advantage of the **LUCAS** survey information by using all “LUCAS crops” (e.g. LUCAS point with crop type information) which were not used as calibration/training data for the year 2018 (3226 LUCAS points when excluding **UK**). The location of the samples is illustrated in Figure 8-2.
- Use of **GSAA** data when and where available: per country, a random sampling of crop parcels (among those not used for calibration/training purposes) is performed to populate the reference database. This sampling is done on the year 2021 when GSAA was available but can be done on years 2020 or 2019 depending on GSAA availability for said country. This is illustrated

on Figure 8-3. The total amount of samples selected from GSAA data is similar to the sampling effort for the year 2018 when considering LUCAS crop sample (~ 3300 samples)




Figure 8-2: Approach II sample distribution depending on LUCAS points containing crop type information for the year 2018



It is important to note that those samplings do not cover the whole EEA38 due to the availability and specifications of GSAA and LUCAS data, so the reporting levels will differ from approach I.

The response design is the protocol used for retrieval of the validation/reference information for all sample units. Two types of datasets are used to perform the interpretation of samples units: guiding data and reference data.

Reference data are complementary and independent data that can provide more spatial details and landscape context:

- **VHR_IMAGE_2018⁴** and **VHR_IMAGE_2021⁵**: very high resolution optical earth observation imagery, covering **EEA38** for the reference years 2018 and 2021 (+1 year). Note that at time of this exercise, the VHR_IMAGE_2021 coverage is not 100% complete. These datasets comprise EO imagery between 2 and 4m spatial resolution.
- Other external datasets:
 - o **Bing maps image/** cartography layer
 - o **Open Street Map** data
 - o **Google Earth Image /** cartography data
 - o **LUCAS** data for the year 2018 (where available)
 - o **GSAA** data for the year 2018 and 2021 (where available)

The interpretation workflow for the verification of the crop group consists of thematic plausibility analysis of the sample units. Plausibility analysis means that the class assigned by the layer is known by the interpreter during the interpretation. The Primary Sampling Units were interpreted at the crop group level (Table 8-2) using guiding and reference datasets, considering the surrounding of the sample (typically crop parcel) to properly account for the MMU of 0.25 ha.

Table 8-2: Crop Types layer nomenclature and aggregation levels

Class code	Land Cover	Crop Group	Level 1(accuracy aggregation)	Crop types (Level 2)
1110	Arable Crops	Cereals	Cereals	Wheat
1120	Arable Crops	Cereals	Cereals	Barley
1130	Arable Crops	Cereals	Maize	Maize
1140	Arable Crops	Cereals	Rice	Rice
1150	Arable Crops	Cereals	Cereals	Other cereals
1210	Arable Crops	Dry pulses & Vegetables	Dry Pulses, Fresh vegetables and Potatoes	Fresh Vegetables
1220	Arable Crops	Dry pulses & Vegetables	Dry Pulses, Fresh vegetables and Potatoes	Dry pulses
1310	Root/tuber crops	Dry Pulses, Fresh vegetables and Potatoes	Potatoes	
1320	Root/tuber crops	Sugar beet	Sugar Beet	
1410	Non-permanent industrial crops	Sunflower	Sunflower	
1420	Non-permanent industrial crops	Soybeans	Soybeans	
1430	Non-permanent industrial crops	Rapeseed	Rapeseed	
1440	Non-permanent industrial crops	Flax, cotton and hemp	Flax, cotton and hemp	

⁴ [Copernicus Data Space Ecosystem – Copernicus Contributing Missions – VHR IMAGE 2018](#)

⁵ [Copernicus Data Space Ecosystem – Copernicus Contributing Missions – VHR IMAGE 2021](#)

Class code	Land Cover	Crop Group	Level 1(accuracy aggregation)	Crop types (Level 2)
2100	Permanent Crops	Permanent Crops	Permanent crops	Grapes
2200	Permanent Crops	Permanent Crops	Permanent crops	Olives
2310	Permanent Crops	Permanent Crops	Permanent crops	Fruits
2320	Permanent Crops	Permanent Crops	Permanent crops	Nuts
3100	Arable Crops	Unclassified arable crop	Unclassified arable crop	Unclassified arable crop
3200	Permanent Crops	Unclassified permanent crop	Unclassified permanent crop	Unclassified permanent crop

For layers where the second sampling design has been applied (see 8.2) and the use of **LUCAS** and **GSAA** data were used, the response design is simplified. No visual interpretation is required as the reference class is extracted from LUCAS/GSAA data to be compared to the **Crop Types** layer without any additional process.

8.4 Statistical Analysis

The thematic accuracy level or the **Crop Types** layer is defined in terms of F-score, which can be derived from confusion matrices. To this end, the row and column totals and the diagonal of the matrix are used to assess two further types of accuracy, the User's and Producer's Accuracy:

- **Producer's Accuracy (PA)** for a given class =

$$a_{ii} / C_i$$

, representing an (inversely proportional) measure of Omission Error. For instance, an observation has been identified as tree-covered in the validation dataset but was actually classified as another class: it has been omitted from the target class.

- **User's Accuracy (UA)** for a given class =

$$a_{ii} / R_i$$

, representing an (inversely proportional) measure of the Commission Error (or contamination risk), i.e. errors due to the wrong allocation of an observation (i.e. mapped landcover) to a landcover class. For instance, an observation is classified as broadleaved tree cover, but identified as belonging to another class during the validation process: this observation has contaminated another class.

F-score is a measure of accuracy calculated as the harmonic mean of the Producer's and the User's Accuracies calculated as:

$$F_1 = 2 \cdot \frac{(PA \cdot UA)}{(PA + UA)}$$

For both **Crop Types** layers 2018 and 2021 the F1 scores are computed derived for each class of crop group level (6 classes) and aggregation level 1 (11 classes) (Table 8-2).

8.5 Verification Results

The assessed accuracies for the **Crop Types** layers are presented at the crop group level (gridbased random sampling) and the aggregation level 1 (11 classes, opportunistic sampling from LUCAS and GSAA data). For the rationale on the different sampling designs please refer to section 8.2)

8.5.1 Crop Group level

Accuracy figures for crop group level are presented in the confusion matrices shown for the EU27 in Table 8-4 and the EEA38 in Table 8-5

At the crop group level the layers should reach an F1-score of 85% for each class. This is the case for all classes for both reference years 2018 and 2021 at EU27 level, even crop types which present a lower number of samples such as root/tuber or dry pulses and vegetables. At EEA38 level, the results are similar with the exception of permanent crops class that reach accuracies slightly below the 85% threshold (83-84%).

Table 8-3 Crop group level class codes

Class code	Crop Group
11	Cereals
12	Dry pulses & Vegetables
13	Root/tuber crops
14	Non-permanent industrial crops
20	Permanent Crops
30	Unclassified crop

Table 8-4 HRL Crop Types accuracies for crop groups, EU27 (for the class codes see Table 8-3)

8.5.2 CTY2018 - EU27 (Weighted)

Product	Ref. 0	Ref. 11	Ref. 12	Ref. 13	Ref. 14	Ref. 20	Ref. 30	Total
0	5156.57	164.00	2.01	0.01	2.00	21.02	1.00	5346.60
11	2.00	816.07						818.07
12			63.01					63.01
13				51.00				51.00
14	1.00	1.00			203.01	1.00		206.01
20	10.01	4.00				160.46	1.00	175.48
30	3.00	25.00				1.00	8.04	37.04
Total	5172.58	1010.07	65.02	51.01	205.01	183.49	10.04	6697.22

	0	11	12	13	14	20	30
Prod. Acc.	99.69%	80.79%	96.91%	99.99%	99.02%	87.45%	80.08%
User Acc.	96.45%	99.76%	100.00%	100.00%	98.54%	91.45%	21.71%
F-score	98.04%	89.28%	98.43%	99.99%	98.78%	89.40%	34.15%

8.5.3 CTY2018 - EU27 (Unweighted)

Product	Ref. 0	Ref. 11	Ref. 12	Ref. 13	Ref. 14	Ref. 20	Ref. 30	Total
0	8047	164	3	1	2	25	1	8243
11	3	833						836
12			65					65
13				51				51
14	1	1			206	1		209
20	13	4				224	1	242
30	3	25				2	15	45
Total	8067	1027	68	52	208	252	17	9691

	0	11	12	13	14	20	30
Prod. Acc.	99.75%	81.11%	95.59%	98.08%	99.04%	88.89%	88.24%
User Acc.	97.62%	99.64%	100.00%	100.00%	98.56%	92.56%	33.33%
F-score	98.68%	89.43%	97.74%	99.03%	98.80%	90.69%	48.39%

8.5.4 CTY2021 - EU27 (Weighted)

Product	Ref. 0	Ref. 11	Ref. 12	Ref. 13	Ref. 14	Ref. 20	Ref. 30	Total
0	5127.29	133.02	0.01	0.00	2.00	19.04	1.00	5282.36
11	6.01	898.75				0.01		904.77
12	1.00		72.16					73.16
13				49.06				49.06
14	3.00	1.00			197.19			201.19
20	10.02	2.01				164.56	0.00	176.59
30	1.00	5.00					6.07	12.07
Total	5148.32	1039.78	72.17	49.06	199.19	183.61	7.07	6699.20

	0	11	12	13	14	20	30
Prod. Acc.	99.59%	86.44%	99.99%	100.00%	99.00%	89.63%	85.80%
User Acc.	97.06%	99.33%	98.63%	100.00%	98.01%	93.19%	50.29%
F-score	98.31%	92.44%	99.31%	100.00%	98.50%	91.37%	63.41%

8.5.5 CTY2021 - EU27 (Unweighted)

Product	Ref. 0	Ref. 11	Ref. 12	Ref. 13	Ref. 14	Ref. 20	Ref. 30	Total
0	7029.00	140.00	1.00	1.00	2.00	24.00	2.00	7199.00
11	10.00	1676.00				2.00		1688.00
12	1.00		134.00					135.00
13				76.00				76.00
14	3.00	2.00			279.00			284.00
20	16.00	3.00				256.00	1.00	276.00

Product	Ref. 0	Ref. 11	Ref. 12	Ref. 13	Ref. 14	Ref. 20	Ref. 30	Total
30		1.00					26.00	32.00
Total	7060.00	1826.00	135.00	77.00	281.00	282.00	29.00	9690.00

	0	11	12	13	14	20	30
Prod. Acc.	99.56%	91.79%	99.26%	98.70%	99.29%	90.78%	89.66%
User Acc.	97.64%	99.29%	99.26%	100.00%	98.94%	92.75%	81.25%
F-score	98.59%	95.39%	99.26%	99.35%	98.76%	91.76%	85.25%

8.5.6 CTY2018 - EEA38 (Weighted)

Product	Ref. 0	Ref. 11	Ref. 12	Ref. 13	Ref. 14	Ref. 20	Ref. 30	Total
0	6810.95	188.08	3.01	0.01	31.00	42.06	1.00	7076.11
11	20.02	960.10		0.00	18.00			998.12
12	2.00	3.00	93.04		1.00			99.04
13		1.00		52.00				53.00
14	1.00	13.00			226.01	1.00		241.01
20	18.02	5.00			2.00	179.51	1.00	205.54
30	5.00	26.01	1.00		4.00	1.00	8.04	45.05
Total	6857.00	1196.19	97.05	52.01	282.02	223.56	10.04	8717.87

	0	11	12	13	14	20	30
Prod. Acc.	99.33%	80.26%	95.87%	99.98%	80.14%	80.29%	80.09%
User Acc.	96.25%	96.19%	93.94%	98.11%	93.78%	87.34%	17.85%
F-score	97.77%	87.51%	94.89%	99.04%	86.42%	83.67%	29.20%

8.5.7 CTY2018 - EEA38 (Unweighted)

Product	Ref. 0	Ref. 11	Ref. 12	Ref. 13	Ref. 14	Ref. 20	Ref. 30	Total
0	10578	219	6	1	32	60	1	10897.00
11	27	989		1	18			1035.00
12	2	4	103		1			110.00
13		1		53				54.00
14	1	13			230	1		245.00
20	26	7			2	257	1	293.00
30	5	28	1		4	2	16	56.00
Total	10639	1261	110	55	287	320	18	12690.00

	0	11	12	13	14	20	30
Prod. Acc.	99.43%	78.43%	93.64%	96.36%	80.14%	80.31%	88.89%
User Acc.	97.07%	95.56%	93.64%	98.15%	93.88%	87.71%	28.57%

	0	11	12	13	14	20	30
F-score	98.24%	86.15%	93.64%	97.25%	86.47%	83.85%	43.24%

8.5.8 CTY2021 - EEA38 (Weighted)

Product	Ref. 0	Ref. 11	Ref. 12	Ref. 13	Ref. 14	Ref. 20	Ref. 30	Total
0	6784.24	157.07	1.01	0.01	23.00	48.07	1.01	7014.40
11	22.07	1041.09	6.00		26.00	0.01		1095.17
12	1.01	4.02	82.18		2.00		1.00	90.21
13	1.00			52.07	1.00		0.00	54.07
14	4.00	20.01	2.00		216.22	1.01		243.24
20	17.02	3.02			2.00	179.61	0.00	201.65
30	2.02	8.00	2.00		3.00		8.07	23.09
Total	6831.37	1233.21	93.19	52.08	273.22	228.70	10.09	8721.85

	0	11	12	13	14	20	30
Prod. Acc.	99.31%	84.42%	88.18%	99.99%	79.14%	78.54%	80.01%
User Acc.	96.72%	96.05%	91.10%	96.30%	88.89%	89.07%	39.96%
F-score	98.00%	89.43%	89.62%	98.11%	83.73%	83.47%	48.66%

8.5.9 CTY2021 - EEA38 (Unweighted)

Product	Ref. 0	Ref. 11	Ref. 12	Ref. 13	Ref. 14	Ref. 20	Ref. 30	Total
0	9386.00	186.00	4.00	2.00	23.00	64.00	5.00	9670.00
11	52.00	1952.00	6.00		26.00	4.00		2040.00
12	6.00	11.00	151.00		2.00		2.00	172.00
13	1.00			84.00	1.00		1.00	87.00
14	5.00	26.00	2.00		310.00	3.00		346.00
20	25.00	8.00			2.00	288.00	1.00	324.00
30	8.00	9.00	2.00		3.00		30.00	52.00
Total	9483.00	2192.00	165.00	86.00	367.00	359.00	39.00	12691.00

	0	11	12	13	14	20	30
Prod. Acc.	98.98%	89.05%	91.52%	97.67%	84.47%	80.22%	76.92%
User Acc.	97.06%	95.69%	87.79%	96.55%	89.60%	88.89%	57.69%
F-score	98.01%	92.25%	89.61%	97.11%	86.96%	84.33%	65.93%

8.5.10 Aggregation level 1

Accuracy figures for Crop Types layers at Aggregation Level 1 are presented in Table 8-6 below, according to the comparisons with LUCAS and GSSA data.

Target accuracy for this level of analysis is an F-Score of 80% for each of the classes. While some crop types exceed the 80% threshold (cereals, maize, rapeseed), other fail to reach it. The rarity of some

classes explains the few numbers of samples concerned and therefore can imply that lower F-score should be taken with caution (flax, cotton, hemp, soybean, and rice classes). Unclassified classes have a null F-score due to their nature, as those classes do not exist in the LUCAS or GSAA reference dataset.

Table 8-6: Accuracies for aggregation level-1 crop types

		All GSAA data (2021-2020-2019)	All GSAA data (2021-2020-2019)	2018 (LUCAS)	2018 (LUCAS)
Class	Code	Nb Samples (according to GSAA)	F-score	Nb Samples(accordin g to LUCAS)	F-score
Other Cereals	11	921	0.850	1571	0.845
Maize	1130	400	0.879	458	0.822
Rice	1140	0	0.000	2	0.000
Dry pulses, Fresh vegetables & potatoes	12	193	0.654	185	0.667
Sugar beet	1320	17	0.686	70	0.915
Sunflower	1410	36	0.771	130	0.792
Soybeans	1420	20	0.654	22	0.500
Rapeseed	1430	84	0.804	148	0.830
Flax, cotton and hemp	1440	4	0.500	10	0.778
Permanent crops	20	300	0.735	385	0.673
Unclassified arable crops	3100	0	0.000	0	0.000
Unclassified permanent crops	3200	0	0.000	0	0.000

9. Terms of use and product technical support

9.1 Terms of use

The Terms of Use for the product(s) described in this document acknowledge the following:

Free, full and open access to the products and services of the Copernicus Land Monitoring Service is made on the conditions that:

1. When distributing or communicating Copernicus Land Monitoring Service products and services (data, software scripts, web services, user and methodological documentation and similar) to the public, users shall inform the public of the source of these products and services and shall acknowledge that the Copernicus Land Monitoring Service products and services were produced “with funding by the European Union”.
2. Where the Copernicus Land Monitoring Service products and services have been adapted or modified by the user, the user shall clearly state this.
3. Users shall make sure not to convey the impression to the public that the user’s activities are officially endorsed by the European Union.

The user has all intellectual property rights to the products he/she has created based on the Copernicus Land Monitoring Service products and services.

Consult [Data policy — Copernicus Land Monitoring Service](#) for further details.

9.2 Citation

When planning a publication (scientific, commercial, etc.), it shall explicitly mention:

“This publication has been prepared using European Union’s Copernicus Land Monitoring Service information; <insert all relevant DOI links here, if applicable>”

When developing a product or service using the products or services of the Copernicus Land Monitoring Service, it shall explicitly mention:

“Generated using European Union’s Copernicus Land Monitoring Service information; <insert all relevant DOI links here, if applicable>”

When redistributing a part of the Copernicus Land Monitoring Service (product, dataset, documentation, picture, web service, etc.), it shall explicitly mention:

“European Union’s Copernicus Land Monitoring Service information; <insert all relevant DOI links here, if applicable>”

Consult [Data policy — Copernicus Land Monitoring Service](#) for further details.

9.3 Product technical support

Product technical support is provided by the product custodian through [Copernicus Land Monitoring Service – Service desk](#). Product technical support does not include software specific user support or general GIS or remote sensing support.

More information on the products can be found on the Copernicus Land Monitoring Service website (<https://land.copernicus.eu/>).

10. List of Abbreviations & Acronyms

Abbreviation	Name
ATBD	Algorithm Theoretical Basis Document
BS	Bare Soil
BVL	Base Vegetation Layer
CAP	Common Agricultural Policy
CL	Confidence Layer
CLC	CORINE Land Cover
CLMS	Copernicus Land Monitoring Service
CLS	Collecte Localisation Satellites
COG	Cloud-Optimized GeoTIFFs
CORINE	Coordination of information on the environment
CP	Cropping Patterns
CPBSB	Bare Soil Before
CPBSBCL	Bare Soil Before Confidence Layer
CPBSA	Bare Soil After
CPBSACL	Bare Soil After Confidence Layer
CPCST	Cropping Seasons Types over 3 years
CPCSY	Cropping Seasons Yearly
CPFLD	Fallow Land Duration
CPFLDCL	Fallow Land Duration Confidence Layer
CPFLP	Fallow Land Presence
CPFLPCL	Fallow Land Presence Confidence Layer
CPMCD	Main Crop Duration
CPMCDCL	Main Crop Duration Confidence Layer
CPMCE	Main Crop Emergence
CPMCECL	Main Crop Emergence Confidence Layer
CPMCH	Main Crop Harvest
CPMCHCL	Main Crop Harvest Confidence Layer
CPSCT	Secondary Crops Type
CPSD	Secondary Crops Duration
CPSCDCL	Secondary Crops Duration Confidence Layer
CPSE	Secondary Crops Emergence
CTY	Crop Types
CTYCL	Crop Types Confidence Layer
DLT	Dominant Leaf Type
DLTC	Dominant Leaf Type Change
DOY	Day Of Year
EAGLE	EIONET Action Group on Land monitoring in Europe
ECOLaSS	Evolution of Copernicus Land Services based on Sentinel Data
EEA	European Environment Agency

Abbreviation	Name
EEA38	The 32 member and 6 cooperating countries of the EEA
EIONET	European Environment Information and Observation Network
EO	Earth Observation
EU	European Union
EU27	The 27 member states of the EU
fAPAR	Fraction of absorbed photosynthetically active radiation
FLP	Fallow Land Presence
FML	Forest Monitoring Law
GIS	Geographic Information System
GSAA	GeoSpatial Aid Application
H2020	Horizon2020
HER	Herbaceous Cover Layer
HR	High Resolution
HRL / HRLs	High Resolution Layer / High Resolution Layers
HRL VLCC	High Resolution Layer – Vegetated Land Cover Characteristics
ID	Identification Number
IPCC	Intergovernmental Panel on Climate Change
JRC	Joint Research Centre
LAEA	Lambert Azimuthal Equal Area projection
LU	Land Use
LUCAS	Land Use / Cover Area frame Survey
LULUCF	Land Use, Land Use Change and Forestry
MC	Main Crops
MMU	Minimum Mapping Unit
OA	Overall Accuracy
PA	Producer Accuracy
PAM	Permanent Auxiliary metadata
SC	Secondary Crops
TCD	Tree Cover Density
TI	Thünen-Institut (Federal Research Institute for Rural Areas, Forestry and Fisheries)
UA	User's Accuracy
UK	United Kingdom
USDA	U.S. Department of Agriculture
UTM	Universal Transverse Mercator
VHR	Very High Resolution
VITO	Vlaamse Instelling voor Technologisch Onderzoek
XML	Extensible Markup Language

11. References

- [1] T. Duan et al., NGBoost: Natural Gradient Boosting for Probabilistic Prediction (2020), ICML 2020
- [2] Eurostat. (2018). LUCAS Survey 2018. [Data set]. <https://ec.europa.eu/eurostat/web/lucas/data/primary-data>.
- [3] Selkowitz, D. J., & Stehman, S. V. (2011). Thematic accuracy of the National Land Cover Database (NLCD) 2001 land cover for Alaska. *Remote Sensing of Environment*, 115(6), 1401-1407.
- [4] Olofsson, P., Foody, G. M., Stehman, S. V., & Woodcock, C. E. (2013). Making better use of accuracy data in land change studies: Estimating accuracy and area and quantifying uncertainty using stratified estimation. *Remote Sensing of Environment*, 129, 122-131.
- [5] ECoLaSS (2019). Deliverable D3.2 - Service Evolution Requirements Report Vol. 2 https://www.ecolass.eu/_files/ugd/c90769_5a431f06039141a6b4db4d6b4596d272.pdf (accessed 28 October 2024)
- [6] Nextspace (2018). Work performed by the Nextspace consortium – Observation Requirements (February 2018); <https://www.copernicus.eu/en/documentation/technicaldocuments/technical-documents> (accessed 28 October 2024)
- [7] CLMS Vegetated HRL ATBD (2024). HRL Algorithm Technical Basis Document

12. Annex I – Colour tables for HRL Croplands

Class Code	Class Name	Red	Green	Blue	
0	No Cropland	240	240	240	
1110	Wheat	238	110	50	
1120	Barley	251	162	74	
1130	Maize	250	220	20	
1140	Rice	233	67	1	
1150	Other Cereals	232	169	149	
1210	Fresh Vegetables	174	199	232	
1220	Dry Pulses	72	151	191	
1310	Potatoes	201	140	67	
1320	Sugar Beet	156	91	12	
1410	Sunflower	255	121	121	

1420	Soybeans	168	106	150	
1430	Rapeseed	227	119	194	
1440	Flax, cotton and hemp	247	182	210	
2100	Grapes	219	219	141	
2200	Olives	193	206	18	
2310	Fruits	121	160	58	
2320	Nuts	90	124	48	
3100	Unclassified arable crop	215	215	215	
3200	Unclassified permanent crop	171	171	171	
65535	Outside area	255	255	255	

Table 0-1: Colour palette and attributes of HRL Crop Types (CTY) layer

Class Code	Class Name	Red	Green	Blue	
0	0% classification confidence	255	0	0	
1-49	1-49% classification confidence	colour shades in between			
50	50% classification confidence	255	255	0	
51-99	51-99% classification confidence	colour shades in between			
100	100% classification confidence	8	99	0	
254	No cropland	240	240	240	
255	Outside area	0	0	0	

Table 0-2: Colour palette and attributes of CTYCL layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
17090	Apr (product year)	255	128	159	
17120	May (product year)	255	176	127	
17151	Jun (product year)	255	223	128	
17181	Jul (product year)	223	255	128	
17212	Aug (product year)	159	255	128	
17243	Sep (product year)	109	220	141	
17273	Oct (product year)	128	255	223	
17304	Nov (product year)	128	223	255	
17334	Dec (product year)	127	189	255	
18365	Jan (product year+1)	64	0	255	
65526	65526	225	225	225	
65527	65527	200	200	200	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65535	Outside area	0	0	0	

Table 0-3: Colour palette and attributes of CPMCH layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
1	1 day uncertainty	202	0	32	
1-19	1-19 days uncertainty	Colour shades in between			
20	20 days uncertainty	247	243	241	
21-39	21-39 days uncertainty	Colour shades in between			
40	40 days uncertainty	5	113	176	
65526	65526	225	225	225	
65527	65527	200	200	200	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-4: Colour palette and attributes of CPMCHCL layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
16214	Aug (product year-1)	72	135	67	
16245	Sep (product year-1)	101	204	129	
16276	Oct (product year-1)	125	252	200	
16306	Nov (product year-1)	128	255	223	
16336	Dec (product year-1)	128	223	255	
16366	Jan (product year)	159	127	255	
17031	Feb (product year)	221	127	255	
17059	Mar (product year)	255	127	223	

17120	Apr (product year)	255	127	161	
17151	May (product year)	255	176	127	
17181	Jun (product year)	255	223	128	
17212	Jul (product year)	223	255	128	
17243	Aug (product year)	159	255	128	
17273	Sep (product year)	109	220	141	
65526	65526	225	225	225	
65527	65527	200	200	200	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65535	Outside area	0	0	0	

Table 0-5: Colour palette and attributes of CPMCE layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
1	1 day uncertainty	202	0	32	
1-19	1-19 days uncertainty	Colour shades in between			
20	20 days uncertainty	247	243	241	
21-39	21-39 days uncertainty	Colour shades in between			
40	40 days uncertainty	5	113	176	
65526	65526	225	225	225	
65527	65527	200	200	200	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-6: Colour palette and attributes of CPMCECL layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
40	40 days	215	25	28	
41-189	41-189 days	Colour shades in between			
190	190 days	255	243	178	
191-365	191-365 days	Colour shades in between			
366	366 days	26	150	65	
65526	65526	225	225	225	
65527	65527	200	200	200	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65535	Outside area	0	0	0	

Table 0-7: Colour palette and attributes of CPMCD layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
1	1	202	0	32	
2-29	2-29 days uncertainty	Colour shades in between			
30	30 days uncertainty	247	245	243	
31-59	31-59 days uncertainty	Colour shades in between			
60	>60 days uncertainty	5	113	176	
65526	65526	225	225	225	
65527	65527	200	200	200	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-8: Colour palette and attributes of CPMCDCL layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
10	10 days	214	44	58	
11-74	11-74 days	Colour shades in between			
75	75 days	247	247	247	
76-149	76-149 days	Colour shades in between			
150	> 150 days	5	113	176	
65526	65526	225	225	225	
65527	65527	200	200	200	
65529	65529	150	150	150	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-9: Colour palette and attributes of CPBSB layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
1	1	202	0	32	
2-29	2-29 days uncertainty	Colour shades in between			
30	30 days uncertainty	247	245	243	
31-59	31-59 days uncertainty	Colour shades in between			
60	> 60 days uncertainty	5	113	176	
65526	65526	225	225	225	

65527	65527	200	200	200	
65529	65529	150	150	150	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-10: Colour palette and attributes of CPBSBCL layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
10	10 days	214	44	58	
11-74	11-74 days	Colour shades in between			
75	75 days	247	247	247	
76-149	76-149 days	Colour shades in between			
150	>150 days	5	113	176	
65526	65526	225	225	225	
65527	65527	200	200	200	
65529	65529	150	150	150	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-11: Colour palette and attributes of CPBSA layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
1	1	202	0	32	
2-29	2-29 days uncertainty	Colour shades in between			
30	30 days uncertainty	247	245	243	
31-59	31-59 days uncertainty	Colour shades in between			
60	>60 days uncertainty	5	113	176	
65526	65526	225	225	225	
65527	65527	200	200	200	
65529	65529	150	150	150	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-12: Colour palette and attributes of CPBSBCL layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
1	Short summer CC	254	203	91	
2	Long summer CC	255	198	1	
3	Short winter CC	99	229	59	
4	Long winter CC	16	100	0	
65526	65526	225	225	225	
65527	65527	200	200	200	
65530	65530	125	125	125	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-13: Colour palette and attributes of CPSCT layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
16366	Jan (product year)	159	127	255	
17031	Feb (product year)	221	127	255	
17059	Mar (product year)	255	127	223	
17090	Apr (product year)	255	127	161	
17120	May (product year)	255	176	127	
17151	Jun (product year)	255	223	128	
17181	Jul (product year)	223	255	128	
17212	Aug (product year)	159	255	128	
17243	Sep (product year)	109	220	141	
17273	Oct (product year)	128	255	223	
17304	Nov (product year)	128	223	255	
17334	Dec (product year)	127	189	255	
65526	65526	222	225	225	
65527	65527	200	200	200	
65530	65530	125	125	125	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65535	Outside area	0	0	0	

Table 0-14: Colour palette and attributes of CPSCE layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
40	40 days	215	25	28	
41-149	41-149 days	Colour shades in between			
150	150 days	247	252	189	
151-199	151-200 days	Colour shades in between			
200	200 days	165	217	165	
201-249	201-249 days	Colour shades in between			
250	>250 days	43	131	186	
65526	65526	225	225	225	
65527	65527	200	200	200	
65530	65530	125	125	125	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65535	Outside area	0	0	0	

Table 0-15: Colour palette and attributes of CPSCD layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
1	1	202	0	32	
2-29	2-29 days uncertainty	Colour shades in between			
30	30 days uncertainty	247	245	243	
31-59	31-59 days uncertainty	Colour shades in between			
60	>60 days uncertainty	5	113	176	
65526	65526	225	225	225	
65527	65527	200	200	200	
65530	65530	125	125	125	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-16: Colour palette and attributes of CPSCDCL layer

Class Code	Class Name	Red	Green	Blue	
0	No fallow land	240	240	240	
1	Fallow	206	139	39	
65535	Outside area	0	0	0	

Table 0-17: Colour palette and attributes of CPFLP layer

Class Code	Class Name	Red	Green	Blue	
0	0% confidence	202	0	32	
1-49	1-49% confidence	Colour shades in between			
50	50% confidence	247	247	247	
51-99	51-99% confidence	Colour shades in between			
100	100% confidence	5	113	176	
253	No fallow land	240	240	240	
65535	Outside area	0	0	0	

Table 0-18: Colour palette and attributes of CPFLPCL layer

Class Code	Class Name	Red	Green	Blue	
0	No fallow land	240	240	240	
40	40 days	215	25	28	
41-948	41-948 days	Colour shades in between			
949	949 days	248	252	185	
950-1825	950-1825 days	Colour shades in between			
1826	1826 days	26	150	65	
65535	Outside area	0	0	0	

Table 0-19: Colour palette and attributes of CPFLD layer

Class Code	Class Name	Red	Green	Blue	
0	No fallow land	240	240	240	
1	1 day uncertainty	202	0	32	
1-19	1-19 days uncertainty	Colour shades in between			
20	20 days uncertainty	247	243	241	
21-39	21-39 days uncertainty	Colour shades in between			
40	>40 days uncertainty	5	113	176	
65534	65534	179	48	179	
65535	Outside area	0	0	0	

Table 0-20: Colour palette and attributes of CPFLDCL layer

Class Code	Class Name	Red	Green	Blue	
0	No annual cropland	240	240	240	
1	1 growing season	231	157	29	
2	2 growing seasons	138	233	55	
65526	65526	225	225	225	
65527	65527	200	200	200	
65531	65531	100	100	100	
65532	65532	75	75	75	
65533	65533	126	52	107	
65535	Outside area	0	0	0	

Table 0-21: Colour palette and attributes of CPCS layer







Class Code	Class Name	Red	Green	Blue	
0	No annual cropland (over 3-years period)	240	240	240	
1	1 annual crop type over three years period	231	157	29	
2	2 annual crop types over three years period	138	233	55	
3	3 annual crop types over three years period	243	58	252	
65528	65528	175	175	175	
65535	Outside area	0	0	0	

Table 0-22: Colour palette and attributes of CPCST layer

13. Document history:

Record of changes made to the document over time after the initial published version, including version numbers, dates of revisions, and a summary of modifications. This table must be only included in the initial PUM published version and without the personal information ("Created by:" filed must be removed).

Revision	Date	Short description of changes
1.0	2025-03-28	Initial published issue