

# POLYTROPIC MODELS OF WHITE DWARFS

## UNC PHYS 331 PROJECT

Erin Conn    Matthew Hurley

April 14, 2014

# TABLE OF CONTENTS

## THEORY

Polytropes

White Dwarfs

## METHODS

## RESULTS

## DISCUSSION

## REFERENCES

## TABLE OF CONTENTS

## THEORY

Polytropes

White Dwarfs

## METHODS

## RESULTS

## DISCUSSION

## REFERENCES

# WHAT ARE POLYTROPES?

# WHAT ARE POLYTROPES?

Solutions to...

## The Lane-Emden Equation

$$\frac{1}{\xi^2} \frac{d}{d\xi} \left( \xi^2 \frac{d\theta}{d\xi} \right) = -\theta^n(\xi)$$

A dimensionless, 2nd order nonlinear differential equation relating the pressure of a spherically-symmetric gas distribution to the radius.

# WHY POLYTROPES?

- Provide simplified stellar models - simple pressure/density relation

# WHY POLYTROPES?

- ▶ Provide simplified stellar models - simple pressure/density relation
- ▶ Easier to solve than full equations of stellar structure

# WHY POLYTROPES?

- ▶ Provide simplified stellar models - simple pressure/density relation
- ▶ Easier to solve than full equations of stellar structure
- ▶ Require less computational effort - some analytic solutions even exist!



# DEFINITIONS

## DEFINITION

**Polytropic fluid** - Fluid with an equation of state that depends on only one variable

# DEFINITIONS

## DEFINITION

**Polytropic fluid** - Fluid with an equation of state that depends on only one variable

## DEFINITION

**Polytropic index** - Constant that relates pressure of a polytropic fluid to its volume (density). It may be any real number.

# DEFINITIONS

## DEFINITION

**Polytropic fluid** - Fluid with an equation of state that depends on only one variable

## DEFINITION

**Polytropic index** - Constant that relates pressure of a polytropic fluid to its volume (density). It may be any real number.

## DEFINITION

**Poisson's equation** Relates a force density function to a potential field

$$\nabla^2\Phi = f$$

## DERIVATION 1: POISSON EQUATION

Can be derived multiple ways. From laws of mass conservation and hydrostatic equilibrium:

## DERIVATION 1: POISSON EQUATION

Can be derived multiple ways. From laws of mass conservation and hydrostatic equilibrium:

$$dM(r) = 4\pi r^2 \rho(r) dr \rightarrow \frac{dM(r)}{dr} = 4\pi r^2 \rho(r)$$

## DERIVATION 1: POISSON EQUATION

Can be derived multiple ways. From laws of mass conservation and hydrostatic equilibrium:

$$\begin{aligned} dM(r) &= 4\pi r^2 \rho(r) dr \rightarrow \frac{dM(r)}{dr} = 4\pi r^2 \rho(r) \\ \frac{dP(r)}{dr} &= -\frac{\rho(r)GM(r)}{r^2} \end{aligned}$$

## DERIVATION 1: POISSON EQUATION

Can be derived multiple ways. From laws of mass conservation and hydrostatic equilibrium:

$$\begin{aligned} dM(r) &= 4\pi r^2 \rho(r) dr \rightarrow \frac{dM(r)}{dr} = 4\pi r^2 \rho(r) \\ \frac{dP(r)}{dr} &= -\frac{\rho(r)GM(r)}{r^2} \end{aligned}$$

These equations are related by multiplying the hydrostatic equation by  $r^2/\rho$  and differentiating:

## DERIVATION 1: POISSON EQUATION

Can be derived multiple ways. From laws of mass conservation and hydrostatic equilibrium:

$$\begin{aligned} \frac{dM(r)}{dr} &= 4\pi r^2 \rho(r) \\ \frac{dP(r)}{dr} &= -\frac{\rho(r)GM(r)}{r^2} \end{aligned} \rightarrow \frac{dM(r)}{dr} = 4\pi r^2 \rho(r)$$

These equations are related by multiplying the hydrostatic equation by  $r^2/\rho$  and differentiating:

$$\frac{d}{dr} \left( \frac{r^2}{\rho(r)} \frac{dP(r)}{dr} \right) = -G \frac{dM(r)}{dr}$$



## DERIVATION 1: POISSON EQUATION

Can be derived multiple ways. From laws of mass conservation and hydrostatic equilibrium:

$$\begin{aligned} \frac{dM(r)}{dr} &= 4\pi r^2 \rho(r) \\ \frac{dP(r)}{dr} &= -\frac{\rho(r)GM(r)}{r^2} \end{aligned} \rightarrow \frac{dM(r)}{dr} = 4\pi r^2 \rho(r)$$

These equations are related by multiplying the hydrostatic equation by  $r^2/\rho$  and differentiating:

$$\frac{d}{dr} \left( \frac{r^2}{\rho(r)} \frac{dP(r)}{dr} \right) = -G \frac{dM(r)}{dr}$$

Yielding Poisson's equation for gravity:

## DERIVATION 1: POISSON EQUATION

Can be derived multiple ways. From laws of mass conservation and hydrostatic equilibrium:

$$\begin{aligned} \frac{dM(r)}{dr} &= 4\pi r^2 \rho(r) \\ \frac{dP(r)}{dr} &= -\frac{\rho(r)GM(r)}{r^2} \end{aligned} \rightarrow \frac{dM(r)}{dr} = 4\pi r^2 \rho(r)$$

These equations are related by multiplying the hydrostatic equation by  $r^2/\rho$  and differentiating:

$$\frac{d}{dr} \left( \frac{r^2}{\rho(r)} \frac{dP(r)}{dr} \right) = -G \frac{dM(r)}{dr}$$

Yielding Poisson's equation for gravity:

$$\frac{1}{r^2} \frac{d}{dr} \left( \frac{r^2}{\rho(r)} \frac{dP(r)}{dr} \right) = -4\pi G \rho(r)$$

## DERIVATION 2: WORKING TOWARDS A DIMENSIONLESS FORM

Define a polytropic state equation:

## DERIVATION 2: WORKING TOWARDS A DIMENSIONLESS FORM

Define a polytropic state equation:

$$P = K\rho^{\frac{n+1}{n}}$$

## DERIVATION 2: WORKING TOWARDS A DIMENSIONLESS FORM

Define a polytropic state equation:

$$P = K \rho^{\frac{n+1}{n}}$$

Make it dimensionless:

$$\theta^n \equiv \frac{\rho}{\rho_c}$$

## DERIVATION 2: WORKING TOWARDS A DIMENSIONLESS FORM

Define a polytropic state equation:

$$P = K \rho^{\frac{n+1}{n}}$$

Make it dimensionless:

$$\theta^n \equiv \frac{\rho}{\rho_c}$$

$$P(r) = K \rho_c^{\frac{n+1}{n}} \theta^{n+1}(r) = P_c \theta^{n+1}(r)$$

## DERIVATION 2: WORKING TOWARDS A DIMENSIONLESS FORM

Define a polytropic state equation:

$$P = K \rho^{\frac{n+1}{n}}$$

Make it dimensionless:

$$\theta^n \equiv \frac{\rho}{\rho_c}$$

$$P(r) = K \rho_c^{\frac{n+1}{n}} \theta^{n+1}(r) = P_c \theta^{n+1}(r)$$

Substitute into Poisson and simplify:

## DERIVATION 2: WORKING TOWARDS A DIMENSIONLESS FORM

Define a polytropic state equation:

$$P = K \rho^{\frac{n+1}{n}}$$

Make it dimensionless:

$$\theta^n \equiv \frac{\rho}{\rho_c}$$

$$P(r) = K \rho_c^{\frac{n+1}{n}} \theta^{n+1}(r) = P_c \theta^{n+1}(r)$$

Substitute into Poisson and simplify:

$$\frac{(n+1)P_c}{4\pi G \rho_c^2} \frac{1}{r^2} \frac{d}{dr} \left( r^2 \frac{d\theta(r)}{dr} \right) = -\theta^n(r)$$



## DERIVATION 3: MORE SIMPLIFICATION

Define a new variable:

## DERIVATION 3: MORE SIMPLIFICATION

Define a new variable:

$$\alpha^2 \equiv \frac{(n+1)P_c}{4\pi G\rho_c^2}$$

## DERIVATION 3: MORE SIMPLIFICATION

Define a new variable:

$$\alpha^2 \equiv \frac{(n+1)P_c}{4\pi G\rho_c^2}$$

Use it to define a dimensionless radius:

## DERIVATION 3: MORE SIMPLIFICATION

Define a new variable:

$$\alpha^2 \equiv \frac{(n+1)P_c}{4\pi G\rho_c^2}$$

Use it to define a dimensionless radius:

$$\xi \equiv \frac{r}{\alpha}$$

## DERIVATION 3: MORE SIMPLIFICATION

Define a new variable:

$$\alpha^2 \equiv \frac{(n+1)P_c}{4\pi G\rho_c^2}$$

Use it to define a dimensionless radius:

$$\xi \equiv \frac{r}{\alpha}$$

Substitute into the simplified Poisson:

## DERIVATION 3: MORE SIMPLIFICATION

Define a new variable:

$$\alpha^2 \equiv \frac{(n+1)P_c}{4\pi G\rho_c^2}$$

Use it to define a dimensionless radius:

$$\xi \equiv \frac{r}{\alpha}$$

Substitute into the simplified Poisson:

$$\frac{1}{\xi^2} \frac{d}{d\xi} \left( \xi^2 \frac{d\theta(\xi)}{d\xi} \right) = -\theta^n(\xi)$$

## TABLE OF CONTENTS

## THEORY

Polytropes

White Dwarfs

## METHODS

## RESULTS

## DISCUSSION

## REFERENCES

# WHY WHITE DWARFS?

- ▶ They are very dense



# WHY WHITE DWARFS?

- ▶ They are very dense
- ▶ So dense that they are completely degenerate

# WHY WHITE DWARFS?

- ▶ They are very dense
- ▶ So dense that they are completely degenerate
- ▶ We'll see why that's important shortly

# DEFINITIONS

## DEFINITION

**Degeneracy** - In quantum mechanics, when 2 or more energy states correspond to the same measured energy

## DEFINITION

**Degenerate Matter** - Quantum version of an ideal gas. Appear under extremely high density or extremely low temperatures.

# DEGENERACY IN WHITE DWARFS

High density results in complete degeneracy of electrons.  
energy state.

# DEGENERACY IN WHITE DWARFS

High density results in complete degeneracy of electrons.  
Pauli exclusion principle prevents more than 2 electrons in each energy state.

# DEGENERACY IN WHITE DWARFS

High density results in complete degeneracy of electrons.  
energy state.

Density of electrons in with a range of momentum  $[p, p + dp]$ :

$$n_e(p, p + dp) \leq \frac{8\pi p^2 dp}{h^3}$$

# DEGENERACY IN WHITE DWARFS

High density results in complete degeneracy of electrons.  
energy state.

Density of electrons in with a range of momentum  $[p, p + dp]$ :

$$n_e(p, p + dp) \leq \frac{8\pi p^2 dp}{h^3}$$

When  $n_e \ll \frac{8\pi p^2 dp}{h^3}$ , behaves as an ideal gas

# DEGENERACY IN WHITE DWARFS

High density results in complete degeneracy of electrons.  
energy state.

Density of electrons in with a range of momentum  $[p, p + dp]$ :

$$n_e(p, p + dp) \leq \frac{8\pi p^2 dp}{h^3}$$

As  $n_e \rightarrow \frac{8\pi p^2 dp}{h^3}$ , degeneracy increases and equation of state becomes:

$$P = K\rho^\gamma$$



# DEGENERACY IN WHITE DWARFS

High density results in complete degeneracy of electrons.  
energy state.

As  $n_e \rightarrow \frac{8\pi p^2 dp}{h^3}$ , degeneracy increases and equation of state becomes:

$$P = K\rho^\gamma$$

As the density increases further, the electrons become relativistic.

# DEGENERACY IN WHITE DWARFS

High density results in complete degeneracy of electrons.  
energy state.

As  $n_e \rightarrow \frac{8\pi p^2 dp}{h^3}$ , degeneracy increases and equation of state becomes:

$$P = K\rho^\gamma$$

In the non-relativistic case,  $\gamma = 5/3$ . In the relativistic case,  
 $\gamma = 4/3$ .

# TABLE OF CONTENTS

## THEORY

Polytropes

White Dwarfs

## METHODS

## RESULTS

## DISCUSSION

## REFERENCES

# REARRANGE THE LANE-EMDEN EQUATION

$$\frac{d^2\theta}{d\xi^2} = -\frac{2}{\xi} \frac{d\theta}{d\xi} - \theta^n(\xi)$$

# TRANSLATING TO A SYSTEM OF 1ST ORDER EQUATIONS

$$\left\{ \begin{array}{lcl} \phi & = & \frac{d\theta}{d\xi} \\ \frac{d\phi}{d\xi} & = & -\frac{2}{\xi}\phi - \theta^n \end{array} \right.$$

# BOUNDARY VALUES

Obtained from central density and hydrostatic equation

$$\xi = 0$$

$$\theta = 1$$

$$\frac{d\theta}{d\xi} = 0$$

# RUNGE-KUTTA SOLUTION

Used a 4th degree Runge-Kutta solver

Did not know the integration range beforehand:

To find the surface with arbitrary precision, backed up a step and halved the step size if  $\theta < 0$  until desired precision reached

# PROBLEM!

Singularity at  $\xi_0 = 0$ :

$$\phi' = - \left( \frac{2}{\xi_0} \phi \right) - \theta_0^n$$

Need to work around this somehow:



# PROBLEM!

Singularity at  $\xi_0 = 0$ :

$$\phi' = - \left( \frac{2}{\xi_0} \phi \right) - \theta_0^n$$

Need to work around this somehow:

- Taylor expand at  $\xi = 0$  and take limit as  $\xi \rightarrow 0$ :  $\phi' \rightarrow -\frac{1}{3}$

# PROBLEM!

Singularity at  $\xi_0 = 0$ :

$$\phi' = - \left( \frac{2}{\xi_0} \phi \right) - \theta_0^n$$

Need to work around this somehow:

- ▶ Taylor expand at  $\xi = 0$  and take limit as  $\xi \rightarrow 0$ :  $\phi' \rightarrow -\frac{1}{3}$
- ▶ Offset the starting point:  $0 < \xi_0 \ll 1$

# GETTING SOMETHING USEFUL

Finding the density & pressure:

$$\frac{\rho}{\rho_c} = \frac{1}{3} \frac{\xi_f}{\theta(\xi_f)}$$

# TABLE OF CONTENTS

## THEORY

Polytropes

White Dwarfs

## METHODS

## RESULTS

## DISCUSSION

## REFERENCES

# SOLUTIONS FOR N=1.5 AND N=3

Parameters for n=1.5, 3 polytropes[1]

$n$	$\xi_f$	$\theta'(\xi_f)$	$\rho_c/\langle\rho\rangle$
1.5	3.6538	-0.20330	5.991
3	6.8969	-0.04243	54.1825


SOLUTIONS FOR  $N=1.5$  AND  $N=3$ 

Image created by presenters in Matlab

# DENSITY PROFILE




Image created by presenters in Matlab

# MASS - RADIUS RELATIONSHIP

We did not get this working correctly yet. We believe we're having trouble with unit conversions



# MASS - RADIUS RELATIONSHIP




Image created by presenters in Matlab

# MASS - RADIUS RELATIONSHIP




Image created by presenters in Matlab

# MASS - RADIUS RELATIONSHIP




Image create by presenters in Matlab, Data from astronomical surveys of white dwarfs in binary systems

# TABLE OF CONTENTS

## THEORY

Polytropes

White Dwarfs

## METHODS

## RESULTS

## DISCUSSION

## REFERENCES

# CHALLENGES

- ▶ Major difficulty was in correctly framing the problem.
- ▶ Initially tried to use a shooting method - but with free boundary it became prohibitively difficult.
- ▶ Rearranged & used known physics to turn into initial value problem.

# CHALLENGES

- ▶ Computing the mass-radius relationship
- ▶ Sources differed on derivation
- ▶ UNITS

# WHERE TO GO FROM HERE?

Fix our calculation of the mass-radius relationship  
Real white dwarfs have a mixed equation of state;  
non-relativistic near surface and highly relativistic near core.  
Approximate this state equation to find behavior near  
Chandrasekhar mass

# QUESTIONS?



Copyright by Abe, <http://downloadwallpaperhd.com> Non-commercial use only



# TABLE OF CONTENTS

## THEORY

Polytropes

White Dwarfs

## METHODS

## RESULTS

## DISCUSSION

## REFERENCES

# REFERENCES I



C. Hansen, S. D. Kawaler, V. Trimble.

*Stellar interiors: physical principles, structure, and evolution.*

Springer-Verlag, 2004.



F. LeBlanc

*An introduction to stellar astrophysics.*

John Wiley & Sons, 2010.



W. S. Jardetsky.

*Theories of figures of celestial bodies.*

Dover Publications, 1958.

## REFERENCES II



E. W. Weisstein.

*Lane-Emden differential equation.*

[http://mathworld.wolfram.com/  
Lane-EmdenDifferentialEquation.html](http://mathworld.wolfram.com/Lane-EmdenDifferentialEquation.html)



V. Dhillon.

Solving the Lane-Emden equation

*PHY 213 - The structure of main-sequence stars*

[http://www.vikdhillon.staff.shef.ac.uk/teaching/  
phy213/phy213\\_le.html](http://www.vikdhillon.staff.shef.ac.uk/teaching/phy213/phy213_le.html)