

Addition with Carries

Eric Errthum

Department of Mathematics and Statistics
Winona State University
Winona, MN

Grade 2 Math: Addition with Carrying

Example

$$\begin{array}{r} 957 \\ + 218 \\ \hline \end{array}$$

Grade 2 Math: Addition with Carrying

Example

$$\begin{array}{r} 1 \\ 957 \\ + 218 \\ \hline 5 \end{array}$$

Grade 2 Math: Addition with Carrying

Example

$$\begin{array}{r} 1 \\ 957 \\ + 218 \\ \hline 75 \end{array}$$

Grade 2 Math: Addition with Carrying

Example

$$\begin{array}{r} 1 \quad 1 \\ 957 \\ + 218 \\ \hline 175 \end{array}$$

Grade 2 Math: Addition with Carrying

Example

$$\begin{array}{r} 1 \quad 1 \\ 957 \\ + 218 \\ \hline 1175 \end{array}$$

Grade 2 Math: Addition with Carrying

Example

$$\begin{array}{r} 1 \quad 1 \\ 957 \\ + 218 \\ \hline 1175 \end{array}$$

Boring...

Grade 2 Fun: Addition with “Wrong” Carrying

Example

$$\begin{array}{r} 957 \\ + 218 \\ \hline \end{array}$$

Grade 2 Fun: Addition with “Wrong” Carrying

Example

$$\begin{array}{r} 3 \\ 957 \\ + 218 \\ \hline 5 \end{array}$$

Grade 2 Fun: Addition with “Wrong” Carrying

Example

$$\begin{array}{r} 3 \\ 957 \\ + 218 \\ \hline 95 \end{array}$$

Grade 2 Fun: Addition with “Wrong” Carrying

Example

$$\begin{array}{r} \\ 957 \\ + 218 \\ \hline 195 \end{array}$$

Grade 2 Fun: Addition with “Wrong” Carrying

Example

$$\begin{array}{r} \\ 957 \\ + 218 \\ \hline 3195 \end{array}$$

Grade 2 Fun: Addition with “Wrong” Carrying

Example

$$\begin{array}{r} 3 3 \\ 957 \\ +_3 218 \\ \hline 3195 \end{array}$$

Grade 2 Fun: Addition with “Wrong” Carrying

Example

$$\begin{array}{r} 3 3 \\ 957 \\ +_3 218 \\ \hline 3195 \end{array}$$

Cool, but I wonder...

What about subtraction?

Example

$$\begin{array}{r} 957 \\ - 3218 \\ \hline \end{array}$$

What about subtraction?

Example

$$\begin{array}{r} 217 \\ 957 \\ -3218 \\ \hline \end{array}$$

What about subtraction?

Example

$$\begin{array}{r} 217 \\ 9\cancel{5}7 \\ -_3 218 \\ \hline 719 \end{array}$$

What about subtraction?

Example

$$\begin{array}{r}
 217 \\
 957 \\
 -3218 \\
 \hline
 719
 \end{array}$$

Example

$$\begin{array}{r}
 3 \\
 719 \\
 +3218 \\
 \hline
 957
 \end{array}$$

Properties

- Commutative.

Properties

- Commutative.
- Inverses (through subtraction) and identity of 0

Properties

- Commutative.
- Inverses (through subtraction) and identity of 0
- Associative?

Properties

- Commutative.
- Inverses (through subtraction) and identity of 0
- Associative? ...Yes.

Properties

- Commutative.
- Inverses (through subtraction) and identity of 0
- Associative? ...Yes.

It's an Abelian Group!

Properties

- Commutative.
- Inverses (through subtraction) and identity of 0
- Associative? ...Yes.

It's an Abelian Group!

Which one is it?

What size are my blocks?

- The digit positions represent generators of the group
- Ten of one generator is equivalent to three of the next generator.
- First (right-most) generator is still just $b_0 = 1$.

What size are my blocks?

- The digit positions represent generators of the group
- Ten of one generator is equivalent to three of the next generator.
- First (right-most) generator is still just $b_0 = 1$.
- The second generator must satisfy $3b_1 = 10 \cdot 1$. Hence $b_1 = \frac{10}{3}$.

What size are my blocks?

- The digit positions represent generators of the group
- Ten of one generator is equivalent to three of the next generator.
- First (right-most) generator is still just $b_0 = 1$.
- The second generator must satisfy $3b_1 = 10 \cdot 1$. Hence $b_1 = \frac{10}{3}$.
- Likewise $3b_2 = 10b_1$. Hence $b_2 = \left(\frac{10}{3}\right)^2$.

What size are my blocks?

- The digit positions represent generators of the group
- Ten of one generator is equivalent to three of the next generator.
- First (right-most) generator is still just $b_0 = 1$.
- The second generator must satisfy $3b_1 = 10 \cdot 1$. Hence $b_1 = \frac{10}{3}$.
- Likewise $3b_2 = 10b_1$. Hence $b_2 = \left(\frac{10}{3}\right)^2$.
- In general

$$b_n = \left(\frac{10}{3}\right)^n.$$

$$957 +_3 218$$

$$957 +_3 218 = \left(9 \left(\frac{10}{3} \right)^2 + 5 \left(\frac{10}{3} \right) + 7 \right) + \left(2 \left(\frac{10}{3} \right)^2 + 1 \left(\frac{10}{3} \right) + 8 \right)$$

$$\begin{aligned}
 957 +_3 218 &= \left(9 \left(\frac{10}{3} \right)^2 + 5 \left(\frac{10}{3} \right) + 7 \right) + \left(2 \left(\frac{10}{3} \right)^2 + 1 \left(\frac{10}{3} \right) + 8 \right) \\
 &= 11 \left(\frac{10}{3} \right)^2 + 6 \left(\frac{10}{3} \right) + 15 \\
 &= 11 \left(\frac{10}{3} \right)^2 + 6 \left(\frac{10}{3} \right) + 3 \left(\frac{10}{3} \right) + 5 \\
 &= 11 \left(\frac{10}{3} \right)^2 + 9 \left(\frac{10}{3} \right) + 5 \\
 &= \left(3 \left(\frac{10}{3} \right) + 1 \right) \left(\frac{10}{3} \right)^2 + 9 \left(\frac{10}{3} \right) + 5
 \end{aligned}$$

When you carry by c with $0 < c < 10$ with $\gcd(c, 10) = 1$, then you have $\mathbb{Z} \left[\frac{10}{c} \right]$.

Theorem

When you carry by c with $0 < c < 10$ with $\gcd(c, 10) = 1$, then you have $\mathbb{Z} \left[\frac{10}{c} \right]$.

Theorem

When you carry by c in base- b with $0 < c < b$ with $\gcd(c, b) = 1$, then you have $\mathbb{Z} \left[\frac{b}{c} \right]$.

Let's Get NUTS!

Let's Get NUTS!

Instead of carrying into one (the next) place, let's carry into a few.

Let's Get NUTS!

Instead of carrying into one (the next) place, let's carry into a few.

Example

Instead of carrying 1's, let's carry the sequence: 1, 7.

$$\begin{array}{r} 957 \\ + (1, 7) \quad 218 \\ \hline \end{array}$$

Let's Get NUTS!

Instead of carrying into one (the next) place, let's carry into a few.

Example

Instead of carrying 1's, let's carry the sequence: 1, 7.

$$\begin{array}{r} 17 \\ 957 \\ + (1, 7) \\ \hline 218 \\ 5 \end{array}$$

Let's Get NUTS!

Instead of carrying into one (the next) place, let's carry into a few.

Example

Instead of carrying 1's, let's carry the sequence: 1, 7.

$$\begin{array}{r} 17 \\ 17 \\ 957 \\ + (1, 7) \quad 218 \\ \hline 35 \end{array}$$

Let's Get NUTS!

Instead of carrying into one (the next) place, let's carry into a few.

Example

Instead of carrying 1's, let's carry the sequence: 1, 7.

$$\begin{array}{r} 17 \\ 17 \\ 17 \\ 957 \\ + (1, 7) \quad 218 \\ \hline 935 \end{array}$$

Let's Get NUTS!

Instead of carrying into one (the next) place, let's carry into a few.

Example

Instead of carrying 1's, let's carry the sequence: 1, 7.

$$\begin{array}{r} 17 \\ 17 \\ 17 \\ 957 \\ + (1, 7) \quad 218 \\ \hline 18935 \end{array}$$

What size are my blocks?

- In this case, $b_n + 7b_{n-1} = 10b_{n-2}$, $b_0 = 1$.
- This is a homogeneous linear recursion with characteristic equation $r^2 + 7r - 10 = 0$.

What size are my blocks?

- In this case, $b_n + 7b_{n-1} = 10b_{n-2}$, $b_0 = 1$.
- This is a homogeneous linear recursion with characteristic equation $r^2 + 7r - 10 = 0$.
- Hence

$$b_n = A \left(\frac{-7 + \sqrt{89}}{2} \right)^n + (1 - A) \left(\frac{-7 - \sqrt{89}}{2} \right)^n.$$

What size are my blocks?

- In this case, $b_n + 7b_{n-1} = 10b_{n-2}$, $b_0 = 1$.
- This is a homogeneous linear recursion with characteristic equation $r^2 + 7r - 10 = 0$.
- Hence

$$b_n = A \left(\frac{-7 + \sqrt{89}}{2} \right)^n + (1 - A) \left(\frac{-7 - \sqrt{89}}{2} \right)^n.$$

- If we force b_1 to be the positive root, then

$$b_n = \left(\frac{-7 + \sqrt{89}}{2} \right)^n$$

and again we have a base-like representation.

Theorem

When you carry by sequence $(c_n, c_{n-1}, \dots, c_1)$, then you have $\mathbb{Z}[\alpha]$ where α satisfies

$$c_n \alpha^n + c_{n-1} \alpha^{n-1} + \dots + c_1 \alpha - 10 = 0.$$

Theorem

When you carry by sequence $(c_n, c_{n-1}, \dots, c_1)$ and use base b digits, then you have $\mathbb{Z}[\alpha]$ where α satisfies

$$c_n \alpha^n + c_{n-1} \alpha^{n-1} + \dots + c_1 \alpha - b = 0.$$

Example

In base $\alpha = \frac{1}{10} (\sqrt{249} - 7)$ you carry the sequence: 5, 7.

$+(5, 7)$

Bad α 's

Example

In base $\alpha = \frac{1}{10} (\sqrt{249} - 7)$ you carry the sequence: 5, 7.

$$\begin{array}{r} 57 \\ + (5, 7) \\ \hline \end{array}$$

In base $\alpha = \frac{1}{10} (\sqrt{249} - 7)$ you carry the sequence: 5, 7.

$$\begin{array}{r} 57 \\ 57 \\ + (5, 7) \\ \hline 2 \end{array}$$

Bad α 's

Example

In base $\alpha = \frac{1}{10} (\sqrt{249} - 7)$ you carry the sequence: 5, 7.

$$\begin{array}{r} 57 \\ 57 \\ 57 \\ + (5, 7) \\ \hline 22 \end{array}$$

Bad α 's

Example

In base $\alpha = \frac{1}{10} (\sqrt{249} - 7)$ you carry the sequence: 5, 7.

$$\begin{array}{r} 57 \\ 57 \\ 57 \\ 57 \\ + (5, 7) \\ \hline 222 \end{array}$$

Bad α 's

Example

In base $\alpha = \frac{1}{10} (\sqrt{249} - 7)$ you carry the sequence: 5, 7.

$$\begin{array}{r} 57 \\ 57 \\ 57 \\ 57 \\ + (5, 7) \quad 5 \\ \hline \dots 22222 \end{array}$$

Bad α 's

Example

In base $\alpha = \frac{1}{10} (\sqrt{249} - 7)$ you carry the sequence: 5, 7.

$$\begin{array}{r} 5\ 7 \\ 5\ 7 \\ 5\ 7 \\ 5\ 7 \\ + (5, 7) \quad 5 \\ \hline \dots\ 2\ 2\ 2\ 2\ 2 \end{array}$$

But $\alpha \approx 0.877973 \dots < 1$

Bad α 's

Example

In base $\alpha = \frac{1}{10} (\sqrt{249} - 7)$ you carry the sequence: 5, 7.

$$\begin{array}{r} 5\ 7 \\ 5\ 7 \\ 5\ 7 \\ 5\ 7 \\ + (5, 7) \quad 5 \\ \hline \dots\ 2\ 2\ 2\ 2\ 2 \end{array}$$

But $\alpha \approx 0.877973 \dots < 1$

Theorem

For $c_i \geq 0$, if $c_n + c_{n-1} + \dots + c_1 > b$, then $0 < \alpha < 1$.

New Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3}$.

New Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3}$.

The α satisfies $3\alpha^2 + 4\alpha - 11 = 0$. So use **base-11 digits**, e.g. 0, 1, ..., 8, 9, a and carry the sequence (3, 4)

New Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3}$.

The α satisfies $3\alpha^2 + 4\alpha - 11 = 0$. So use **base-11 digits**, e.g. 0, 1, ..., 8, 9, a and carry the sequence (3, 4)



New Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3}$.

The α satisfies $3\alpha^2 + 4\alpha - 11 = 0$. So use **base-11 digits**, e.g. 0, 1, ..., 8, 9, a and carry the sequence (3, 4)

		9	12	
				38
				5

New Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3}$.

The α satisfies $3\alpha^2 + 4\alpha - 11 = 0$. So use **base-11 digits**, e.g. 0, 1, ..., 8, 9, a and carry the sequence (3, 4)

	3	4		
		9	12	
				38
			1	5

New Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3}$.

The α satisfies $3\alpha^2 + 4\alpha - 11 = 0$. So use **base-11 digits**, e.g. 0, 1, ..., 8, 9, a and carry the sequence (3, 4)

3	4			
	3	4		
		9	12	
				38
		2	1	5

New Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3}$.

The α satisfies $3\alpha^2 + 4\alpha - 11 = 0$. So use **base-11 digits**, e.g. 0, 1, ..., 8, 9, a and carry the sequence (3, 4)

3	4			
	3	4		
		9	12	
				38
3	7	2	1	5

New Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3}$.

The α satisfies $3\alpha^2 + 4\alpha - 11 = 0$. So use **base-11 digits**, e.g. 0, 1, ..., 8, 9, a and carry the sequence (3, 4)

3		4						
		3		4				
				9		12		
								38
3		7		2		1		5

Homework: Check $3\alpha^4 + 7\alpha^3 + 2\alpha^2 + \alpha + 5 = 38$.

Standard Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3} \approx 1.36092\dots$

Standard Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3} \approx 1.36092\dots$

$$\alpha^{11} < 38 < \alpha^{12}$$

Standard Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3} \approx 1.36092\dots$

$$\alpha^{11} < 38 < \alpha^{12}$$

$$\alpha^6 < 38 - \alpha^{11} < \alpha^7$$

Standard Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3} \approx 1.36092\dots$

$$\alpha^{11} < 38 < \alpha^{12}$$

$$\alpha^6 < 38 - \alpha^{11} < \alpha^7$$

$$\alpha^2 < 38 - \alpha^{11} - \alpha^6 < \alpha^3$$

Standard Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3} \approx 1.36092\dots$

$$\alpha^{11} < 38 < \alpha^{12}$$

$$\alpha^6 < 38 - \alpha^{11} < \alpha^7$$

$$\alpha^2 < 38 - \alpha^{11} - \alpha^6 < \alpha^3$$

$$\alpha^{-7} < 38 - \alpha^{11} - \alpha^6 - \alpha^2 < \alpha^{-6}$$

$$\vdots$$

Standard Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3} \approx 1.36092\dots$

$$\alpha^{11} < 38 < \alpha^{12}$$

$$\alpha^6 < 38 - \alpha^{11} < \alpha^7$$

$$\alpha^2 < 38 - \alpha^{11} - \alpha^6 < \alpha^3$$

$$\alpha^{-7} < 38 - \alpha^{11} - \alpha^6 - \alpha^2 < \alpha^{-6}$$

$$\vdots$$

$$38 = 100001000100.0000001000001\dots$$

Standard Approach to Nonintegral Bases

Example

Put 38 into base $\alpha = \frac{\sqrt{37} - 2}{3} \approx 1.36092\dots$

$$\alpha^{11} < 38 < \alpha^{12}$$

$$\alpha^6 < 38 - \alpha^{11} < \alpha^7$$

$$\alpha^2 < 38 - \alpha^{11} - \alpha^6 < \alpha^3$$

$$\alpha^{-7} < 38 - \alpha^{11} - \alpha^6 - \alpha^2 < \alpha^{-6}$$

$$\vdots$$

$$38 = 100001000100.0000001000001\dots$$

And the addition rules are even worse.

So the new way is better

- Pros

So the new way is better

- Pros
 - Every integer has a finite expression

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons

Example

$$\begin{array}{r} 20 \\ -3 1 \\ \hline \end{array}$$

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons

Example

$$\begin{array}{r}
 -1 \ 10 \\
 2 \ 0 \\
 \hline
 -3 \qquad 1
 \end{array}$$

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons

Example

$$\begin{array}{r}
 -1 \ 10 \\
 \cancel{2} \ \emptyset \\
 1 \\
 \hline
 -3 \\
 9
 \end{array}$$

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons

Example

$$\begin{array}{r}
 9 \\
 -3 \quad \cancel{-1} \quad 10 \\
 \quad \quad \cancel{2} \quad \emptyset \\
 \quad \quad \quad 1 \\
 \hline
 -3 \quad \quad \quad 9
 \end{array}$$

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons

Example

$$\begin{array}{r}
 9 \\
 -3 \quad \cancel{-1} \quad 10 \\
 \quad \quad \cancel{2} \quad \emptyset \\
 \quad \quad \quad 1 \\
 \hline
 -3 \quad \quad \quad 9 \quad 9
 \end{array}$$

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons

Example

$$\begin{array}{r}
 7 9 \\
 -3 \cancel{-3} \cancel{-1} 10 \\
 \cancel{2} \cancel{0} \\
 1 \\
 \hline
 -3 \\
 7 9 9
 \end{array}$$

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons

Example

$$\begin{array}{r}
 \dots \quad 7 \quad 7 \quad 9 \\
 \dots \quad \cancel{-3} \quad \cancel{-3} \quad \cancel{-1} \quad 10 \\
 \phantom{\dots \quad \cancel{-3} \quad \cancel{-3} \quad} 2 \quad 0 \\
 \phantom{\dots \quad \cancel{-3} \quad \cancel{-3} \quad} 1 \\
 \hline
 \dots 7 \quad 7 \quad 7 \quad 9 \quad 9
 \end{array}$$

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons

Example

$$\begin{array}{r}
 \dots \quad 7 \quad 7 \quad 9 \\
 \dots \quad \cancel{-3} \quad \cancel{-3} \quad \cancel{-1} \quad 10 \\
 \phantom{\dots \quad \cancel{-3} \quad \cancel{-3} \quad} 2 \quad 0 \\
 \phantom{\dots \quad \cancel{-3} \quad \cancel{-3} \quad} \underline{-3 1} \\
 \dots 7 \quad 7 \quad 7 \quad 9 \quad 9
 \end{array}$$

Note: $2 \cdot \frac{10}{3} - 1 = \frac{17}{3} = 9 + 9 \cdot \frac{10}{3} + 7 \left(\frac{10}{3} \right)^2 \left(\frac{1}{1 - \frac{10}{3}} \right)$

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons
 - Subtraction might require infinite digits to the left.

So the new way is better

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons
 - Subtraction might require infinite digits to the left.
 - Not easily extended to represent all of \mathbb{Q}

So the new way is better ... right?

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons
 - Subtraction might require infinite digits to the left.
 - Not easily extended to represent all of \mathbb{Q}

So the new way is better ... right?

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons
 - Subtraction might require infinite digits to the left.
 - Not easily extended to represent all of \mathbb{Q}

Take something elementary and explore mathematical silliness.

So the new way is better ... right?

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons
 - Subtraction might require infinite digits to the left.
 - Not easily extended to represent all of \mathbb{Q}

Take something elementary and explore mathematical silliness.

Applications: Fun with kids.

So the new way is better ... right?

- Pros
 - Every integer has a finite expression
 - Easy to add
 - IT'S FUN!
- Cons
 - Subtraction might require infinite digits to the left.
 - Not easily extended to represent all of \mathbb{Q}

Take something elementary and explore mathematical silliness.

Applications: Fun with kids.

Credit to Daniel C. Isaksen "A Cohomological Viewpoint on Elementary School Arithmetic"

Thanks for staying late on Saturday!