Equilibrium

Question1

The molar solubility(s) of zirconium phosphate with molecular formula $\left(Zr^{4+}\right)_3\left(PO_4^{3-}\right)_4$ is given by relation :

JEE Main 2025 (Online) 22nd January Evening Shift

Options:

A. $\left(\frac{K_{sp}}{5348}\right)^{\frac{1}{6}}$

B. $\left(\frac{K_{sp}}{8435}\right)^{\frac{1}{7}}$

C. $\left(\frac{K_{sp}}{6912}\right)^{\frac{1}{7}}$

D. $\left(\frac{K_{sp}}{9612}\right)^{\frac{1}{3}}$

Answer: C

Solution:

$$Zr_3(PO_4)_4(s) \rightleftharpoons \underset{3s}{3Zr^{4+}}(aq) + \underset{4s}{4PO_4^{3-}}(aq)$$

$$K_{sp} = (3\,s)^3 \cdot (4\,s)^4$$

$$K_{sp} = 6912\,s^7$$

$$s = \left(\frac{K_{sp}}{6912}\right)^{\frac{1}{7}}$$

Question2

Which of the following happens when NH_4OH is added gradually to the solution containing 1 M A^{2+} and $1MB^{3+}$ ions?

Given : $K_{sp}\ \ [A(OH)_2] = 9 \times 10^{-10}$ **and** $K_{sp}\,[B(OH)_3] = 27 \times 10^{-18}$ **at 298 K.**

JEE Main 2025 (Online) 23rd January Morning Shift

Options:

A. $A(OH)_2$ will precipitate before $B(OH)_3$

B. $A(OH)_2$ and $B(OH)_3$ will precipitate together

C. Both $A(OH)_2$ and $B(OH)_3$ do not show precipitation with NH_4OH

D. $B(OH)_3$ will precipitate before $A(OH)_2$

Answer: D

Solution:

Condition for precipitation $Q_{ip} > K_{sp}$

For $[A(OH)_2]$

$$[A^{2+}][OH^-]^2 > 9 \times 10^{-10}$$
$$\left[A^{+2}\right] = 1M$$
$$\Rightarrow [OH^-] > 3 \times 10^{-5}M$$

For $[B(OH)_3]$

$$[B^{3+}][OH^-]^3 > 27 \times 10^{-18}$$
$$\left[B^{3+}\right] = 1M$$
$$\Rightarrow [OH^-] > 3 \times 10^{-6}M$$

So, $B(OH)_3$ will precipitate before $A(OH)_2$

Question3

pH of water is 7 at $25^\circ C$. If water is heated to $80^\circ C$., it's pH will :

JEE Main 2025 (Online) 23rd January Evening Shift

Options:

A. Decrease

B. Remains the same

C. Increase

D. H^+concentration increases, OH^-concentration decreases

Answer: A

Solution:

When water is heated, its self-ionization increases. At 25°C, the water ionization constant is given by

$K_w = [\mathrm{H^+}][\mathrm{OH^-}] = 1.0 \times 10^{-14}$,

so in pure water,

$[\mathrm{H^+}] = [\mathrm{OH^-}] = 1.0 \times 10^{-7}\,\mathrm{M}$,

which corresponds to a pH of 7.

As water is heated to 80°C, the value of K_w increases due to the endothermic nature of water's autoionization. This means that both $[\mathrm{H^+}]$ and $[\mathrm{OH^-}]$ increase. However, since their concentrations remain equal (which defines neutrality at that temperature), the pH isn't 7 anymore. Instead, the increased concentration of $\mathrm{H^+}$ ions leads to a pH value lower than 7.

Key points:

Heating water increases K_w.

Higher K_w means higher $[\mathrm{H^+}]$ (and equally higher $[\mathrm{OH^-}]$).

Thus, the pH decreases (e.g., it might be around 6.5–6.6 at 80°C) even though the water is still neutral.

So, the correct answer is:

Option A – Decrease.

Question4

$\mathrm{K_{sp}}$ for $\mathrm{Cr(OH)_3}$ is 1.6×10^{-30}. What is the molar solubility of this salt in water?

JEE Main 2025 (Online) 24th January Morning Shift

Options:

A. $\sqrt[5]{1.8 \times 10^{-30}}$

B. $\frac{1.8 \times 10^{-30}}{27}$

C. $\sqrt[4]{\frac{1.6 \times 10^{-30}}{27}}$

D. $\sqrt[2]{1.6 \times 10^{-30}}$

Answer: C

Solution:

To find the molar solubility of $\mathrm{Cr(OH)_3}$ in water, we start with the dissolution equilibrium:

$$\mathrm{Cr(OH)_{3(\,s)} \rightleftharpoons \underset{s}{Cr^{3+}_{(aq)}} + \underset{3s}{3OH^-_{(aq)}}}$$

At equilibrium, the solubility product $\mathrm{K_{sp}}$ is given by:

$$\mathrm{K_{sp} = (s) \cdot (3\,s)^3 = 27\,s^4}$$

Substituting the given $\mathrm{K_{sp}}$ value:

$27\,s^4 = 1.6 \times 10^{-30}$

Solving for s, we divide both sides by 27:

$s = \left(\frac{1.6 \times 10^{-30}}{27}\right)^{1/4}$

Thus, the molar solubility of $Cr(OH)_3$ is:

$\left(\frac{1.6}{27} \times 10^{-30}\right)^{1/4}$

Question5

A weak acid HA has degree of dissociation x . Which option gives the correct expression of (pH - pK_a)?

JEE Main 2025 (Online) 28th January Morning Shift

Options:

A. $\log\left(\frac{1-x}{x}\right)$

B. 0

C. $\log(1+2x)$

D. $\log\left(\frac{x}{1-x}\right)$

Answer: D

Solution:

$HA \rightleftharpoons H^{\oplus} + A^{\Theta}$
$t = 0 \quad a$
$t = t \quad a(1-x) \quad ax \quad ax$
$K_a = (ax)\frac{(x)}{1-x}; \left[H^+\right] = ax$
$-\log(K_a) = -\log(ax) - \log\left(\frac{x}{1-x}\right)$
$pKa = pH - \log\left(\frac{x}{1-x}\right)$
$pH - pKa = \log\left(\frac{x}{1-x}\right)$

Question6

Arrange the following in increasing order of solubility product :

$Ca(OH)_2, AgBr, PbS, HgS$

JEE Main 2025 (Online) 28th January Evening Shift

Options:

A. $\mathrm{PbS} < \mathrm{HgS} < \mathrm{Ca(OH)_2} < \mathrm{AgBr}$

B. $\mathrm{HgS} < \mathrm{AgBr} < \mathrm{PbS} < \mathrm{Ca(OH)_2}$

C. $\mathrm{HgS} < \mathrm{PbS} < \mathrm{AgBr} < \mathrm{Ca(OH)_2}$

D. $\mathrm{Ca(OH)_2} < \mathrm{AgBr} < \mathrm{HgS} < \mathrm{PbS}$

Answer: C

Solution:

Based on the Ksp values and salt analysis cation identification, we can say that order of Ksp value is:

$\mathrm{HgS} < \mathrm{PbS} < \mathrm{AgBr} < \mathrm{Ca(OH)_2}$

Ksp values

$\mathrm{HgS} \rightarrow 4 \times 10^{-53}$
$\mathrm{PbS} \rightarrow 8 \times 10^{-28}$
$\mathrm{AgBr} \rightarrow 5 \times 10^{-13}$
$\mathrm{Ca(OH)_2} \rightarrow 5.5 \times 10^{-6}$

Question7

If equal volumes of AB_2 and XY (both are salts) aqueous solutions are mixed, which of the following combination will give a precipitate of $\mathrm{AY_2}$ at 300 K ?(Given $\mathrm{K_{sp}}$ (at 300 K) for $\mathrm{AY_2} = 5.2 \times 10^{-7}$)

JEE Main 2025 (Online) 2nd April Morning Shift

Options:

A. $2.0 \times 10^{-4} \mathrm{MAB_2}, 0.8 \times 10^{-3} \mathrm{MXY}$

B. $2.0 \times 10^{-2} \mathrm{MAB_2}, 2.0 \times 10^{-2} \mathrm{MXY}$

C. $1.5 \times 10^{-4} \mathrm{MAB_2}, 1.5 \times 10^{-3} \mathrm{MXY}$

D. $3.6 \times 10^{-3} \mathrm{MAB_2}, 5.0 \times 10^{-4} \mathrm{MXY}$

Answer: B

Solution:

When equal volumes of solutions are mixed, the molarity of each solution halves. To determine if a precipitate will form, calculate the ionic product $\mathrm{Q_{SP}}$ using the formula:

$Q_{SP} = [A^{2+}][Y^{-}]^2$

A precipitate of AY_2 will form if $Q_{SP} > K_{SP}$, where K_{SP} is given as 5.2×10^{-7}.

Option 1 Calculation:

$Q_{SP} = \left(1.8 \times 10^{-3}\right)\left(\frac{5}{2} \times 10^{-4}\right)^2 =$ Calculated value $< K_{SP}$

Option 2 Calculation:

$Q_{SP} = \left(10^{-4}\right)\left(0.4 \times 10^{-3}\right)^2 =$ Calculated value $< K_{SP}$

Option 3 Calculation:

$Q_{SP} = \left(10^{-2}\right)\left(10^{-2}\right)^2 =$ Calculated value $> K_{SP}$


Option 4 Calculation:

$Q_{SP} = \left(\frac{1.5}{2} \times 10^{-4}\right)\left(\frac{1.5}{2} \times 10^{-3}\right)^2 =$ Calculated value $< K_{SP}$

From the calculations, the solution in Option 3 will form a precipitate since the ionic product Q_{SP} is greater than the solubility product K_{SP}.

Question8

40 mL of a mixture of CH_3COOH and HCl (aqueous solution) is titrated against 0.1 M NaOH solution conductometrically. Which of the following statement is correct?



JEE Main 2025 (Online) 3rd April Evening Shift

Options:

A. The concentration of CH_3COOH in the original mixture is 0.005 M

B. The concentration of HCl in the original mixture is 0.005 M

C. CH_3COOH is neutralised first followed by neutralisation of HCl

D. Point ' C ' indicates the complete neutralisation of HCl

Answer: B

Solution:

For HCl:

Moles of HCl = Moles of NaOH used

$M \times 40 = 0.1 \times 2$

$M = 0.005$

For CH_3COOH:

Moles of CH_3COOH = Moles of NaOH used

$M \times 40 = 0.1 \times 3$

$M = 0.0075$

Since HCl is a strong acid, it is neutralized first.

Question9

An aqueous solution of HCl with pH 1.0 is diluted by adding equal volume of water (ignoring dissociation of water). The pH of HCl solution would

(**Given** $\log 2 = 0.30$)

JEE Main 2025 (Online) 7th April Morning Shift

Options:

A. increase to 1.3

B. reduce to 0.5

C. increase to 2

D. remain same

Answer: A

Solution:

An aqueous solution of HCl initially has a pH of 1.0, meaning the hydrogen ion concentration is $[H^+] = 10^{-1}$ M.

When we dilute the solution by adding an equal volume of water, the concentration of hydrogen ions will be halved:

$[H^+]_{new} = \frac{10^{-1}}{2} = 5 \times 10^{-2}$ M

To find the new pH, we use the pH formula:

$pH = -\log[H^+]$

Substituting the new hydrogen ion concentration gives:

$pH = -\log(5 \times 10^{-2})$

Using the logarithm property $\log(a \times b) = \log a + \log b$:

$pH = -(\log 5 + \log 10^{-2}) = -\log 5 + 2$

Given $\log 2 = 0.30$, we need $\log 5$, which can be calculated as:

$\log 10 = \log(2 \times 5) = \log 2 + \log 5 \Rightarrow \log 5 = \log 10 - \log 2 = 1 - 0.30 = 0.70$

Thus:

$\mathrm{pH} = -(0.70) + 2 = 1.3$

Therefore, the pH of the solution increases to 1.3.

Question10

If 1 mM solution of ethylamine produces $\mathrm{pH} = 9$, then the ionization constant $(\mathrm{K_b})$ of ethylamine is 10^{-x}. The value of x is _____ (nearest integer).

[The degree of ionization of ethylamine can be neglected with respect to unity.]

JEE Main 2025 (Online) 23rd January Morning Shift

Answer: 7

Solution:

$\mathrm{C_2H_5NH_2(aq) + H_2O \rightleftharpoons C_2H_2NH_3^+ + \overset{\ominus}{O}\,H}$

$$\begin{array}{ccccc} \mathrm{C = 10^{-3}M} & & - & & - \\ \mathrm{C(1-\alpha)} & & \mathrm{C\alpha} & & \mathrm{C\alpha} \\ \Rightarrow \mathrm{C = 10^{-3}} & & = 10^{-5} & & = 10^{-5} \end{array}$$

$1 - \alpha \simeq 1$

Given, $\mathrm{P^H = 9 \Rightarrow P^{OH} = 5 \Rightarrow [\overset{\ominus}{O}\,H] = 10^{-5}M}$

Now, $K_b = \frac{[\mathrm{C_2H_5NH_3^+}][\overset{\ominus}{\mathrm{OH}}]}{[\mathrm{C_2H_5NH_2}]}$

$\Rightarrow \mathrm{K_b} = \frac{10^{-5} \times 10^{-5}}{10^{-3}} = 10^{-7}$

Question11

The observed and normal molar masses of compound $\mathrm{MX_2}$ are 65.6 and 164 respectively. The percent degree of ionisation of $\mathrm{MX_2}$ is _____%. (Nearest integer)

JEE Main 2025 (Online) 24th January Evening Shift

Answer: 75

Solution:

To calculate the percent degree of ionization of the compound MX_2, we begin by considering its dissociation process:

$MX_2 \rightarrow M^{+2} + 2X^-$

Next, we use the van't Hoff factor (i), calculated as the ratio of the normal molar mass to the observed molar mass:

$i = \frac{\text{normal molar mass}}{\text{observed molar mass}} = \frac{164}{65.6}$

Now, expressing the relationship of ionization in terms of the van't Hoff factor, we have:

$1 + (3 - 1)\alpha = \frac{164}{65.6}$

Simplifying this equation gives us:

$2\alpha = \frac{164}{65.6} - 1 = \frac{98.4}{65.6}$

Solving for α:

$\alpha = 0.75$

Therefore, the percent dissociation, or the percent degree of ionization, is:

Percent dissociation $= 75\%$

Question12

The pH of a 0.01 M weak acid HX $\left(K_a = 4 \times 10^{-10}\right)$ is found to be 5 . Now the acid solution is diluted with excess of water so that the pH of the solution changes to 6 . The new concentration of the diluted weak acid is given as $x \times 10^{-4} M$. The value of x is _____ (nearest integer)

JEE Main 2025 (Online) 4th April Morning Shift

Answer: 25

Solution:

$HX_{(aq)} \rightleftharpoons H^+_{(aq)} + X^-_{(aq)} K_a = 4 \times 10^{-10}$

$0.01(1-\alpha) \quad 0.01\alpha \quad 0.01\alpha \quad$ Not justified
$\Rightarrow 0.01\alpha = 10^{-5} \Rightarrow \alpha = 10^{-3}$
$K_a = 0.01\alpha^2 = 10^{-8}$

On dilution let final concentration of $HX = cM$
$Hx_{(aq)} \rightleftharpoons H^+_{(aq)} + X^-_{(aq)}$

$C(1-\alpha) \quad C\alpha \quad C\alpha$

$\Rightarrow C\alpha = 10^{-6}$ ..... (1)

$\frac{C\alpha^2}{1-\alpha} = \mathrm{K_a} = 10^{-8}$ ..... (2)

$\Rightarrow \frac{10^{-6}\alpha}{1-\alpha} = 10^{-8}$

Data given is inconsistent & contradictory. This should be bonus.

Question13

$x$mg of $\mathrm{Mg(OH)_2}$(molar mass $= 58$) is required to be dissolved in 1.0 L of water to produce a pH of 10.0 at 298 K . The value of x is _____ mg. (Nearest integer)

(Given : $\mathrm{Mg(OH)_2}$ is assumed to dissociate completely in $\mathrm{H_2O}$]

JEE Main 2025 (Online) 4th April Evening Shift

Answer: 3

Solution:

To determine the mass of $Mg(OH)_2$ that needs to be dissolved to achieve a pH of 10.0, follow these steps:

Given:

pH = 10

Therefore, pOH = 14 - pH = 4

$[OH^-] = 10^{-4}$ M

First, calculate the number of moles of OH^- ions:

Number of moles of $OH^- = 10^{-4}$

Since $Mg(OH)_2$ dissociates completely in water, from one mole of $Mg(OH)_2$, you get two moles of OH^-:

Number of moles of $Mg(OH)_2 = (10^{-4}) / 2 = 5 \times 10^{-5}$ moles

Next, calculate the mass of $Mg(OH)_2$:

Molar mass of $Mg(OH)_2$ = 58 g/mol

Convert the number of moles to mass:

Mass of $Mg(OH)_2$ = (5×10^{-5} moles) × (58 g/mol) = 2.9×10^{-3} g

Convert to milligrams: 2.9×10^{-3} g = 2.9 mg

Therefore, the calculated mass of $Mg(OH)_2$ required is approximately 2.9 mg.

Question14

The molar conductance of an infinitely dilute solution of ammonium chloride was found to be $185\ \mathrm{S\ cm^2\ mol^{-1}}$ and the ionic conductance of hydroxyl and chloride ions

are 170 and $70\ \mathrm{S\ cm^2\ mol^{-1}}$, respectively. If molar conductance of 0.02 M solution of ammonium hydroxide is $85.5\ \mathrm{S\ cm^2\ mol^{-1}}$, its degree of dissociation is given by $x \times 10^{-1}$.The value of x is _____ . (Nearest integer)

JEE Main 2025 (Online) 4th April Evening Shift

Answer: 3

Solution:

$\lambda_{\mathrm{m}}^{\circ}$ of $\mathrm{NH_4Cl} = 185$

$(\lambda_{\mathrm{m}}^{\circ})_{\mathrm{NH_4^+}} + (\lambda_{\mathrm{m}}^{\circ})_{\mathrm{Cl^-}} = 185$

$(\lambda_{\mathrm{m}}^{\circ})_{\mathrm{NH_4^+}} = 185 - 70 = 115 \mathrm{Scm^2\ mol^{-1}}$

$(\lambda_{\mathrm{m}}^{\circ})_{\mathrm{NH_4OH}} = (\lambda_{\mathrm{m}}^{\circ})_{\mathrm{NH_4^+}} + (\lambda_{\mathrm{m}}^{\circ})_{\mathrm{OH^-}}$

$= 115 + 170$

$(\lambda_{\mathrm{m}}^{\circ})_{\mathrm{NH_4OH}} = 285 \mathrm{Scm^2\ mol^{-1}}$

$$\begin{aligned}\text{degree of dissociation} &= \frac{(\lambda_{\mathrm{m}})_{\mathrm{NH_4OH}}}{(\lambda_{\mathrm{m}}^{\circ})_{\mathrm{NH_4OH}}} \\ &= \frac{85.5}{285} \\ &= 0.3 \\ &= 3 \times 10^{-1}\end{aligned}$$

Question15

The percentage dissociation of a salt $(\mathrm{MX_3})$ solution at given temperature (van't Hoff factor $\mathrm{i} = 2$) is _____ %(Nearest integer)

JEE Main 2025 (Online) 7th April Morning Shift

Answer: 33

Solution:

To determine the percentage dissociation of the salt $\mathrm{MX_3}$, consider the following dissociation reaction:

$\mathrm{MX_3} \rightarrow \mathrm{M^{+3}} + 3\mathrm{X^-}$

The van't Hoff factor, i, is given as 2. The formula relating i to the degree of dissociation, α, is:

$\mathrm{i} = 1 + (\mathrm{n} - 1)\alpha$

Here, n represents the total number of ions produced per formula unit of the salt. For $\mathrm{MX_3}$, dissociation yields:

1 $\mathrm{M^{+3}}$ ion

3 X^- ions

Thus, n = 4. Substituting into the formula, we have:

$2 = 1 + (4 - 1)\alpha$

Simplifying, we solve for α:

$2 = 1 + 3\alpha$

$3\alpha = 1$

$\alpha = \frac{1}{3} \approx 0.3333$

As a percentage, this degree of dissociation is:

$\alpha \times 100\% \approx 33.33\%$

Rounding to the nearest integer gives:

33%

Therefore, the percentage dissociation of the salt MX_3 is approximately 33%.

Question16

One litre buffer solution was prepared by adding 0.10 mol each of NH_3 and NH_4Cl in deionised water. The change in pH on addition of 0.05 mol of HCl to the above solution is _____ $\times 10^{-2}$.

(Nearest integer)

Given : pK_b of $NH_3 = 4.745$ and $\log_{10} 3 = 0.477$

JEE Main 2025 (Online) 7th April Evening Shift

Answer: 48

Solution:

$$pOH = pK_b + \log \frac{[NH_4^+]}{[NH_3]}$$

$\mathbf{pOH} = 4.745$
on adding 0.05 mole HCl

NH_3	+ $H^{\oplus}$	$\rightarrow$	$NH_4^{\oplus}$
0.1	0.05		0.1
0.05	0		0.15

$pOH' = 4.745 + \log 3$
$pOH' - pOH = 0.477$
$14 - pH' - 14 + pH = 0.477$
$\Delta pH = 0.477$
$= 47.7 \times 10^{-2} \approx 48 \times 10^{-2}$

Question17

A vessel at 1000 K contains CO_2 with a pressure of 0.5 atm . Some of CO_2 is converted into CO on addition of graphite. If total pressure at equilibrium is 0.8 atm , then K_p is :

JEE Main 2025 (Online) 22nd January Morning Shift

Options:

A. 0.18 atm

B. 0.3 atm

C. 3 atm

D. 1.8 atm

Answer: D

Solution:

The reaction is:

$\mathrm{CO}_2(g) + \mathrm{C}(s) \rightleftharpoons 2\mathrm{CO}(g)$

Initially, the pressure of CO_2 is 0.5 atm, and the pressure of CO is 0 atm. Let 'x' be the change in pressure of CO_2 at equilibrium. Since the stoichiometric coefficient of CO is twice that of CO_2, the pressure of CO formed at equilibrium is 2x.

At equilibrium:

Pressure of $\mathrm{CO}_2 = 0.5 - x$

Pressure of $\mathrm{CO} = 2x$

The total pressure at equilibrium is given as 0.8 atm. Therefore:

$(0.5 - x) + 2x = 0.8$

$0.5 + x = 0.8$

$x = 0.8 - 0.5 = 0.3$

So, at equilibrium:

Pressure of $\mathrm{CO}_2 = 0.5 - 0.3 = 0.2$ atm

Pressure of $\mathrm{CO} = 2(0.3) = 0.6$ atm

The equilibrium constant K_p is given by:

$K_p = \frac{P_{\mathrm{CO}}^2}{P_{\mathrm{CO}_2}}$

$K_p = \frac{(0.6)^2}{0.2} = \frac{0.36}{0.2} = 1.8$ atm

Therefore, the value of K_p is 1.8 atm.

Question18

Consider the reaction

$$\mathrm{X_2Y(\ g) \rightleftharpoons X_2(\ g) + \frac{1}{2}Y_2(\ g)}$$

The equation representing correct relationship between the degree of dissociation (x) of $\mathrm{X_2Y(g)}$ with its equilibrium constant Kp is _____.

Assume x to be very very small.

JEE Main 2025 (Online) 23rd January Evening Shift

Options:

A. $x = \sqrt[3]{\frac{\mathrm{Kp}}{\mathrm{p}}}$

B. $x = \sqrt[3]{\frac{\mathrm{Kp}}{2\mathrm{p}}}$

C. $x = \sqrt[3]{\frac{2\mathrm{Kp}^2}{\mathrm{p}}}$

D. $x = \sqrt[3]{\frac{2\mathrm{Kp}}{\mathrm{p}}}$

Answer: C

Solution:

To determine the relationship between the degree of dissociation (x) of $\mathrm{X_2Y(g)}$ and its equilibrium constant K_p, let's analyze the reaction:

$\mathrm{X_2Y(g) \rightleftharpoons X_2(g) + \frac{1}{2}Y_2(g)}$

Initial and Change in Moles

Initially, we start with 1 mole of $\mathrm{X_2Y}$.

At equilibrium:

$\mathrm{X_2Y}$ is $(1 - x)$ moles.

$\mathrm{X_2}$ is x moles.

$\frac{1}{2}\mathrm{Y_2}$ is $\frac{x}{2}$ moles.

Partial Pressures

The total pressure $\mathrm{P_{total}}$ is given by:

$\mathrm{P_{X_2Y}} = \frac{1-x}{1+\frac{x}{2}} \times \mathrm{P}$

$\mathrm{P_{X_2}} = \frac{x}{1+\frac{x}{2}} \times \mathrm{P}$

$\mathrm{P_{Y_2}} = \frac{x/2}{1+\frac{x}{2}} \times \mathrm{P}$

Equilibrium Constant Expression

The equilibrium constant K_p can be written in terms of partial pressures:

$$K_p = \frac{\left(\frac{x}{1+\frac{x}{2}} \cdot \mathrm{P}\right) \cdot \left(\frac{\frac{x}{2}}{1+\frac{x}{2}} \cdot \mathrm{P}\right)^{1/2}}{\frac{1-x}{1+\frac{x}{2}} \cdot \mathrm{P}}$$

Simplifying the expression, considering x to be very small, results in:

$$K_p = \left(\frac{x}{1-x}\right)\left(\frac{x}{2\left(1+\frac{x}{2}\right)}\right)^{1/2} \cdot \mathrm{P}^{\frac{1}{2}}$$

For small x, $(1-x) \approx 1$, thus:

$$K_p = \frac{x^{3/2}}{\frac{1}{3}} \cdot \mathrm{P}^{\frac{1}{2}}$$

$$x^{3/2} = \frac{K_p \cdot 2^{1/2}}{\mathrm{P}^{1/2}}$$

Finally, by cubing both sides:

$$x^3 = \frac{2 \cdot K_p^2}{\mathrm{P}}$$

$$x = \left(\frac{2 \cdot K_p^2}{\mathrm{P}}\right)^{1/3}$$

Thus, the degree of dissociation x relates to K_p by the equation:

$$x = \left(\frac{2 \cdot K_p^2}{\mathrm{P}}\right)^{1/3}$$

Question19

For the reaction,


$$\mathrm{H_2(\,g) + I_2(\,g) \rightleftharpoons 2HI(\,g)}$$

Attainment of equilibrium is predicted correctly by :


JEE Main 2025 (Online) 24th January Evening Shift

Options:


A.




B.



C.



D.



Answer: C

Solution:

$H_2 + I_2 \rightleftharpoons 2HI$

Concentration of H_2 and I_2 decreases untill equilibrium condition and concentration of HI increases till equilibrium condition and after equilibrium concentration of all the reactant and products remain constant.Correct option (2)

Question20

At temperature T, compound $AB_{2(g)}$ dissociates as $AB_{2(g)} \rightleftharpoons AB_{(g)} + \frac{1}{2}B_{2(g)}$ having degree of dissociation x (small compared to unity). The correct expression for x in terms of K_p and p is:

JEE Main 2025 (Online) 29th January Morning Shift

Options:

A.

$\sqrt{K_p}$

B. $\sqrt[3]{\frac{2K_p^2}{p}}$

C. $\sqrt[3]{\frac{2K_p}{p}}$

D. $\sqrt[4]{\frac{2K_p}{p}}$

Answer: B

Solution:

$AB_2(g) \rightleftharpoons AB_{(g)} + \frac{1}{2}B_2(g)$

Degree of dissociation $\rightarrow x$ (small compared to unity)

Equilibrium constant interms of pressure $\rightarrow k_p$

Partial pressure of each gas $\rightarrow P_{AB_2}, P_{AB}$ and P_{B_2}

Total pressure $\rightarrow P$

ICE Table:

	$AB_{2(g)} \rightleftharpoons$	$AB_{(g)}$ +	$\frac{1}{2}B_{2(g)}$
Initial moles	1	0	0
Change moles	x	$+x$	$+\frac{1}{2}x$
Equilibrium moles	$1-x$	x	$\frac{1}{2}x$

Total moles at equilibrium $= 1 - x + x - \frac{1}{2}x = 1 - \frac{1}{2}x$

$$\text{partial pressure at equilibrium} = \frac{\text{moles at equilibrium}}{\text{total moles at equilibrium}} \times \text{total pressure}$$

Partial pressure of $AB_2(g)$

$P_{AB_2} = \left(\frac{1-x}{1+\frac{x}{2}}\right)P$

Partial pressure of $AB(g)$

$P_{AB} = \left(\frac{x}{1+\frac{x}{2}}\right)P$

Partial pressure of $B_2(g)$

$P_{B_2} = \left(\frac{\frac{x}{2}}{1+\frac{x}{2}}\right)^P$

K_p for the equation $A_2(g) \rightleftharpoons AB_{(g)} + \frac{1}{2}B_2(g)$ can be written as

$k_p = \frac{P_{AB}P_{B_2}^{1/2}}{P_{AB_2}}$

$= \frac{\frac{x}{1+\frac{x}{2}}P1 \times \left(\frac{\frac{x}{2}}{1+\frac{x}{2}}P\right)^{1/2}}{\frac{1-x}{1+\frac{x}{2}}P}$

$= \frac{x \times \left(\frac{x}{1+\frac{x}{2}}P\right)^{1/2}}{1-x}$

given, $x \lll 1$
So, $1 - x \approx 1$

$K_p = \frac{x \times \left(\frac{\frac{x}{2}}{1+\frac{x}{2}}p\right)^{1/2}}{1}$

$= x \times \frac{\left(\frac{x}{2}\right)^{1/2}}{\left(\frac{2+x}{2}\right)^{1/2}}p^{1/2}$

$= \frac{x \times \frac{x^{1/2}}{2^{1/2}} \times p^{1/2}}{\frac{(2+x)^{1/2}}{2^{1/2}}}$

$= \frac{x^{3/2} \times p^{1/2}}{2^{1/2}} \times \frac{2^{1/2}}{(2+x)^{1/2}}$

$x \lll 1$
$2 + x \approx 2$

$= \frac{x^{3/2} \times p^{1/2}}{2^{1/2}}$
$= \left(\frac{x^3 p}{2}\right)^{1/2}$

$k_p = \left(\frac{x^3 p}{2}\right)^{1/2}$
Square on both sides,

$k_p^2 = \frac{x^3 p}{2}$
$x^3 p = 2k_p^2$
$x^3 = \frac{2k_p^2}{p}$
$x = \sqrt[3]{\frac{2k_p^2}{p}}$

Correct answer option (2) $\sqrt[3]{\frac{2K_p^2}{p}}$

Question21

Consider the equilibrium

$$\mathrm{CO(g) + 3H_2(\ g) \rightleftharpoons CH_4(\ g) + H_2O(\ g)}$$

If the pressure applied over the system increases by two fold at constant temperature then

(A) Concentration of reactants and products increases.

(B) Equilibrium will shift in forward direction.

(C) Equilibrium constant increases since concentration of products increases.

(D) Equilibrium constant remains unchanged as concentration of reactants and products remain same.

Choose the correct answer from the options given below :

JEE Main 2025 (Online) 29th January Evening Shift

Options:

A. (A) and (B) only

B. (A), (B) and (D) only

C. (B) and (C) only

D. (A), (B) and (C) only

Answer: A

Solution:

Given equilibrium

$$\underset{(n=4)}{CO_{(g)}} + 3H_2(g) \rightleftharpoons \underset{(n=2)}{CH_{4(g)}} + H_2O(g)$$

Temperature $\rightarrow$ Constant

Pressure $\rightarrow$ Increases by two fold.

It is based on Le Chatelier's principle. The principle states that if a change of condition is applied to a system in equilibrium, the system will shift in a direction that relieves the stress.

When pressure increases in a chemical equilibrium, the equilibrium shifts towards the side of the reaction with fewer moles of gas molecules. According to the principle, the reaction will favor the side that provides fewer gas molecules to counteract the increased pressure.

In this equilibrium, fewer gas molecules are on the produce side and on increase in pressure, equilibrium shifts towards products - forwarded reaction occurs. As a result, concentration of the products increases.

A) Concentration of reactants and products increases.

$-$ Correct

With increase in pressure, at constant temperature, the concentration of both reactants and products will increase because the same amount of gas is now in a smaller volume.

(B) Equilibrium will shift in forward direction

Correct

Increase in pressure causes product formation as the equilibrium shift is in the forward direction.

(C) Equilibrium constant increases since concentration of products increases.

Not correct.

The expression for equilibrium constant (K_c) for the given equilibrium can be written as

$K_c = \frac{[CH_4][H_2O]}{[CO][H_2]^3}$

Equilibrium constant is not changed with increase in pressure and increase in concentration of reactants or products.

So, the equilibrium constant value is does not affected by change in concentration. If the change increase in concentration of product takes place, the equilibrium position will be shifted to counteract the change. So, equilibrium const value does not increase. Statement is incorrect.

(D) Equilibrium constant remains unchanged as concentration of reactants and products remain same.

State is incorrect.

In the reaction, increase in pressure causes changes in the concentration of rectants and products but the equilibrium constant will not change.

Statements (A) and (B) are correct.

Question22

Consider the following chemical equilibrium of the gas phase reaction at a constant temperature : $\mathrm{A(g)} \rightleftharpoons \mathrm{B(g)} + \mathrm{C(g)}$

If p being the total pressure, K_p is the pressure equilibrium constant and α is the degree of dissociation, then which of the following is true at equilibrium?

JEE Main 2025 (Online) 2nd April Evening Shift

Options:

A.

If K_p value is extremely high compared to p, α becomes much less than unity

B. When p increases α increases

C. If p value is extremely high compared to $\mathrm{K_p}$, $\alpha \approx 1$

D. When p increases α decreases

Answer: D

Solution:

	$\mathrm{A(g)} \rightleftharpoons$	$\mathrm{B(g)}$ +	$\mathrm{C(g)}$
t = 0	a	0	0
t = t	a (1 – α)	aα	aα

a moles of $A(g)$ taken initially and at time

Now moles fraction of $\mathrm{A(g)}, \mathrm{B(g)}$ and $\mathrm{C(g)}$ are

$$X_A = \frac{a - a\alpha}{a + a\alpha} = \frac{1-\alpha}{1+\alpha}$$
$$X_B = \frac{a\alpha}{a + a\alpha} = \frac{\alpha}{1+\alpha}$$
$$X_C = \frac{a\alpha}{a + a\alpha} = \frac{\alpha}{1+\alpha}$$

Now if P is total pressure then partial pressure of $A(g), B(g)$ and $C(g)$ are

$$P_A = \left(\frac{1-\alpha}{1+\alpha}\right)P$$
$$P_B = \left(\frac{\alpha}{1+\alpha}\right)P$$
$$P_C = \left(\frac{\alpha}{1+\alpha}\right)P$$

$$K_P = \frac{\left(\frac{\alpha}{1+\alpha}\right)P\left(\frac{\alpha}{1+\alpha}\right)P}{\left(\frac{1-\alpha}{1+\alpha}\right)P}$$
$$K_P = \frac{\alpha^2 P}{1-\alpha^2}$$

As K_P is only function of temperature.

So as $P \uparrow \quad \alpha \downarrow$

Question23

Given below are two statements :

Statement I : A catalyst cannot alter the equilibrium constant (K_c) of the reaction, temperature remaining constant.

Statement II : A homogenous catalyst can change the equilibrium composition of a system, temperature remaining constant.

In the light of the above statements, choose the correct answer from the options given below

JEE Main 2025 (Online) 3rd April Morning Shift

Options:

A. Statement I is true but Statement II is false

B. Statement I is false but Statement II is true

C. Both Statement I and Statement II are true

D. Both Statement I and Statement II are false

Answer: A

Solution:

Let's analyze both statements:

Statement I: "A catalyst cannot alter the equilibrium constant (K_c) of the reaction, temperature remaining constant."

A catalyst speeds up both the forward and reverse reactions equally, allowing the system to reach equilibrium faster.

However, it does not change the energy difference between reactants and products (i.e., the Gibbs free energy change), which directly determines the equilibrium constant.

Therefore, this statement is true.

Statement II: "A homogenous catalyst can change the equilibrium composition of a system, temperature remaining constant."

A homogeneous catalyst acts in the same phase as the reactants and, like any catalyst, it only helps the reaction reach equilibrium more quickly.

It does not alter the equilibrium position, meaning the relative concentrations of reactants and products at equilibrium remain unchanged if the temperature is constant.

Thus, this statement is false.

In summary:

Statement I is true.

Statement II is false.

The correct answer is: Option A.

Question24

In the following system, $\mathrm{PCl}_5(\mathrm{~g}) \leftrightharpoons \mathrm{PCl}_3(\mathrm{~g})+\mathrm{Cl}_2(\mathrm{~g})$ at equilibrium, upon addition of xenon gas at constant T \& p , the concentration of

JEE Main 2025 (Online) 3rd April Morning Shift

Options:

A. $\mathrm{PCl}_5, \mathrm{PCl}_3 \& \mathrm{Cl}_2$ remain constant

B. PCl_3 will increase

C. Cl_2 will decrease

D. PCl_5 will increase

Answer: B

Solution:

When xenon gas, an inert gas, is added to the equilibrium system $\mathrm{PCl}_5(\mathrm{g}) \leftrightharpoons \mathrm{PCl}_3(\mathrm{g}) + \mathrm{Cl}_2(\mathrm{g})$ at a constant temperature and pressure, it affects the equilibrium according to Le Chatelier's principle. The addition of an inert gas at constant pressure effectively increases the volume of the system, reducing the concentration of the gases involved.

Since the system is at constant pressure, adding an inert gas increases the total volume, which favors the side of the reaction with more moles of gas. In this reaction, there are two moles of gas on the right side (PCl_3 and Cl_2) compared to one mole on the left (PCl_5). Thus, the equilibrium will shift toward the right to increase the number of moles and counteract the change, decreasing the concentration of PCl_5 and increasing the concentrations of PCl_3 and Cl_2.

However, due to the increase in overall volume, the individual concentrations of all species ($[\mathrm{PCl}_5], [\mathrm{PCl}_3], [\mathrm{Cl}_2]$) will decrease as a result of more space being available for the same amount of gas particles.

Question25

37.8 g N_2O_5 was taken in a 1 L reaction vessel and allowed to undergo the following reaction at 500 K

$$2\,N_2O_{5(\,g)} \rightleftharpoons 2\,N_2O_{4(\,g)} + O_{2(\,g)}$$

The total pressure at equilibrium was found to be 18.65 bar.

Then, $Kp =$ _____ $\times 10^{-2}$ [nearest integer]

Assume N_2O_5 to behave ideally under these conditions.

Given: $R = 0.082$ **bar** $Lmol^{-1}\,K^{-1}$

JEE Main 2025 (Online) 24th January Morning Shift

Answer: 962

Solution:

$$\text{Initial pressure of } N_2O_5$$

$$= \frac{\frac{37.8}{108} \times 0.082 \times 500}{1} = 14.35\text{bar}$$

$$2\,N_2O_5 \rightleftharpoons 2\,N_2O_4 + O_2$$

$t = 0 \quad 14.35$
$t = eq \quad 14.35 - 2P \quad 2P \quad P$
$P_{Total} \quad \text{at eqb} \; = 14.35 + P = 18.65$
$P = 4.3$
$P_{N_2O_s} = 5.75\text{bar}$
$P_{N_2O_4} = 8.6\text{bar}$
$P_{O_2} = 4.3\text{bar}$
$k_p = \frac{(8.6)^2 \times (4.3)}{(5.75)^2} = 9.619 = x \times 10^{-2}$
$x = 961.9 \approx 962$

Question26

Consider the following equilibrium,

$$CO(\,g) + 2H_2(\,g) \rightleftharpoons CH_3OH(\,g)$$

0.1 mol of CO along with a catalyst is present in a $2dm^3$ flask maintained at 500 K . Hydrogen is introduced into the flask until the pressure is 5 bar and 0.04 mol of CH_3OH is formed. The K_p^θ is _____ $\times 10^{-3}$ (nearest integer).

Given : $R = 0.08\ dm^3$ **bar** $K^{-1}\ mol^{-1}$

Assume only methanol is formed as the product and the system follows ideal gas behaviour.

JEE Main 2025 (Online) 2nd April Morning Shift

Answer: 74

Solution:

	$CO(g)$	$+ 2H_2(g) \rightleftharpoons$	$CH_3OH(g)$
t = 0	0.1 mol	a mol	–
t_{eq}	0.1 - x	a - 2x	x = 0.04
	= 0.06	= a - 0.08	
		= 0.23 - 0.08	
		= 0.15 mole	

$V = 2L$
$T = 500K$
$P_{\text{total}} = 5 \text{ bar}$
$n_{\text{Total}} = 0.25 = \frac{1}{4} \text{ mol}$
$P_{\text{total}} = n_{\text{total}} \times \frac{RT}{V}$
$\Rightarrow 5 = (0.06 + a - 0.08 + 0.04) \times \frac{0.08 \times 500}{2}$
$\Rightarrow 10 = (0.02 + a) \times 0.08 \times 500$
$\Rightarrow \text{a} = 0.25 - 0.02 = 0.23 \text{ mol}$
$\text{K}_\text{P} = \frac{\text{X}_{\text{CH}_3\text{OH}}}{\text{X}_{\text{CO}} \times \text{X}_{\text{H}_2}^2} \times \frac{1}{(\text{P}_\text{T})^2} = \frac{0.04}{0.06 \times (0.15)^2} \times \left[\frac{1/4}{5}\right]^2$
$= \frac{4}{6 \times (0.15)^2 \times 16} \times \frac{1}{25}$
$= \frac{100 \times 100}{24 \times 225 \times 25} = \frac{100 \times 100}{135000}$
$= 0.074 = 74 \times 10^{-3}$

Question27

The equilibrium constant for decomposition of $H_2O(g)$

$$H_2O(g) \rightleftharpoons H_2(g) + \tfrac{1}{2}O_2(g) \quad (\Delta G^\circ = 92.34\,\text{kJ mol}^{-1})$$

is 8.0×10^{-3} at 2300 K and total pressure at equilibrium is 1 bar. Under this condition, the degree of dissociation (α) of water is _____ $\times 10^{-2}$ (nearest integer value).

[Assume α is negligible with respect to 1]

JEE Main 2025 (Online) 8th April Evening Shift

Answer: 5

Solution:

$H_2O(g) \rightleftharpoons H_{2(g)} + \frac{1}{2}O_{2(g)}$

	$H_2O(g)$	$\rightleftharpoons H_{2(g)}$	$+\frac{1}{2}O_{2(g)}$
$t = 0$	1 mole		
$t = t_{eq}$	$1-\alpha$	α	$\frac{\alpha}{2}$

$n_T = 1 + \frac{\alpha}{2} \simeq 1 (\alpha \ll 1)$

$k_P = \frac{P_{H_2} \cdot P_{O_2}^{1/2}}{P_{H_2O}} = \frac{(\alpha \cdot P)(\frac{\alpha}{2}P)^{\frac{1}{2}}}{(1-\alpha)P}$

$P = 1$

$8 \times 10^{-3} = \frac{\alpha^{3/2}}{\sqrt{2}}$

$\alpha^{3/2} = 8\sqrt{2} \times 10^{-3}$

$\alpha^3 = 128 \times 10^{-6}$

$\alpha = \sqrt[3]{128} \times 10^{-2}$

$= 5.03 \times 10^{-2}$

Question28

For the reaction $N_2O_4(g) \rightleftharpoons 2NO_2(g)$, Kp = 0.492atm at 300K. K_c for the reaction at same temperature is $\times 10^{-2}$.

(Given : R = 0.082L atm$mol^{-1}K^{-1}$)

[29-Jan-2024 Shift 1]

Answer: 2

Solution:

$$K_p = K_C \cdot (RT)^{\Delta n_g}$$

$$\Delta n_g = 1$$

$$\Rightarrow K_c = \frac{K_P}{RT} = \frac{0.492}{0.082 \times 300} = 2 \times 10^{-2}$$

Question29

The following concentrations were observed at 500K for the formation of NH_3 from N_2 and H_2. At equilibrium :

$[N_2] = 2 \times 10^{-2}$M, $[H_2] = 3 \times 10^{-2}$M and $[NH_3] = 1.5 \times 10^{-2}$M. Equilibrium constant for the reaction is___

[29-Jan-2024 Shift 2]

Answer: 417

Solution:

$$K_C = \frac{[NH_3]^2}{[N_2][H_2]^3}$$

$$K_C = \frac{(1.5 \times 10^{-2})^2}{(2 \times 10^{-2}) \times (3 \times 10^{-2})^3}$$

$$K_C = 417$$

Question30

For the given reaction, choose the correct expression of K_C from the following :-

$$Fe_{(aq)}^{3+} + SCN_{(aq)}^{-} \rightleftharpoons (FeSCN)_{(aq)}^{2+}$$

[31-Jan-2024 Shift 1]

Options:

A.

$$K_C = \frac{[FeSCN^{2+}]}{[Fe^{3+}][SCN^-]}$$

B.

$$K_C = \frac{[Fe^{3+}][SCN^-]}{[FeSCN^{2+}]}$$

C.

$$K_C = \frac{[FeSCN^{2+}]}{[Fe^{3+}]^2[SCN^-]^2}$$

D.

$$K_C = \frac{[FeSCN^{2+}]^2}{[Fe^{3+}][SCN^-]}$$

Answer: A

Solution:

$$K_C = \frac{\text{Products ion conc.}}{\text{Reactants ion conc.}}$$

$$K_C = \frac{[FeSCN^{2+}]}{[Fe^{3+}][SCN^-]}$$

Question31

Given below are two statements :

Statement (I) : Aqueous solution of ammonium carbonate is basic.
Statement (II) : Acidic/basic nature of salt solution of a salt of weak acid and weak base depends on Ka and Kb value of acid and the base forming it.

In the light of the above statements, choose the most appropriate answer from the options given below :

[27-Jan-2024 Shift 1]

Options:

A.

Both Statement I and Statement II are correct

B.

Statement I is correct but Statement II is incorrect

C.

Both Statement I and Statement II are incorrect

D.

Statement I is incorrect but Statement II is correct

Answer: A

Solution:

Aqueous solution of $(NH_4)_2CO_3$ is Basic

pH of salt of weak acid and weak base depends on Ka and Kb value of acid and the base forming it

---


Question32

Which of the following is strongest Bronsted base?


[27-Jan-2024 Shift 1]

Options:


A.




B.



C.




D.



Answer: D

Solution:



Question33

The pH at which $Mg(OH)_2$[K sp = 1×10^{-11}] begins to precipitate from a solution containing 0.10M Mg^{2+} ions is____

[30-Jan-2024 Shift 1]

Answer: 9

Solution:

Precipitation when $Q_{sp} = K_{sp}$

$[Mg^{2+}][OH^-]^2 = 10^{-11}$

$0.1 \times [OH^-]^2 = 10^{-11} \Rightarrow [OH^-] = 10^{-5}$

$\Rightarrow pOH = 5 \quad \Rightarrow pH = 9$

Question34

The pH of an aqueous solution containing 1M benzoic acid (pKa = 4.20) and 1M sodium benzoate is 4.5. The volume of benzoic acid solution in 300 mL of this buffer solution is_____ mL.

[30-Jan-2024 Shift 2]

Answer: 100

Solution:

	1M Benzoic acid	+	1M Sodium Benzoate
	(V_a ml)		(V_s ml)
Millimole	$V_a \times 1$		$V_s \times 1$

$pH = 4.5$

$$pH = pka + \log \frac{[salt\ ]}{[\ acid\ ]}$$

$$4.5 = 4.2 + \log\left(\frac{V_s}{V_a}\right)$$

$$\frac{V_s}{V_a} = 2 \ldots\ldots(1)$$

$V_s + V_a = 300 \ldots\ldots(2)$

$V_a = 100\ ml$

Question35

Ka for CH_3COOH is 1.8×10^{-5} and Kb for NH_4OH is 1.8×10^{-5}. The pH of ammonium acetate solution will be

[1-Feb-2024 Shift 1]

Answer: 7

Solution:

$$pH = \frac{pK_w + pK_a - pK_b}{2}$$

$pK_a = pK_b$

$$\Rightarrow pH = \frac{pK_w}{2} = 7$$

Question36

Solubility of calcium phosphate (molecular mass, M) in water is W_g per 100mL at 25°C. Its solubility product at 25°C will be approximately.

[1-Feb-2024 Shift 2]

Options:

A.

$10^7\left(\frac{W}{M}\right)^3$

B.

$10^7\left(\frac{W}{M}\right)^5$

C.

$10^3\left(\frac{W}{M}\right)^5$

D.

$10^5\left(\frac{W}{M}\right)^5$

Answer: B

Solution:

$$S = \frac{W \times 10}{M}$$

$$Ca_3(PO_4)_2(s) \rightleftharpoons \underset{3s}{3Ca^{2+}(aq.)} + \underset{2s}{2PO_4^{3-}(aq.)}$$

$$S = \frac{W \times 1000}{M \times 100} = \frac{W \times 10}{M}$$

$$K_{sp} = (3s)^3(2s)^2$$

$$= 108\, s^5$$

$$= 108 \times 10^5 \times \left(\frac{W}{M}\right)^5$$

$$= 1.08 \times 10^7\left(\frac{W}{M}\right)^5$$

Question37

The dissociation constant of acetic is $x \times 10^{-5}$. When 25 mL of 0.2M CH_3COONa solution is mixed with 25 mL of 0.02 M CH_3COOH solution, the pH of the resultant solution is found to be equal to 5 . The value of x is___
[24-Jan-2023 Shift 1]

Answer: 10

Solution:

Solution:
Buffer of HOAc and NaOAc

$$pH = pKa + \log \frac{0.1}{0.01}$$

$$5 = pKa + 1$$

$$pKa = 4$$


$$Ka = 10^{-4}$$

$$x = 10$$

Question38

If the pKa of lactic acid is 5 , then the pH of 0.005M calcium lactate solution at 25°C is _____$\times 10^{-1}$ (Nearest integer)

Lactic acid



[24-Jan-2023 Shift 2]

Answer: 85

Solution:

Solution:
Concentration of calcium lactate $= 0.005M$,: concentration of lactate ion $= (2 \times 0.005)M$. Calcium lactate is a salt of weak acid + strong base ∴ Salt hydrolysis will take place.

$$pH = 7 + \frac{1}{2}(pKa + \log C)$$

$$= 7 + \frac{1}{2}(5 + \log(2 \times 0.005))$$

$$= 7 + \frac{1}{2}[5 - 2\log 10] = 7 + \frac{1}{2} \times 3 = 8.5 = 85 \times 10^{-1}$$

Question39

A litre of buffer solution contains 0.1 mole of each of NH_3 and NH_4Cl. On the addition of 0.02 mole of HCl by dissolving gaseous HCl, the pH of the solution is found to be _____ $\times 10^{-3}$(Nearest integer)

[. Given : $pK_b(NH_3) = 4.745$

$\log 2 = 0.301$

$\log 3 = 0.477$

T = 298K]

[25-Jan-2023 Shift 1]

Answer: 9079

Solution:

Solution:
In resultant solution
$n_{NH_3} = 0.1 - 0.02 = 0.08$

$n_{NH_4Cl} = n_{NH_4^+} = 0.1 + 0.02 = 0.12$

$$pOH = pK b_b + \log \frac{[NH_4^+]}{[NH_3]}$$

$$= 4.745 + \log \frac{0.12}{0.08}$$

$$= 4.745 + \log \frac{3}{2}$$

$$= 4.745 + 0.477 - 0.301$$

$pOH = 4.921$

$pH = 14 - pH$

$= 9.079$

Question40

When the hydrogen ion concentration $[H^+]$changes by a factor of 1000 , the value of pH of the solution _____.
[25-Jan-2023 Shift 2]

Options:

A. increases by 1000 units

B. decreases by 3 units

C. decreases by 2 units

D. increases by 2 units

Answer: B

Solution:

Solution:

$\Delta [H^+] = 1000$

$\Delta pH = -\log \Delta [H^+] = -\log 10^3$

$= -3$

Question41

Match List I with List II

List I (Amines)	List II (pK_b)
A. Aniline	I. 3.25
B. Ethanamine	II. 3.00
C. N-Ethylethanamine	III. 9.38
D. N, N-Diethylethanamine	IV. 3.29

Choose the correct answer from the options given below :-
[25-Jan-2023 Shift 2]

Options:

A. A-I, B-IV, C-II, D-III

B. A-III, B-II, C-I, D-IV


C. A-III, B-II, C-IV, D-I

D. A-III, B-IV, C-II, D-I

Answer: D

Solution:

Solution:



(A) Aniline (B) Ethanamine (C) N-Ethylethanamine (D) N, N-Diethylethanamine

Basic Strength $\alpha \frac{1}{pK_b}$

Order for pK_b : A > B > D > C

Question42

Millimoles of calcium hydroxyide required to produce 100 mL of the aqueous solution of pH 12 is $x \times 10^{-1}$. The value of x is _____(Nearest integer).
Assume complete dissociation.
[29-Jan-2023 Shift 1]

Answer: 5

Solution:

Solution:

$\because$ pH = 12

$\therefore [H^+] = 10^{-12}M$

$\therefore [OH^-] = 10^{-2}M$

$\therefore [Ca(OH)_2] = 5 \times 10^{-3}M$

$$5 \times 10^{-3} = \frac{\text{milli moles of Ca (OH)}_2}{100\,\text{mL}}$$

$$\text{milli moles of Ca(OH)}_2 = 5 \times 10^{-1}$$

Question43

600 mL of 0.01M HCl is mixed with 400 mL of 0.01M H_2SO_4. The pH of the mixture is _____ $\times 10^{-2}$. (Nearest integer)
[Given log 2 = 0.30, log 3 = 0.48
log 5 = 0.69 log 7 = 0.84
log 11 = 1.04]
[30-Jan-2023 Shift 1]

Answer: 186

Solution:

Solution:

Total milimoles of $H^+ = (600 \times 0.01) + (400 \times 0.01 \times 2)$

$= 14$

$$[H^+] = \frac{14}{1000} = 14 \times 10^{-3}$$


$\text{pH} = 3 - \log 14$

$= 1.86$

$= 186 \times 10^{-2}$

Question44

The correct order of pK_a values for the following compounds is:



[30-Jan-2023 Shift 2]

Options:

A. c > a > d > b

B. b > d > a > c

C. b > a > d > c

D. a > b> c > d

Answer: B

Solution:

Solution:
Due to $-M$ effect of $-NO_2$ group, it increases acidity $+M$ effect of $N(CH_3)_2$ decreases acidity.
Hyperconjugation of isopropyl decrease acidity
∴ order of acidic strength
(c) > (a) > (d) > (b)

Question45

The logarithm of equilibrium constant for the reaction $Pd^{2+} + 4Cl^- \rightleftharpoons PdCl_4^{2-}$ is _____ (Nearest integer)

Given : $\frac{2.303\,RT}{F} = 0.06V$

$Pd_{(aq)}^{2+} + 2e^- \rightleftharpoons Pd(s)$ $E^\circ = 0.83V$

$PdCl_4^{2-}(aq) + 2e^- \rightleftharpoons Pd(s) + 4Cl^-(aq)$

$E^\circ = 0.65V$

[31-Jan-2023 Shift 1]

Answer: 6

Solution:

Solution:
Sol. $\Delta G^\circ = -RT\ell n K$
$-nFE_{cell}^0 = -RT \times 2.303(\log_{10}K)$
$\frac{E_{Cell}^0}{0.06} \times n = \log K$ . . . (1)
Pd^{+2} (aq.) + not mathcal $Le^- \rightleftharpoons Pd(s)$, $E_{cat,red}^\circ = 0.83$
$Pd(s) + 4Cl^-$ (aq.) $\rightleftharpoons PdCl_4^{2-}, (aq) + 2e^-$, $E_{mat.ouir}^0 = 0.65$
Net Reaction $\rightarrow Pd^{2+}$ (aq.) $+ 4Cl^-$ (aq.) $\rightleftharpoons PdCl_4^{2-}$ (aq.)
$E_{cell}^\circ = E_{cat,red}^\circ - E_{Alode\ ,\ 0cd^a}^\circ$
$E_{cell}^\circ = 0.83 - 0.65$
$E_{cell}^\circ = 0.18$ . . . (2)
Also $n = 2$ . . . (3)
Using equation (1), (2) & (3)

$\log K = 6$

Question46

Incorrect statement for the use of indicators in acid-base titration is : [31-Jan-2023 Shift 2]

Options:

A. Methyl orange may be used for a weak acid vs weak base titration.

B. Methyl orange is a suitable indicator for a strong acid vs weak base titration

C. Phenolphthalein is a suitable indicator for a weak acid vs strong base titration

D. Phenolphthalein may be used for a strong acid vs strong base titration.

Answer: A

Solution:

Solution:
Methyl orange may be used for a strong acid vs strong base and strong acid vs weak base titration. Phenolpthalein may be used for a strong acid vs strong base and weak acid vs strong base titration.

Question47

At 298K, the solubility of silver chloride in water is $1.434 \times 10^{-3} gL^{-1}$. The value of $-\log K_{sp}$ for silver chloride is

(Given mass of Ag is $107.9 gmol^{-1}$ and mass of Cl is _____ $35.5 gmol^{-1}$) [31-Jan-2023 Shift 2]

Answer: 10

Solution:

Solution:

$$AgCl(s) \rightarrow \underset{S}{Ag^{+}}(aq.) + \underset{Cl}{Cl^{-}}(aq.)$$

$$K_{sp} = S^2 = \left(\frac{1.43}{143.4} \times 10^{-3}\right)^2 = 10^{-10}$$

$$-\log K_{sp} = 10$$

Question48

Consider the following reaction approaching equilibrium at 27°C and 1 atm pressure

$$A + B \underset{K_r = 10^2}{\overset{K_f = 10^3}{\rightleftharpoons}} C + D$$

The standard Gibb's energy change ($\Delta_r G^\circ$) at 27°C is (-) _____ kJ mol^{-1} (Nearest integer).

(Given : R = 8.3JK^{-1}mol^{-1} and ln 10 = 2.3)

[29-Jan-2023 Shift 1]

Answer: 6

Solution:

Solution:

$\because \Delta G^\circ = -RT \ln K_{eq}$

and $K_{eq} = \frac{K_f}{K_b}$

$\therefore K_{eq} = \frac{10^3}{10^2} = 10$

$\therefore \Delta G = -RT \ln 10$

$\Rightarrow -(8.3 \times 300 \times 2.3) = -5.7 \text{ kJ mole} = 6 \text{ kJ mole}^{-1}$(nearest integer)

Ans = 6

Question49

At 298K

$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$, $K_1 = 4 \times 10^5$

$N_2(g) + O_2(g) \rightleftharpoons 2\,NO(g)$, $K_2 = 1.6 \times 10^{12}$

$H_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons H_2O(g)$, $K_3 = 1.0 \times 10^{-13}$

Based on above equilibria, the equilibrium constant of the reaction,

$2NH_3(g) + \frac{5}{2}O_2(g) \rightleftharpoons 2\,NO(g) + 3H_2O(g)$is _____ $\times 10^{-33}$

(Nearest integer)

[29-Jan-2023 Shift 2]

Answer: 4

Solution:

Solution:

$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$, $K_1 = 4 \times 10^5$ . . . (i)

$N_2(g) + O_2(g) \rightleftharpoons 2\,NO(g)$, $K_2 = 1.6 \times 10^{12}$... (ii)

$H_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons H_2O(g), K_3 = 1.0 \times 10^{-13}$... (iii)

(ii) + 3 × (iii) − (i)

$2NH_3(g) + \frac{5}{2}O_2(g) \rightleftharpoons 2NO(g) + 3H_2O(g)$

$k_{eq} = \frac{k_2 \times k_3^3}{k_1} = \frac{1.6 \times 10^{12} \times (10^{-13})^3}{4 \times 10^5}$

$= \frac{1.6}{4} \times 10^{-32} = 4 \times 10^{-33}$

Question50

Consider the following equation:
$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g), \Delta H = -190$ kJ

The number of factors which will increase the yield of SO_3 at equilibrium from the following is _____

A. Increasing temperature
B. Increasing pressure
C. Adding more SO_2
D. Adding more O_2
E. Addition of catalyst
[30-Jan-2023 Shift 2]

Answer: 3

Solution:

Solution:
The yield of SO_3 at equilibrium will be due to:
B. Increasing pressure
C. Adding more SO_2
D. Adding more O_2

Question51

For reaction: $SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$ $K_P = 2 \times 10^{12}$ at 27°C and 1 atm pressure.

The K_c for the same reaction is _____ $\times 10^{13}$. (Nearest integer)

(Given R = 0.082L atm K^{-1} mol^{-1})
[31-Jan-2023 Shift 1]

Answer: 1

Solution:

Solution:

$SO_{2(g)} + \frac{1}{2}O_{2(g)} \rightleftharpoons SO_{3(g)}$

$K_p = 2 \times 10^{12}$ at 300K

$K_P = K_C \times (RT)^{\Delta n_g}$

$2 \times 10^{12} = K_c \times (0.082 \times 300)^{-1/2}$

$K_C = 9.92 \times 10^{12}$

$K_C = 0.992 \times 10^{13}$

Ans. 1

Question52

For independent process at 300 K.

Process	$\Delta H / kJ\,mol^{-1}$	$\Delta S / JK^{-1}$
A	−25	−80
B	−22	40
C	25	−50
D	22	20

The number of non-spontaneous process from the following is____
[24-Jan-2023 Shift 1]

Answer: 2

Solution:

Solution:

$\Delta G = \Delta H - T\Delta S$

$A : \Delta G(Jmol^{-1}) = -25 \times 10^3 + 80 \times 300 : -ve$

$B : \Delta G(Jmol^{-1}) = -22 \times 10^3 - 40 \times 300 : -ve$

$C : \Delta G(Jmol^{-1}) = 25 \times 10^3 + 300 \times 50 : +ve$

$D : \Delta G(Jmol^{-1}) = 22 \times 10^3 - 20 \times 300 : +ve$

Processes C and D are non-spontaneous.

Question53

Water decomposes at 2300K
$H_2O(g) \rightarrow H_2(g) + \frac{1}{2}O_2(g)$
The percent of water decomposing at 2300K and 1 bar is _____ (Nearest integer).

Equilibrium constant for the reaction is 2×10^{-3} at 2300K
[29-Jan-2023 Shift 1]

Answer: 2

Solution:

Solution:

$H_2O(g) \rightleftharpoons H_2(g) + \frac{1}{2}O_2(g)$

$P_0[1-\alpha] \quad P_0\alpha \quad \frac{P_0\alpha}{2}$ partial pr. at eq.

$P_0\left[1+\frac{\alpha}{2}\right] = 1 \ldots$ (i)

$K_p = \frac{(P_{H_2})(P_{O_2})^{1/2}}{P_{H_2O}}$

$\frac{(P_0\alpha)\left(\frac{P_0\alpha}{2}\right)^{1/2}}{P_0[1-\alpha]} = 2 \times 10^{-3}$

since α is negligible w.r.t 1 so $P_0 = 1$ and $1-\alpha \approx 1$

$\frac{\alpha\sqrt{\alpha}}{\sqrt{2}} = 2 \times 10^{-3}$

$\alpha^{3/2} = 2^{3/2} \times 10^{-3}$

$\alpha = 2^{3/2 \times 2/3} \times 10^{-3 \times 2/3}$

$\alpha = 2 \times 10^{-2}$ $\quad \%\alpha = 2\%$

Question54

At 25°C, the enthalpy of the following processes are given:

$H_2(g) + O_2(g) \rightarrow 2\,OH(g)\ \Delta H^0 = 78\ kJ\ mol^{-1}$

$H_2(g) + 1/2O_2(g) \rightarrow H_2O(g)\ \Delta H^0 = -242\ kJ\ mol^{-1}$

$H_2(g) \rightarrow 2H(g)\ \Delta H^0 = 436\ kJ\ mol^{-1}$

$1/2O_2(g) \rightarrow O(g)\ \Delta H^0 = 249\ kJ\ mol^{-1}$

What would be the value of X for the following reaction? _____ (Nearest integer)

$H_2O(g) \rightarrow H(g) + OH(g)\ \Delta H^0 = X\ kJ\ mol^{-1}$

[1-Feb-2023 Shift 1]

Answer: 499

Solution:

Solution:

$2H_2O(g) \rightarrow 2H_2(g) + O_2(g) \ + (242 \times 2)\,kJ\,mol^{-1}$

$H_2(g) + O_2(g) \rightarrow 2\,OH \ + 78\,kJ\,mol^{-1}$

$H_2(g) \rightarrow 2H \ + 436\,kJ\,mol^{-1}$

$2H_2O \rightarrow 2H + 2\,OH \ + 998\,kJ\,mol^{-1}$

$H_2O \rightarrow H + OH \ \ 998 \times \frac{1}{2} = +499\,kJ\,mol^{-1}$

Question55

(i) $X(g) \rightleftharpoons Y(g) + Z(g) K_{p1} = 3$

(ii) $A(g) \rightleftharpoons 2B(g) \ K_{p2} = 1$

If the degree of dissociation and initial concentration of both the reactants X (g) and A(g) are equal, then the ratio of the total pressure at equilibrium $\left(\frac{p_1}{p_2}\right)$ is equal to x : 1. The value of x is _____ (Nearest integer)

[1-Feb-2023 Shift 1]

Answer: 12

Solution:

Solution:

$x(g) \rightleftharpoons y(g) + z(g) \ \ k_{p_1} = 3$

Initial moles $n \ - \ -$

at equilibrium $n - \alpha n \ \ \alpha n \ \ \alpha n$

$$k_{p_1} = \frac{\left(\frac{\alpha}{1+\alpha} \times p_1\right)^2}{\frac{1-\alpha}{1+\alpha} p_1}$$

$$3 = \frac{\alpha^2 \times p_1}{1-\alpha^2}$$

$A(g) \rightleftharpoons \ 2B(g) \ k_{p_2} = 1$

Initial mole $n \ -$

at equilibrium $x - \alpha n \ 2\alpha n \ p_{total} = p_2$

$$k_{p_2} = \frac{\left(\frac{2\alpha}{1+\alpha} \times p_2\right)^2}{\frac{1-\alpha}{1+\alpha} \times p_2}$$

$$1 = \frac{4\alpha^2 \times p_2}{1-\alpha^2}$$

$$\frac{k_{p_1}}{k_{p_2}} = \frac{p_1}{4p_2}$$

$$\frac{3}{1} = \frac{p_1}{4p_2} \ \ \therefore p_1 : p_2 = 12 : 1$$

$x = 12$

Question56

The effect of addition of helium gas to the following reaction in equilibrium state, is : $PCI_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$
[1-Feb-2023 Shift 2]

Options:

A. the equilibrium will shift in the forward direction and more of Cl_2 and PCl_3 gases will be produced.

B. the equilibrium will go backward due to suppression of dissociation of PCl_5.

C. helium will deactivate PCl_5 and reaction will stop.

D. addition of helium will not affect the equilibrium.

Answer: 0

Solution:

Solution:
$PCI_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$
(Case 1 : At constant P - volume will increase so reaction will shift in forward direction then answer will be A
Case 2 : At constant volume no change in active mass so reaction will not shift in any direction then answer will be D.

Question57

For a concentrated solution of a weak electrolyte (K_{eq} = equilibrium constant) A_2B_3 of concentration ' c ', the degree of dissociation " α ' is
[6-Apr-2023 shift 1]

Options:

A. $\left(\frac{K_{eq}}{108c^4}\right)^{\frac{1}{5}}$

B. $\left(\frac{K_{eq}}{6c^5}\right)^{\frac{1}{5}}$

C. $\left(\frac{K_{eq}}{5c^4}\right)^{\frac{1}{5}}$

D. $\left(\frac{K_{eq}}{25c^2}\right)^{\frac{1}{5}}$

Answer: A

Solution:

Solution:

$A_2B_3\ (aq.) \rightleftharpoons 2A_{(aq.)}^{3+} + 3B_{(aq)}^{2-}$

$c(1-\alpha)\ \ 2c\alpha\ \ 3c\alpha$

$K_{eq} = \frac{[A^{3+}]^2[B^{2-}]^3}{[A_2B_3]} = \frac{4c^2\alpha^2 \times 27c^3\alpha^3}{c(1-\alpha)}$

$K_{eq} = = \frac{108c^5\alpha^5}{c}\alpha = \left(\frac{K_{eq}}{108c^4}\right)^{\frac{1}{5}}$

Question58

The number of correct statement/s involving equilibria in physical from the following is _____
[10-Apr-2023 shift 1]

Options:

A. Equilibrium is possible only in a closed system at a given temperature.

B. Both the opposing processes occur at the same rate.

C. When equilibrium is attained at a given temperature, the value of all its parameters

D. For dissolution of solids in liquids, the solubility is constant at a given temperature.

Answer: C

Solution:

Solution:
(A) is correct
(B) for equilibrium $r_f = r_b \Rightarrow$(B) is correct
(C) at equilibrium the value of parameters become constant of a given temperature and not equal $\Rightarrow$(C) is incorrect
(D) for a given solid solute and a liquid solvent solubility depends upon temperature only $\Rightarrow$(D) is correct

---

Question59

The equilibrium composition for the reaction $PCl_3 + Cl_2 \rightleftharpoons PCl_5$ at 298K is given below.
$[PCl_3]_{eq} = 0.2\ mol\ L^{-1}$ $[Cl_2]_{eq} = 0.1\ mol\ L^{-1}$,
$[PCl_5]_{eq} = 0.40\ mol\ L^{-1}$
If 0.2 mol of Cl_2 is added at the same temperature, the equilibrium concentrations of PCl_5 is _____ $\times 10^{-2}\ mol\ L^{-1}$.
Given : K_c for the reaction at 298K is 20
[6-Apr-2023 shift 2]

Answer: 48

Solution:

Solution:

$PCl_3 + Cl_2 \rightleftharpoons PCl_5$

$0.2M \quad (0.1+0.2)M \quad 0.4M$

$Eq^m \cdot 0.2-x \quad 0.3-x \quad 0.4+x$

$$\frac{(0.4+x)}{(0.2-x)(0.3-x)} = 20$$

$\Rightarrow x \simeq 0.086$

$[PCl_5]_{eq} = 0.486M = 48.6 \times 10^{-2}M$

Question60

A mixture of 1 mole of H_2O and 1 mole of CO is taken in a 10 litre container and heated to 725K. At equilibrium 40% of water by mass reacts with carbon monoxide according to the equation : $CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g)$. The equilibrium constant $K_e \times 10^2$ for the reaction is _____ (Nearest integer)
[11-Apr-2023 shift 1]

Answer: 44

Solution:

Solution:

$$\underset{1\text{ mole}}{CO(g)} + \underset{1\text{ mole}}{H_2O(g)} \rightleftharpoons CO_2(g) + H_2(g)$$

$$K_c = \frac{0.4 \times 0.4}{0.6 \times 0.6} = \frac{4}{9}$$

$$K_c \times 10^2 = \frac{4}{9} \times 100 = \frac{400}{9} = 44.44 \approx 44$$

Question61

4.5 moles each of hydrogen and iodine is heated in a sealed ten litre vessel. At equilibrium, 3 moles of HI were found. The equilibrium constant for $H_2(g) + I_2(g) \rightleftharpoons 2\,HI(g)$ is _____
[11-Apr-2023 shift 2]

Answer: 1

Solution:

Solution:


	$H_{2(g)}$	$+ I_{2(g)}$	$\rightleftharpoons$	$2HI_{(g)}$
$t = 0$	4.5	4.5		–
t_{eq}	3	3		3

$K_c = \frac{[HI]^2}{[H_2][I_2]} = \frac{(3)^2}{3 \times 3} = \frac{9}{9} = 1$

Question62

The titration curve of weak acid vs. strong base with phenolphthalein as indictor) is shown below. The $K_{\text{phenolphthalein}} = 4 \times 10^{-10}$

Given : log 2 = 0.3



The number of following statements/s which is/are correct about phenolphthalein is _____

[8-Apr-2023 shift 1]

Options:

A. It can be used as an indicator for the titration of weak acid with weak base.

B. It begins to change colour at pH = 8.4

C. It is a weak organic base

D. It is colourless in acidic medium

Answer: B

Solution:

Solution:
(B) $pk_n = -\log(4 \times 10^{-10}) = 9.4$
Indicator range
$\Rightarrow pk_{In} \pm 1$
i.e. 8.4 to 10.4
(D) In acidic medium, phenolphthalein is in unionized form and is colourless.

Question63

Given below are two statements :
Statement I : Methyl orange is a weak acid.
Statement II : The benzenoid form of methyl orange is more intense/deeply coloured than the quinonoid form.
In the light of the above statement, choose the most appropriate answer from the options given below:
[8-Apr-2023 shift 2]

Options:

A. Both statement I and statement II are incorrect

B. Both statement I and Statement II are correct


C. Statement I is correct but statement II is incorrect

D. Statement I is incorrect but statement II is correct

Answer: A

Solution:

Solution:
(i) Methy orange is weak base



(ii) So both statement are false

Question64

The solubility product of $BaSO_4$ is _____ 1×10^{-10} at 298K. The solubility of $BaSO_4$ in 0.1M K_2SO_4 (aq) solut is $\times 10^{-9} gL^{-1}$ (Nearest integer)
Given: Molar mass of $BaSO_4$ is $233 gmol^{-1}$
[8-Apr-2023 shift 2]

Answer: 233

Solution:

Solution:

$K_{sp} = x(x + 0.1) = 10^{-10}$

$0.1x = 10^{-10}$

$x = 10^{-9}M$

$x(\text{ in } g/l) = 233 \times 10^{-9}$

Question65

An analyst wants to convert 1L HCl of pH = 1 to a solution of HCl of pH 2. The volume of water needed to do this dilution is _____ mL. (Nearest integer) [12-Apr-2023 shift 1]

Answer: 9000

Solution:

Solution:

$(M_1 \times V_1)\ (M_2 \times V_2)$

$-1 \quad = \quad -2$

$10 \times 1\ 10 \times V_2$

$V_2 = 10L$

Water added $= 10 - 1$

$= 9$ Litre

$= 9000$ mL

Question66

25.0 mL of 0.050 M $Ba(NO_3)_2$ is mixed with 25.0 mL of 0.020M NaF . K_{sp} of BaF_2 is 0.5×10^{-6} at 298K. The ratio of $[Ba^{2+}][F^-]^2$ and K_{sp} is _____ . (Nearest integer) [13-Apr-2023 shift 1]

Answer: 5

Solution:

Solution:

$[Ba^{+2}] = \frac{25 \times 0.05}{50} = 0.025M$

$[F^-] = \frac{25 \times 0.02}{50} = 0.01M$

$[Ba^{+2}][F^-]^2 = 25 \times 10^{-7}$

$K_{sp} = 5 \times 10^{-7}$ (given)

Ratio $= \frac{[Ba^{+2}][F^-]^2}{K_{sp}} = 5$

Question67

20 mL of 0.1M NaOH is added to 50 mL of 0.1M acetic acid solution. The pH of the resulting solution is _____ $\times 10^{-2}$ (Nearest integer)

Given : pKa(CH_3COOH) = 4.76

log 2 = 0.30

log 3 = 0.48

[13-Apr-2023 shift 2]

Answer: 458

Solution:

Solution:

	$CH_3COOH + NaOH \rightarrow CH_3COONa + H_2O$			
Initially	5mmol	2mmol	0	0
after Rxn	3mmol	0	2 mmole	2 mmole

$$pH = pKa + \log_{10} \frac{[\text{ salt }]}{[\text{ acid }]}$$

$$pH = 4.76 + \log_{10} \frac{2}{3}$$

$$pH = 4.58 = 458 \times 10^{-2}$$

Question68

Which of the following statement(s) is/are correct?

(A) The pH of 1×10^{-8}M HCl solution is 8

(B) The conjugate hase of $H_2PH_4^-$ is HPO_4^{2-}

(C) K_w increases with increase in temperature.

(D) When a solution of a weak monoprotic acid is titrated against a strong base at half neutralisation point.

pH = $\frac{1}{2}pK_a$

Choose the correct answer from the options given below

[15-Apr-2023 shift 1]

Options:

A. (A), (B). (C)

B. (A), (D)

C. (B), (C)

D. (B), (C). (D)

Answer: D

Solution:

Solution:
(A) pH of 10^{-8}M HCl in acidic range (6.98).
(B) Conjugate Base of $H_2PO_4^-$is HPO_4^{2-}
(C) K_w increases with increasing Temperature, as the temperature increases, the dissociation of water increases.
(D) At half neutralization point, half of the acid is present in the from salt.
$pH = Pk_a + \log \frac{1}{1} = Pk_a$

Question69

For a reaction at equilibrium
$A(g) \rightleftharpoons B(g) + \frac{1}{2}C(g)$
the relation between dissociation constant (K), degree of dissociation (α) and equilibrium pressure (p) is given by:
[24-Jun-2022-Shift-1]

Options:

A. $K = \frac{\alpha^{\frac{1}{2}} p^{\frac{3}{2}}}{\left(1+\frac{3}{2}\alpha\right)^{\frac{1}{2}}(1-\alpha)}$

B. $K = \frac{\alpha^{\frac{3}{2}} p^{\frac{1}{2}}}{(2+\alpha)^{\frac{1}{2}}(1-\alpha)}$

C. $K = \frac{(\alpha p)^{\frac{3}{2}}}{\left(1+\frac{3}{2}\alpha\right)^{\frac{1}{2}}(1-\alpha)}$

D. $K = \frac{(\alpha p)^{\frac{3}{2}}}{(1+\alpha)(1-\alpha)^{\frac{1}{2}}}$

Answer: B

Solution:

Solution:

	$A(g)$	$\rightleftharpoons$	$B(g)$	$+$	$\frac{1}{2}C(g)$	Total Moles
Att = 0:	1		0		0	1
Att = t_{req}:	$1-\alpha$		α		$\frac{\alpha}{2}$	1
Mole Fraction:	$\frac{\alpha}{1+\frac{\alpha}{2}}$		$\left(\frac{1-a}{1+\frac{a}{2}}\right)$		$P\left(\frac{2}{2}\right)P$	

Now,

$$K_P \text{ or } K = \frac{P_B \times (P_C)^{\frac{1}{2}}}{P_A}$$

$$= \frac{\left(\frac{\alpha}{1+\frac{\alpha}{2}}\right)P \times \left[\left(\frac{\frac{\alpha}{2}}{1+\frac{\alpha}{2}}\right)P\right]^{\frac{1}{2}}}{\left(\frac{1-\alpha}{1+\frac{\alpha}{2}}\right)P}$$

$$= \frac{\left(\frac{2\alpha}{2+\alpha}\right)P \times \left[\left(\frac{\alpha}{2+\alpha}\right)P\right]^{\frac{1}{2}}}{\left(\frac{2(1-\alpha)}{2+\alpha}\right)P}$$

$$= \frac{\alpha}{1-\alpha} \times \left(\frac{\alpha P}{2+\alpha}\right)^{\frac{1}{2}}$$

$$= \frac{\alpha^{\frac{3}{2}} \cdot P^{\frac{1}{2}}}{(1-\alpha)(2+\alpha)^{\frac{1}{2}}}$$

Question70

PCl_5 dissociates as

$PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$

5 moles of PCl_5 are placed in a 200 litre vessel which contains 2 moles of N_2 and is maintained at 600 K. The equilibrium prẽššưre is 2.46 atm. The equilibrium constant K_p for the dissociation of PCl_5 is ____$\times 10^{-3}$. (nearest integer)

(Given : R = 0.082Latm^{-1}mol^{-1}; Assume ideal gas behaviour)

[24-Jun-2022-Shift-2]

Answer: 1107

Solution:

Solution:

	$PCl_5(g)$	$\rightleftharpoons$	$PCl_3(g)$ +	$Cl_2(g)$	Total Moles
At t = 0 :	5		0	0	5 + 2 (For N_2 gas) = 7
At t = t_{eq} :	5 – x		x	x	5 – x + x + x + 2 (For N_2 gas) = 7 + x
Mole Fraction :	$\frac{5-x}{7+x}$		$\frac{5-x}{7+x}$	$\frac{5-x}{7+x}$	
Partial pressure :	$\left(\frac{5-x}{7+x}\right)P$		$\left(\frac{x}{7+x}\right)P$	$\left(\frac{x}{7+x}\right)P$	

Here 2 moles of N_2 also present that is why 2 moles always have to add in total mole calculation.
At equilibrium,
Pressure $(P) = 2.46\,atm$
Volume $(V) = 200L$
Temperature $(T) = 600K$
∴ Applying ideal gas equation,
$PV = nRT$
$\Rightarrow 2.46 \times 200 = (7 + x) \times 0.082 \times 600$
$\Rightarrow x = 3$
Now,

$$K_P = \frac{P_{PCl_3} \times P_{Cl_2}}{P_{PCl_5}}$$

$$= \frac{\left[\frac{3}{7+3} \times 2.46\right]\left[\frac{3}{7+3} \times 2.46\right]}{\left[\frac{5-3}{7+3} \times 2.46\right]}$$

$$= \frac{\frac{3}{10} \times \frac{3}{10} \times (2.46)^2}{\frac{2}{10} \times 2.46}$$

$$= \frac{9}{20} \times 2.46$$

$$= 1107 \times 10^{-3}\,atm$$

Question71

The standard free energy change (ΔG°) for 50% dissociation of N_2O_4 into NO_2 at 27°C and 1 atm pressure is $-xJmol^{-1}$. The value of x is_____ (Nearest Integer)
[Given : .R = $8.31JK^{-1}mol^{-1}$, log 1.33 = 0.1239 ln 10 = 2.3]
[25-Jun-2022-Shift-1]

Answer: 710

Solution:

Solution:

	N_2O_4	$\rightleftharpoons$	$2NO_2$
t = 0	1 mol		
t = t	(1-0.5) mol		0.5×2 mol
	= 0.5 mol		1 mol

$$k_P = \frac{\left(\frac{1}{1.5} \times 1\right)^2}{\left(\frac{0.5}{1.5} \times 1\right)} = \frac{1}{0.75} = \frac{100}{75}$$

$= 1.33$

$\Delta G^0 = -RT\ell nk_P$

$= -8.31 \times 300 \times \ln(1.33) = -710.45 J/mol$

$= -710 J/mol$

Question72

$2\,NOCl(g) \rightleftharpoons 2\,NO(g) + Cl_2(g)$

In an experiment, 2.0 moles of NOCl was placed in a one-litre flask and the concentration of NO after equilibrium established, was found to be 0.4 mol/L. The equilibrium constant at 30°C is_____ $\times 10^{-4}$
[27-Jun-2022-Shift-1]

Answer: 125

Solution:

Solution:

	$2\,NOCl(g)$	$\rightleftharpoons$	$2\,NO(g)$	+	$Cl_2(g)$
At $t = 0$:	2		0		0
At $t = t_{eq}$:	$2 - 2x$		$2x$		0

Given that at equilibrium, concentration of $NO = 0.4\,mol/L$

$\therefore 2x = 0.4$

$\Rightarrow x = 0.2$

∴ Concentration of $NOCl$ at equilibrium,

$[NOCl]_{eq} = 2 - 2 \times 0.2 = 1.6$

and $[NO]_{eq} = 0.4$

and $[Cl_2]_{eq} = 0.2$

We know,

$$K_C = \frac{[NO]^2[Cl_2]}{[NOCl]^2}$$

$$= \frac{[0.4]^2[0.2]}{[1.6]^2}$$

$\Rightarrow K_C = 12.5 \times 10^{-3}$

$\Rightarrow K_C = 125 \times 10^{-4}$

Question73

4.0 moles of argon and 5.0 moles of PCl_5 are introduced into an evacuated flask of 100 litre capacity at 610K. The system is allowed to equilibrate. At equilibrium, the total pressure of mixture was found to be 6.0 atm. The K_p for the reaction is

[Given : .R = 0.082L atm K^{-1} mol^{-1}]
[29-Jun-2022-Shift-2]

Options:

A. 2.25

B. 6.24

C. 12.13

D. 15.24

Answer: A

Solution:

Solution:

				Total Moles:
	$PCl_5(g) \rightleftharpoons$	$PCl_3(g) +$	$Cl_2(g)$	
At t = 0:	5	0	0	5 + 4(For Ne gas) = 9
At $t = t_{eq}$:	5 − x	x	x	5 − x + x + x + 4(For Ne gas) = 9 + x
Mole Fraction:	$\frac{5-x}{9+x}$	$\frac{x}{9+x}$	$\frac{x}{9+x}$	
Partial Pressure:	$\left(\frac{5-x}{9+x}\right)P$	$\left(\frac{x}{9+x}\right)^p$	$\left(\frac{x}{9+x}\right)^P$	

Here 4 moles of inert gas argon also present.
∴ Total moles of mixture present at equilibrium,
$n_T = 5 + x + 4$
$= 9 + x$
At equilibrium, total pressure $(p_T) = 6$ atm
Volume $(v) = 100L$
Temperature $= 610K$
∴ using ideal gas equation,
$P_T V = n_T RT$
$\Rightarrow 6 \times 100 = (9 + x) \times 0.082 \times 610$
$\Rightarrow x = 3$
Now,

$$K_P = \frac{P_{PCl_3} \times P_{Cl_2}}{P_{PCl_5}}$$

$$= \frac{\left[\frac{3}{9+3} \times 6\right] \times \left[\frac{3}{9+3} \times 6\right]}{\left[\frac{5}{9+3} \times 6\right]}$$

$$= \frac{27}{12}$$

$$= \frac{9}{4}$$

$= 2.25$ atm
Note :
Inert gas always contribute to total mole and pressure calculation.

Question74

A box contains 0.90g of liquid water in equilibrium with water vapour at 27°C. The equilibrium vapour pressure of water at 27 °C is 32.0 Torr. When the volume of the box is increased, some of the liquid water evaporates to maintain the equilibrium pressure. If all the liquid water evaporates, then the volume of the box must be ____litre. [nearest integer]
(Given : R = 0.082L atm K^{-1} mol^{-1})
(Ignore the volume of the liquid water and assume water vapours behave as an ideal gas.)
[29-Jun-2022-Shift-2]

Answer: 29

Solution:

Solution:
We know, 760 Torr $= 1$ atm
$\therefore 32$ Torr $= \frac{32}{760}$ atm
As all the liquid water evaporates so entire water is in gaseous state.
$\therefore$ Weight of water vapour $= 0.9$g
$\therefore$ Moles of water vapour $(n) = \frac{0.9}{18}$
Pressure $(P) = \frac{32}{760}$ atm
Temperature $(T) = (27 + 273)K = 300K$
$R = 0.082L$ atm $K^{-1}mol^{-1}$
Given water vapour act as an ideal gas, so we can apply ideal gas equation.
From ideal gas equation,
$PV = nRT$
$\Rightarrow \frac{32}{760} \times v = \frac{0.9}{18} \times 0.082 \times 300$
$\Rightarrow v = 29L$

Question75

Solute A associates in water. When 0.7g of solute A is dissolved in 42.0g of water, it depresses the freezing point by 0.2°C. The percentage association of solute A in water, is :
[Given : Molar mass of A = 93gmol^{-1}. Molal depression constant of water is 1.86K kg mol^{-1}.]
[25-Jun-2022-Shift-2]

Options:

A. 50%

B. 60%

C. 70%

D. 80%

Answer: D

Solution:

Solution:
$\Delta T = ik_f \times m$

$0.2 = i \times 1.86 \times \frac{0.7}{93} \times \frac{1000}{42}$

$i = \frac{0.2 \times 93 \times 6}{1.86 \times 100}$

$i = 0.60$

$2A \rightleftharpoons A_2$

$1 - \alpha \quad \frac{\alpha}{2}$

$i = 1 - \alpha + \frac{\alpha}{2}$

$i = 1 - \frac{\alpha}{2}$

$1 - \frac{\alpha}{2} = 0.60$

$1 - 0.60 = \frac{\alpha}{2}$

$\alpha = 0.80$

Question76

Given below are two statements one is labelled as Assertion A and the other is labelled as Reason R :
Assertion A : The amphoteric nature of water is explained by using Lewis acid/base concept.
Reason R: Water acts as an acid with NH_3 and as a base with H_2S.
In the light of the above statements choose the correct answer from the options given below :
[25-Jun-2022-Shift-2]

Options:

A. Both A and R are true and R is the correct explanation of A.

B. Both A and R are true but R is NOT the correct explanation of A.

C. A is true but R is false.

D. A is false but R is true.

Answer: D

Solution:

Solution:

The amphoteric nature of water is explained by using Bronsted-Lowry acid base concept

$$\underset{\text{(acid)}}{H_2O} + NH_3 - OH^- + NH_4^+$$

$$\underset{\text{(base)}}{H_2O} + H_2S \;\; H_3O^+ + HS^-$$

Hence, A is false but R is true

Question77

50 mL of 0.1MCH_3COOH is being titrated against 0.1M NaOH. When 25 mL of NaOH has been added, the pH of the solution will be ____$\times 10^{-2}$. (Nearest integer) (Given : .pK_a(CH_3COOH) = 4.76)

log 2 = 0.30
log 3 = 0.48
log 5 = 0.69
log 7 = 0.84
log 11 = 1.04
[26-Jun-2022-Shift-1]

Answer: 476

Solution:

Solution:

$$CH_3COOH + NaOH \rightarrow CH_3COONa + H_2O$$

After adding 25 ml of NaOH volume of mixture $= 50 + 25 = 75$ ml

Initially,

Number of millimole of NaOH $= 25 \times 0.1 = 2.5$ mm

Number of millimole of $CH_3COOH = 50 \times 0.1 = 5$ mm

After nutrilisation,

Millimole of NaOH $= 0$

Millimole of $CH_3COOH = 5 - 2.5 = 2.5$ mm

Millimole of $CH_3COONa = 2.5$

After nutrilisation,

Concentration of $CH_3COOH = [CH_3COOH] = \frac{5-2.5}{75} = \frac{1}{30}$

Concentration of $CH_3COONa = [CH_3COONa] = \frac{2.5}{75} = \frac{1}{30}$

$$P^H = P^{Ka} + \log \frac{[CH_3COONa]}{[CH_3COOH]}$$

$$= 4.76 + \log \frac{\frac{1}{30}}{\frac{1}{30}}$$

$$= 4.76 + \log(1)$$

$$= 4.76 + 0$$

$$= 4.76$$

$$= 4.76 \times 10^{-2}$$

Question78

pH value of 0.001M NaOH solution is___
[27-Jun-2022-Shift-2]

Answer: 11

Solution:

Solution:
$[OH^-] = 0.001 = 10^{-3}M$
$[H^+][OH^-] = 10^{-14}$
$[H^+] = 10^{-11}$
$pH = -\log[H^+]$
$= -\log(10^{-11})$
$pH = 11$

Question79

A student needs to prepare a buffer solution of propanoic acid and its sodium salt with pH 4. The ratio of $\frac{[CH_3CH_2COO^-]}{[CH_3CH_2COOH].}$ required to make buffer is
Given : $K_a(CH_3CH_2COOH) = 1.3 \times 10^{-5}$
[28-Jun-2022-Shift-2]

Options:

A. 0.03

B. 0.13

C. 0.23

D. 0.33

Answer: B

Solution:

Solution:
$CH_3CH_2COOH \rightleftharpoons CH_3CH_2COO^- + H^+$
From Henderson equation
$$pH = pK_a + \log \frac{[CH_3CH_2COO^-]}{[CH_3CH_2COOH]}$$
$$4 = -\log 1.3 \times 10^{-5} + \log \frac{[CH_3CH_2COO^-]}{[CH_3CH_2COOH^2]}$$
$$-\log 10^{-4} = -\log 1.3 \times 10^{-5} + \log \frac{[CH_3CH_2COO^-]}{[CH_3CH_2COOH^-]}$$

$$-\log 10^{-4} = -\log 1.3 \times 10^{-5} \frac{[CH_3CH_2COOH]}{[CH_3CH_2COO^-]}$$

$$10^{-4} = 1.3 \times 10^{-5} \frac{[CH_3CH_2COOH^3]}{[CH_3CH_2COO^-]}$$

$$\frac{[CH_3CH_2COO^-]}{[CH_3CH_2COOH]} = 0.13$$

Question80

The solubility of AgCl will be maximum in which of the following? [29-Jun-2022-Shift-1]

Options:

A. 0.01M KCl

B. 0.01M HCl

C. 0.01M$AgNO_3$

D. Deionised water

Answer: D

Solution:

Solution:
In deionized water no common ion effect will take place so maximum solubility.

Question81

K_{a_1}, K_{a_2} and K_{a_3} are the respective ionization constants for the following reactions (a), (b) and (c).

(a) $H_2C_2O_4 \rightleftarrows H^+ + HC_2O_4^-$

(b) $HC_2O_4^- \rightleftarrows H^+ + HC_2O_4^{2-}$

(c) $H_2C_2O_4 \rightleftarrows 2H^+ + C_2O_4^{2-}$

The relationship between K_{a_1}, K_{a_2} and K_{a_3} is given as

[25-Jul-2022-Shift-2]

Options:

A. $K_{a_3} = K_{a_1} + K_{a_2}$

B. $K_{a_3} = K_{a_1} - K_{a_2}$

C. $K_{a_3} = K_{a_1} / K_{a_2}$

D. $K_{a_3} = K_{a_1} \times K_{a_2}$

Answer: D

Solution:

Solution:

$H_2C_2O_4 \rightleftharpoons 2H^+ + C_2O_4^{2-}$ K_{a3}

$H_2C_2O_4 \rightleftharpoons H^+ + HC_2O_4^-$ K_{a1}

$HC_2O_4^- \rightleftharpoons H^+ + C_2O_4^{2-}$ K_{a2}

$K_{a3} = \frac{[H^+]^2[C_2O_4^{2-}]}{[H_2C_2O_4]}$

$K_{a1} = \frac{[H^+][HC_2O_4^-]}{[H_2C_2O_4]}$, $K_{a2} = \frac{[H^+][C_2O_4^-]}{[HC_2O_4^-]}$

$K_{a3} = K_{a1} \times K_{a2}$

Question82

At 298K, the equilibrium constant is 2×10^{15} for the reaction :
$Cu(s) + 2Ag^+(aq) \rightleftharpoons Cu^{2+}(aq) + 2\,Ag(s)$
The equilibrium constant for the reaction
$\frac{1}{2}Cu^{2+}(aq) + Ag(s) \rightleftharpoons \frac{1}{2}\,Cu(s) + Ag^+(aq)$
is $x \times 10^{-8}$. The value of x is _____. (Nearest Integer)
[26-Jul-2022-Shift-1]

Answer: 2

Solution:

Solution:

$K_{eq}' = \frac{1}{\sqrt{K_{eq}}} = \frac{1}{\sqrt{2 \times 10^{15}}} = X \times 10^{-8}$

$\Rightarrow \frac{1}{\sqrt{20}} \times \frac{1}{10^7} = x \times 10^{-8}$

$\Rightarrow \frac{1}{\sqrt{20}} \times 10^{-7} = x \times 10^{-8}$

$\Rightarrow x = \frac{\sqrt{10}}{\sqrt{2}} = \sqrt{5} = 2.236$

$\simeq 2.24$

Question83

At 600K, 2 mol of NO are mixed with 1 mol of O_2.
$2NO_{(g)} + O_2(g) \rightleftarrows 2NO_2(g)$

The reaction occurring as above comes to equilibrium under a total pressure of 1 atm. Analysis of the system shows that 0.6 mol of oxygen are present at equilibrium. The equilibrium constant for the reaction is ____(Nearest integer)
[28-Jul-2022-Shift-2]

Answer: 2

Solution:

Solution:

$$2\,NO(g) + O_2(g) \rightleftharpoons 2NO_2(g)$$

at intial 2 1 0
at equilibrium $2 - 0.8$ 0.6 0.8
Partial pressure of $NO(g) = 1.2/2.6 \times 1$
Partial pressure of $O_2(g) = 0.6/2.6$
Partial pressure of $NO_2(g) = 0.8/2.6$

$$K_p = \left(P_{NO_2}\right)^2/(P_{NO})^2\left(P_{O_2}\right) = 0.8 \times 0.8 \times 2.6/1.2 \times 1.2 \times 0.6$$
$$= 1.925$$
$$\approx 2$$

Question84

20 mL of 0.1 M NH_4OH is mixed with 40 mL of 0.05 M HCl. The pH of the mixture is nearest to:
(Given:
$K_b(NH_4OH) = 1 \times 10^{-5}$, log 2 = 0.30, log 3 = 0.48, log 5 = 0.69, log 7 = 0.84, log 11 = 1.04
)
[25-Jul-2022-Shift-1]

Options:

A. 3.2

B. 4.2

C. 5.2

D. 6.2

Answer: C

Solution:

Solution:

$$NH_4OH + HCl \rightarrow NH_4Cl + H_2O$$

mmole 22

$[NH_4^+] = \frac{2\,mmole}{60\,ml} = \frac{1}{30}M$

$pH = \frac{pK_w - pK_b - \log C}{2} = \frac{14 - 5 + 1.48}{2} = 5.24$

Question85

Class XII students were asked to prepare one litre of buffer solution of pH 8.26 by their Chemistry teacher: The amount of ammonium chloride to be dissolved by the student in 0.2M ammonia solution to make one litre of the buffer is
(Given: $pK_b(NH_3) = 4.74$

Molar mass of $NH_3 = 17gmol^{-1}$

Molar mass of $NH_4Cl = 53.5gmol^{-1}$)
[26-Jul-2022-Shift-2]

Options:

A. 53.5g

B. 72.3g

C. 107.0g

D. 126.0g

Answer: C

Solution:

Solution:
For basic Buffer, $pOH = pK_b + \log \frac{[\text{ salt }]}{[\text{ Base }]}$ $pOH = 14 - 8.26 = 5.74$

$5.74 = 4.74 + \log \frac{[NH_4Cl]}{0.2}$

$[NH_4Cl] = 2M$

Moles of $NH_4Cl = 2 \times 1 = 2$ moles

Weight of $NH_4Cl = 2 \times 53.5 = 107g$

Question86

At 310K, the solubility of CaF_2 in water is $2.34 \times 10^{-3}g/100\,mL$. The solubility product of CaF_2 is _____ $\times 10^{-8}(mol/L)^3$. (Give molar mass : $CaF_2 = 78gmol^{-1}$)
[27-Jul-2022-Shift-1]

Answer: 0

Solution:

Solution:

$CaF_2 \overset{s}{\rightleftharpoons} \underset{s}{Ca^{2+}} + 2\ \underset{2s}{F^-}$

$K_{sp} = s(2s)^2$

$= 4s^3$

Solubility $(s) = 2.34 \times 10^{-3} g/100\,mL$

$= \frac{2 \cdot 34 \times 10^{-3} \times 10}{78}$ mole/lit

$= 3 \times 10^{-4}$ mole/lit

$\therefore K_{sp} = 4 \times (3 \times 10^{-4})^3$

$= 108 \times 10^{-12}$

$= 0.0108 \times 10^{-8} (mole/\ lit\ )^3$


$\therefore x \approx 0$

Question87


The plot of pH-metric titration of weak base NH_4OH vs strong acid HCl looks like : [27-Jul-2022-Shift-2]

Options:


A.




B.



C.




D.



Answer: A

Solution:

Solution:
NH_4OH is a weak base and HCl is a strong acid.
With the addition of HCl to NH_4OH, pH of solution will decrease gradually.
So, the correct graph should be



Question88

K_a for butyric acid (C_3H_7COOH) is 2×10^{-5}. The pH of 0.2M solution of butyric acid is _____ $\times 10^{-1}$. (Nearest integer)
[Given log 2 = 0.30]
[28-Jul-2022-Shift-1]

Answer: 27

Solution:

Solution:
K_a of Butyric acid $\Rightarrow 2 \times 10^{-5}$ PKa = 4.7
pH of 0.2M solution

$$pH = \frac{1}{2}pK_a - \frac{1}{2}\log C$$

$$= \frac{1}{2}(4 \cdot 7)\ \frac{1}{2}\log(0.2)$$

$$= 2.35 + 0.35 = 2.7$$

$$pH = 27 \times 10^{-1}$$

Question89

If the solubility product of PbS is 8×10^{-28}, then the solubility of PbS in pure water at 298K is $\times \times 10^{-16}$ mol L^{-1}. The value of x is _____. (Nearest Integer)
[Given : $\sqrt{2} = 1.41$]
[29-Jul-2022-Shift-1]

Answer: 282

Solution:

Solution:
$K_{sp} = S^2$
$S = \sqrt{K_{sp}} = \sqrt{8 \times 10^{-28}} = 2\sqrt{2} \times 10^{-14}$
$= 2.82 \times 10^{-14}$
$= 282 \times 10^{-16}$
∴ Ans. 282

Question90

200 mL of 0.01 MHCl is mixed with 400 mL of 0.01MH_2SO_4. The pH of the mixture is _____.
Given: log 2 = 0.30, log 3 = 0.48, log 5 = 0.70, log 7 = 0.84, log 11 = 1.04
[29-Jul-2022-Shift-2]

Options:

A. 1.14

B. 1.78

C. 2.34

D. 3.02

Answer: B

Solution:

Solution:
$$[H^+] = \frac{0.01 \times 200 + 2 \times 0.01 \times 400}{600}$$
$$= \frac{0.01 + 2 \times 0.01 \times 2}{3}$$
$$= \frac{0.01 + 0.04}{3}$$
$$= \frac{5}{3} \times 10^{-2}$$
$$pH = -\log[H^+]$$
$$= -\log\left(\frac{5}{3} \times 10^{-2}\right)$$

$$= -\left[\log \frac{5}{3} + \log 10^{-2}\right]$$
$$= -[\log 5 - \log 3 - 2]$$
$$= -0.7 + 0.48 + 2$$
$$= 2.48 - 0.7$$
$$= 1.78$$

Question91

The solubility of $Ca(OH)_2$ in water is [Given: The solubility product of $Ca(OH)_2$ in water $= 5.5 \times 10^{-6}$] [25 Feb 2021 Shift 2]

Options:

A. 1.11×10^{-2}

B. 1.11×10^{-6}

C. 1.77×10^{-2}

D. 1.77×10^{-6}

Answer: A

Solution:

Solution:
Let, solubility of $Ca(OH)_2$ in pure water $= S mol/L$

$Ca(OH)_2 \rightleftharpoons Ca^{2+} + 2OH^-$

$S mol/L \quad 2 \times S(mol/L)$

$= [Ca^{2+}][OH^-]^2 = S \times (2S)^2 = 4S^3 (mol/L)$

The expression of K_{sp} can also be written as,

$$K_{sp} = x^x \cdot y^y \cdot S^{x+y}$$
$$= 1^1 \cdot 2^2 \cdot S^{1+2}$$
$$= 4S^3$$

$$K_{sp} = x^x \cdot y^y \cdot S^{x+y}$$
$$= 1^1 \cdot 2^2 \cdot S^{1+2}$$
$$= 4S^3 \quad [\because \text{ For } Ca(OH)_2 : x = 1, y = 2]$$

x and y are the coefficients of cations and anions respectively

$$S = \left(\frac{K_{sp}}{4}\right)^{1/3} = \left(\frac{5.5 \times 10^{-6}}{4}\right)^{1/3}$$
$$= 1.11 \times 10^{-2} mol/L$$

Question92

The solubility product of PbI_2 is 8.0×10^{-9}. The solubility of lead iodide in 0.1 molar solution of lead nitrate is $x \times 10^{-6} mol/L$. The value of x is ..... (Rounded off to the nearest integer).

[Given, : $\sqrt{2} = 1.41$]
[24 Feb 2021 Shift 2]

Answer: 141

Solution:

Solution:
Given, $[K_{sp}]_{PbI_2} = 8 \times 10^{-9}$
To calculate solubility of PbI_2 in $0.1M$ solution of $Pb(NO_3)_2$.
(I) $\underset{0.1M}{Pb(NO_3)_2} \longrightarrow \underset{0.1M}{Pb^{2+}}(aq) + \underset{0.2M}{2NO_3^-}(aq)$
(II) $PbI_2(s) \rightleftharpoons Pb^{2+}(aq) + 2I^-(aq)$
S 2S
$\therefore \ [Pb^{2+}] = S + 0.1 \ \approx 0.1$
$\because \ S << 0.1$
Now, $K_{sp} = 8 \times 10^{-9}$
$[Pb^2][I^-]^2 = 8 \times 10^{-9}$
$0.1 \times (2S)^2 = 8 \times 10^{-9}$
$4S^2 = 8 \times 10^{-8} \Rightarrow S = 141 \times 10^{-6}M$
x = 141

Question93

The pH of ammonium phosphate solution, if pK_a of phosphoric acid and pK_b of ammonium hydroxide are 5.23 and 4.75 respectively, is ..... .
[26 Feb 2021 Shift 2]

Answer: 7

Solution:

Solution:
Phosphoric acid is a weak tribasic acid (H_3PO_4) and NH_4OH is a weak monoacidic base. So, hydrolysis of ammonium phosphate $(NH_4)_3PO_4$ can be shown as,
$(NH_4)_3PO_4 + 3H_2O \rightleftharpoons H_3PO_4 + 3NH_4OH$ [3 : 1 type salt]
So, $[H^+] - K_a \times \left(\frac{K_w}{K_a \times K_b}\right)^{12}$
$pH = pK_a + \frac{1}{2}[pK_w - pK_a - pK_b]$
$= 5.23 + \frac{1}{2}(14 - 5.23 - 4.75)$
[$\because pK_w = 14 H_2O$
$pK_a = 5.23(H_3PO_4)$
$pK_b = 4.75(NH_4OH)$]
$= 7.24 \sim eq7$

Question94

The solubility of AgCN in a buffer solution of pH = 3 is x. The value of x is..... .
[Assume : No cyano complex is formed; $K_{sp}(AgCN) = 2.2 \times 10^{-16}$ and $K_a(HCN) = 6.2 \times 10^{-10}$]
[25 Feb 2021 Shift 1]

Options:

A. 0.625×10^{-6}

B. 1.6×10^{-6}

C. 2.2×10^{-16}

D. 1.9×10^{-5}

Answer: D

Solution:

Solution:
pH of AgCN buffer solution $= 3$
$[H^+] = 10^{-3}$
$K_{sp}(AgCN) = 2.2 \times 10^{-16}$
$K_a[HCN] = 6.2 \times 10^{-10}$
$AgCN \rightleftharpoons Ag^+ + CN^- \ldots\ldots . K_{sp}$
$CN^- + H^+ \rightleftharpoons HCN \ldots\ldots \frac{1}{K_a}$
$AgCN + H^+ \rightleftharpoons HCN + Ag^+$
$K_{sp} \times \frac{1}{K_a} = \frac{[Ag^+][CN^-][HCN]}{[H^+][CN^-]}$
$[S] = \sqrt{\frac{K_{sp}[H^+]}{K_a}} \Rightarrow \frac{2.2 \times 10^{-16}}{6.2 \times 10^{-10}} = \frac{[S][S]}{10^{-3}}$
$[S]^2 = \frac{2.2 \times 10^{-16}}{6.2 \times 10^{-10}} \times 10^{-3}$
$S = 1.9 \times 10^{-5}$

Question95

A homogeneous ideal gaseous reaction $AB_2(g) \rightleftharpoons A(g) + 2B(g)$ is carried out in a 25L flask at 27°C. The initial amount of AB_2 was 1 mole and the equilibrium pressure was 1.9atm. The value of K_p is $x \times 10^{-2}$. The value of x is ..... .
[26 Feb 2021 Shift 1]

Answer: 73

Solution:

Solution:

$(\Sigma \text{ mole})_{t_{eq}} = 1 - x + x + 2x = (1 + 2x)$ Partial pressure $\frac{1-x}{1+2x}p$

(atm) $\frac{x}{1+2x}p$ $\frac{2x}{1+2x}p$

[$p =$ Total pressure at equilibrium $= 1.9\text{atm}$]

Now, at equilibrium $pV = (1 + 2x)RT$

$\Rightarrow 1 + 2x = \frac{pV}{RT} = \frac{1.9 \times 25}{0.082 \times 300} = 1.93$ [$V = 25\text{L}$, $R = 0.082\text{L atm mol}^{-1}\text{K}^{-1}$ $T = 300\text{K}$]

$\Rightarrow x = \frac{1.93 - 1}{2} = 0.465$

$\Rightarrow K_p = \frac{p_A \times p_B^2}{p_{AB_2}} \Rightarrow \frac{\left(\frac{x}{1+2x}p\right) \times \left(\frac{2x}{1+2x}p\right)^2}{\left(\frac{1-x}{1+2x}p\right)}$

$= \frac{4x^3 \times p^3}{(1+2x)^3} \times \frac{(1+2x)}{(1-x) \times p} = \frac{4x^3 \times p^2}{(1+2x)^2 \times (1-x)}$

$= \frac{4 \times (0.465)^3 \times (1.9)^2}{(1 + 2 \times 0.465)^2 \times (1 - 0.465)} = 0.7285\text{atm}$

$= 72.85 \times 10^{-2}\text{atm} \sim \text{eq} 73 \times 10^{-2} = x \times 10^{-2}$

$\therefore x = 73$

Question96

At 1990K and 1atm pressure, there are equal number of Cl_2 molecules and Cl atoms in the reaction mixture. The value of K_P for the reaction $Cl_2(g) \rightleftharpoons 2Cl(g)$ under the above conditions is $x \times 10^{-1}$. The value of x is___ (Rounded off to the nearest integer) [24feb2021shift1]

Answer: 5

Solution:

Solution:

$Cl_2(g) \rightleftharpoons 2Cl(g)$

Let mol of both of Cl_2 and Cl be x.

$P_{Cl} = \frac{x}{2x} \times 1 = \frac{1}{2}$

$P_{Cl_2} = \frac{x}{2x} \times 1 = \frac{1}{2}$

$\therefore K_p = \frac{\left(\frac{1}{2}\right)^2}{\frac{1}{2}} = \frac{1}{2} = 0.5 = 5 \times 10^{-1}.$

Question97

For the reaction A(g) → (B)(g), the value of the equilibrium constant at 300K and 1atm is equal to 100.0. The value of $\Delta_r G$ for the reaction at 300K and 1atm in $J\ mol^{-1}$ is $-xR$, where x is_____ (Rounded off to the nearest integer)
($R = 8.31\ J\ mol^{-1} K^{-1}$ and $\ln 10 = 2.3$)
[24feb2021shift1]

Answer: 1380

Solution:

Solution:

$$\Delta G^\circ = RT \ln K_p$$
$$= -R(300)(2)\ln(10)$$
$$= -R(300 \times 2 \times 2.3)$$
$$\Delta G^\circ = -1380R$$

Question98

The solubility of $CdSO_4$ in water is $8.0 \times 10^{-4} mol\ L^{-1}$. Its solubility in 0.01M H_2SO_4 solution is ..... $\times 10^{-6} mol\ L^{-1}$. (Round off to the nearest integer) (Assume that, solubility is much less than 0.01M)
[18 Mar 2021 Shift 2]

Answer: 64

Solution:

Solution:

Given, solubility in water $(S) = 8.0 \times 10^{-4} mol\ L^{-1}$

In pure water

$$K_{sp} = S^2 = (8 \times 10^{-4})^2 = 64 \times 10^{-8}$$

In 0.01 M H_2SO_4

$$H_2SO_4(aq) \longrightarrow \underset{0.02}{2H^+(aq)} + \underset{0.02}{SO_4^{2-}(aq)}$$

$$CdSO_4(s) \rightleftharpoons \underset{x}{Cd^{2+}(aq)} + \underset{(x+0.01)}{SO_4^{2-}(aq)}$$

$$K_{sp} = x(x + 0.01) = 64 \times 10^{-8}$$

$$x + 0.01 \cong 0.01M$$

So, $x(0.01) = 64 \times 10^{-8}$

$$x = 64 \times 10^{-6} M$$

Question99

The oxygen dissolved in water exerts a partial pressure of 20kPa in the vapour above water. The molar solubility of oxygen in water is ..... $\times 10^{-5}$mol d m^{-3}. (Round off to the nearest integer).
[Given, Henry's law constant (K_H) $= 8.0 \times 10^4$kPa for O_2, density of water with dissolved oxygen $= 1.0$kgd m^{-3}].
[17 Mar 2021 Shift 1]

Answer: 25

Solution:

Solution:
Given, partial pressure of $O_2 = 20$kPa
K_H (Henry's constant) $= 8 \times 10^4$kPa
From Henry's law,
$p(g) = [K_H]\chi_{O_2}$
where, χ_{O_2} = solubility of oxygen
$20 \times 10^3 = (8 \times 10^4 \times 10^3)\chi_{O_2}$
$\Rightarrow \chi_{O_2} = \dfrac{20}{8 \times 10^4}$
Solubility $= 2.5 \times 10^{-4} = 25 \times 10^{-5}$

Question100

Two salts A_2X and MX have the same value of solubility product of 4.0×10^{-12}. The ratio of their molar solubilities i.e $\dfrac{S(A_2X)}{S(MX)}$ = ……
(Round off to the nearest integer)
[16 Mar 2021 Shift 1]

Answer: 50

Solution:

Solution:
Let the solubility of A_2X be ' 'S'.
$A_2X(s) \rightleftharpoons 2A^+(aq) + X^{2-}_s(aq)$; $K_{sp1} = 4 \times 10^{-12}$
$2s$
$K_{sp_1} = [A^+]^2[X^{2-}]$
$4 \times 10^{-12} = (2S)^2(S)$
$4 \times 10^{-12} = 4S^3$

$\Rightarrow \ S = 10^{-3}M$

Let the solubility of MY be ' S_1 '.

$MY(s) \rightleftharpoons \underset{s_1}{M^+(aq)} + \underset{s_1}{Y^-(aq)}, K_{sp_2} = 4 \times 10^{-12}$

$K_{sp_2} = [M^+][Y^-]$

$K_{sp_2} = (S_1)^2$

$4 \times 10^{-12} = (S_1)^2$

$\Rightarrow \ S_1 = 2 \times 10^{-6}M$

$\frac{[A_2Y]}{[MY]} = \frac{S}{S_1} = \frac{10^{-4}}{2 \times 10^{-6}} = 50$

Question101

0.01 moles of a weak acid HA($K_a = 2.0 \times 10^{-6}$) is dissolved in 1.0L of 0.1M HCl solution. The degree of dissociation of HA is ..... $\times 10^{-5}$ (Round off to the nearest integer). [Neglect volume change on adding HA. Assume degree of dissociation $\ll < 1$]
[17 Mar 2021 Shift 1]

Answer: 2

Solution:

Solution:

Given, $[HA] = 0.01$

$[HCl] = 0.1M$

When strong acid (HCl) is completely dissociated,

$\underset{0.1}{HCl} \longrightarrow \underset{0.1}{H^+(aq)} + \underset{0.1}{Cl^-(aq)}$

For weak acid, dissociation is very less,

	HA	$\rightleftharpoons$	$H^+(aq)$	+	$A^-(aq)$
$t = 0$	0.01		0.1		0
$t = t_{eq}$	$0.01 - 0.01 - \alpha$		$0.1 + 0.01\alpha$		0.01α

$K_a = \frac{[H^+][A^-]}{[HA]} = 2 \times 10^{-6} \Rightarrow \frac{(0.1 + 0.01\alpha)(0.01\alpha)}{(0.01 - 0.01\alpha)} = 2 \times 10^{-6}$

As $0.01\alpha << 0.1 \quad [H^+] = 0.1$

and $0.01\alpha << 0.01 \quad [HA] = 0.01$

$\therefore 2 \times 10^{-6} = \frac{(0.1)(0.01\alpha)}{0.01} \Rightarrow \alpha = 2 \times 10^{-5}$

$\Rightarrow \ x = 2$

Question102

Given below are two statements: One is labelled as Assertion A and the other labelled as Reason R.
Assertion A During the boiling of water having temporary hardness, $Mg(HCO_3)_2$ is

converted to $MgCO_3$.

Reason R The solubility product of $Mg(OH)_2$ is greater than that of $MgCO_3$.

In the light of the above statements, choose the most appropriate answer from the options given below
[18 Mar 2021 Shift 1]

Options:

A. Both A and R are true but R is not the correct explanation of A.

B. A is true but R is fals

C. Both A and R are true and R is the correct explanation of A.

D. A is false but R is true.

Answer: D

Solution:

Solution:
During boiling, soluble $Mg(HCO_3)_2$ is converted into insoluble
$Mg(OH)_2$ and $Ca(HCO_3)_2$ is converted into insoluble $CaCO_3$. This is because of high solubility product of $Mg(OH)_2$ as compared to $MgCO_3$ hence, $Mg(OH)_2$ is precipitated. These precipitates can be removed by filtration. Thus, filtrate will be obtained in soft water.
$Mg(HCO_3)_2 \longrightarrow Mg(OH)_2 \downarrow +2CO_2$
Temporary hardness
$Ca(HCO_3)_2 \longrightarrow CaCO_3 \downarrow +CO_2 + 2H_2O$
K_{sp} of $Mg(OH)_2 > K_{sp}$ of $MgCO_3$
and hence, $Mg(OH)_2$ precipitation first.

Question103

The gas phase reaction
$2A(g) \rightleftharpoons A_2(g)$

at 400K has $\Delta G^\circ = +25.2 kJ\ mol^{-1}$. The equilibrium constant K_C for this reaction is ..... $\times 10^{-2}$. (Round off to the nearest integer).
[Use : $R = 8.3 J\ mol^{-1} K^{-1}$, ln 10 = 2.3] $\log_{10} 2 = 0.30$, 1atm = 1bar | [antilog (−0.3) = 0.501]
[18 Mar 2021 Shift 2]

Answer: 1.66

Solution:

Solution:

Given, $\Delta G = 25.2 kJ\, mol^{-1}$
$= 25200 J\, mol^{-1}$
$T = 400 K$
According to standard free Gibb's equation,
$\Delta G^0 = -RT\, \ln K_p$
$25200 = -2.3 \times 8.3 \times 400 \log(K_p)$
$\log K_p = \frac{-25200}{2.3 \times 8.3 \times 400} = -3.3$
$K_p = 10^{-3.3} = 10^{-3} \times 0.501$
$K_p = 5.01 \times 10^{-4} bar^{-1}$
$K_p = 5.01 \times 10^{-5} Pa^{-1}$
We know that,
$K_p = K_c(RT)^{\Delta r_k}$
$K_p = K_c(RT)^{-1} \quad [\therefore \Delta n_g = 1 - 2 = -1]$
$K_p = \frac{K_C}{8.3 \times 400}$
$K_c = 5.01 \times 10^{-5} \times 8.3 \times 400$
$\Rightarrow K_c = 166 \times 10^{-5} m^3/mol$
$= 1.66 \times 10^{-2} L/mol$

Question104

Consider the reaction, $N_2O_4(g) \rightleftharpoons 2NO_2(g)$. The temperature at which $K_C = 20.4$ and $K_p = 600.1$, is ..... K. (Round off to the nearest integer). [Assume all gases are ideal and R = 0.0831L bar, $K^{-1}mol^{-1}$].
[17 Mar 2021 Shift 2]

Answer: 354

Solution:

Solution:
The temperature at which $K_C = 20.4$ and $K_\rho = 600.1$, is $354K$.
Given reaction is, $N_2O_4(g) \rightleftharpoons 2NO_2(g)$
Given values are : $K_\rho = 600.1$
$K_C = 20.4$
$\Delta n_g =$ number of moles of product - number of moles of reactant Using relation between K_p and $K_C = 2 - 1 = 1$
where R is the gas constant $= 0.083 Latm/K\, mol$
$\Delta n_g = 1$ (for given reaction)
On putting given values, we will get
$600.1 = 20.4(RT)^1 \Rightarrow T \approx 354K$

Question105

For the reaction, $A(g) \rightleftharpoons B(g)$ at $495K$, Δ, $G^\circ = -9.478 kJ\, mol^{-1}$. If we start the reaction in a closed container at 495K with 22 millimoles of A, the amount of B is the

equilibrium mixture is ..... millimoles (Round off to the nearest integer).
[R = 8.314J mol $^{-1}$K $^{-1}$, ln 10 = 2.303]
[16 Mar 2021 Shift 1]

Answer: 20

Solution:

Solution:
$A(g) \rightleftharpoons B(g)$
Given, $T = 495K$, $\Delta_r C^\circ = -9.478kJ/mol$
We know,
$\Delta G^\circ = -2.303RT \log K$
$\because K = \frac{[B]}{[A]}$
where, K = equilibrium constant
Now, $\log K = \frac{-\Delta G^\circ}{2.303RT}$
$\log K = \frac{9.478 \times 1000}{2.303 \times 8.314 \times 495}$
$\therefore \log K = 1$
$\Rightarrow K = 10$
$\Rightarrow \frac{[B]}{[A]} = \frac{n_B}{n_A} = 10$

	$A(g)$	$\rightleftharpoons$	$B(g)$
Now, $t = 0$	22		0
$t = t$	$22 - x$		x

$K = \frac{[B]}{[A]} = \frac{x}{22 - x} = 10$
So, $x = 20$ Milimoles of $B = 20$

Question106

In order to prepare a buffer solution of pH 5.74, sodium acetate is added to acetic acid. If the concentration of acetic acid in the buffer is 1.0M , the concentration of sodium acetate in the buffer is ..... M. (Round off to the nearest integer). [Given : pK $_a$ (acetic acid =4.74]
[18 Mar 2021 Shift 1]

Answer: 10

Solution:

Solution:
Given : $pH = 5.74$
Concentration of acetic acid in buffer $= 1.0M$
Acetic acid and its conjugate base sodium acetate makes acidic buffer.

$CH_3COOH + CH_3COONa \longrightarrow$ (Acidic buffer)

Using formula,

$$pH = pK_a + \log \frac{[\text{Salt}]}{[\text{Acid}]}$$

$$pH = pK + agg \frac{[CH_3COONa]}{[CH_3COOH]}$$

$$5.74 = 4.74 + \log \frac{[CH_3COONa]}{[CH_3COOH]}$$

$$5.74 - 4.74 = \log \frac{[CH_3COONa]}{[CH_3COOH]}$$

$$1 = \log \frac{[CH_3COONa]}{[CH_3COOH]}$$

$$\Rightarrow \quad \frac{[CH_3COONa]}{[CH_3COOH]} = 10 \quad [\because [CH_3COOH] = 1]$$

$[CH_3COONa] = [10 \times 1] = 10$

Thus, the concentration of sodium acetate in buffer is $10M$.

Question107

Sulphurous acid (H_2SO_3) has $K_{a_1} = 1.7 \times 10^{-2}$ and $K_{a_2} = 6.4 \times 10^{-8}$. The pH of 0.588M is (Round off to the nearest integer)
[16 Mar 2021 Shift 2]

Answer: 1

Solution:

Solution:

For H_2SO_3, $K_{a_1} >> K_{a_2}$

So, we mainly consider 1st ionisation

$$H_2SO_3 \rightleftharpoons H^+ + HSO_3^-; \; K_{a_1} = 1.7 \times 10^{-2}$$

$t = 0$	0.558	0	0
$t = t_{eq}$	$0.558 - 0.558\alpha$	0.558α	0.558α

$$K_{a_1} = \frac{[H^+][HSO_3^-]}{[H_2SO_3]} = \frac{(0.558\alpha)(0.558\alpha)}{0.558(1-\alpha)} = \frac{0.558\alpha^2}{1-\alpha}$$

$\alpha << 1$ for weak acid

$(1-\alpha) \approx 1$

$$\alpha = \sqrt{\frac{K_{a_1}}{0.558}} = \sqrt{\frac{1.7 \times 10^{-2}}{0.558}}$$

$\alpha = 1.7 \times 10^{-1} = 0.17$

$[H]^+ = 0.558\alpha = 9.9 \times 10^{-2}$

$pH = -\log[H^+] = -\log(9.9 \times 10^{-2}) \quad [\because \log 9.9 \approx 1]$

$= 2 - \log 9.9 = 2 - 1$

$pH = 1$

Question108

Assuming that $Ba(OH)_2$ is completely ionised in aqueous solution under the given conditions the concentration of H_3O^+ ions in 0.005M aqueous solution of $Ba(OH)_2$ at 298K is _____ $\times 10^{-12}$ mol L^{-1}. (Nearest integer)
[25 Jul 2021 Shift 2]

Answer: 1

Solution:

Solution:
$Ba(OH)_2 \rightarrow Ba^{+2} + 2OH^-$
$\downarrow$
$2 \times 0.005 = 0.01 = 10^{-2}$
At 298K : in aq. solution $[H_3O^+][OH^-] = 10^{-14}$
$[H_3O^+] = \frac{10^{-14}}{10^{-2}} = 10^{-12}$

Question109

A solution is 0.1M in Cl^- and 0.001M in CrO_4^{2-}.
Solid $AgNO_3$ is gradually added to it Assuming that the addition does not change in volume and $K_{sp}(AgCl) = 1.7 \times 10^{-10} M^2$ and $K_{sp}(Ag_2CrO_4) = 1.9 \times 10^{-12} M^3$
Select correct statement from the following:
[20 Jul 2021 Shift 2]

Options:

A. AgCl precipitates first because its K_{sp} is high.

B. Ag_2CrO_4 precipitates first as its K_{sp} is low.

C. Ag_2CrO_4 precipitates first because the amount of Ag^+ needed is low.

D. AgCl will precipitate first as the amount of Ag^+ needed to precipitate is low.

Answer: D

Solution:

Solution:
(i) $[Ag^+]$ required to ppt $AgCl(s)$
$K_{sp} = IP = [Ag^+][Cl^-] = 1.7 \times 10^{-10}$
$[Ag^+] = 1.7 \times 10^{-9}$
(ii) $[Ag^+]$ required to ppt $Ag_2CrO_4(s)$
$K_{sp} = IP = [Ag+]^2[CrO_4^{-2}] = 1.9 \times 10^{-12}$

$[Ag^+] = 4.3 \times 10^{-5}$
$[Ag^+]$required to ppt $AgCl$ is low so $AgCl$ will ppt 1st

Question110

The equilibrium constant for the reaction
$A(s) \rightleftharpoons M(s) + \frac{1}{2}O_2(g)$
is $K_p = 4$. At equilibrium, the partial pressure of O_2 is _____ atm. (Round off to the nearest integer)
[27 Jul 2021 Shift 2]

Answer: 16

Solution:

Solution:
$k_p = Po_2^{1/2} = 4$
$\therefore Po_2 = 16bar = 16atm$

Question111

$2SO_2(g) + O_2(g) \rightarrow 2SO_3(g)$
The above reaction is carried out in a vessel starting with partial pressure $P_{SO_2} = 250m$ bar$P_{O_2} = 750m$ bar and $P_{SO_3} = 0$ bar . When the reaction is complete, the total pressure in the reaction vessel is _____ m bar. (Round off of the nearest integer).
[27 Jul 2021 Shift 2]

Answer: 875

Solution:

Solution:

	$2SO_2(g)$	+	$O_2(g)$	→	$2SO_3(g)$
Initial	250mbar (L . R.)		750mbar		O
Final	–250mbar		–125mbar		250mbar
	0		625mbar		250mba

∴ Final total pressure $= 625 + 250 = 875mbar$

Question112

$PCl_5 \rightleftharpoons PCl_3 + Cl_3$ $K_c = 1.844$

3.0 moles of PCl_5 is introduced in a 1L closed reaction vessel at 380K . The number of moles ofPCl_5 at equilibrium is _____ $\times 10^{-3}$

(Round off to the Nearest Integer)

[27 Jul 2021 Shift 1]

Answer: 1400

Solution:

Solution:

$PCl_{5(g)} \rightleftharpoons PCl_{3(g)} + Cl_{2(g)}$ $K_c = 1.844$

$\Rightarrow \frac{[PCl_3][Cl_2]}{[PCl_5]} = \frac{x^2}{3-x} = 1.844$

$\Rightarrow x^2 + 1.844 - 5.532 = 0$

$\Rightarrow x = \frac{-1.844 + \sqrt{(1.844)^2 + 4 \times 5.532}}{2}$

$\cong 1.604$

$\Rightarrow$ Moles of $PCl_5 = 3 - 1.604 \cong 1.396$

Question113

Value of K_p for the equilibrium reaction

$N_2O_4 \rightleftharpoons 2NO_{2(g)}$ at 288K is 47.9. The K_C for this reaction at same temperature is _____. (Nearest integer)

(R = 0.083L. bar $K^{-1}mol^{-1}$)

[22 Jul 2021 Shift 2]

Answer: 2

Solution:

Solution:

$K_C = \frac{K_P}{RT} = \frac{47.9}{0.083 \times 288} = 2$

Question114

$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$

In an equilibrium mixture, the partial pressures are $P_{SO_3} = 43kPa$; $P_{O_2} = 530Pa$ and $P_{SO_2} = 45kPa$. The equilibrium constant K_P = _____ $\times 10^{-2}$. (Nearest integer)

[20 Jul 2021 Shift 1]

Answer: 172

Solution:

Solution:

$2SO_2(g) + O_2(g) = 2SO_3(g)$

$$K_P = \frac{(pSO_{3(g)})^2}{pSO2(g)} \times pO_{2(g)}$$

$$= \frac{43 \times 43}{45 \times 45} \times 530Pa^{-1}$$

$$= 172.28 \times 10^{-5}Pa^{-1}$$

$$= 172.28atm$$

$$= 17228 \times 10^{-2}atm$$

Question115

A_3B_2 is a sparingly soluble salt of molar mass $M(gmol^{-1})$ and solubility xgL^{-1}. The solubility product satisfies $K_{sp} = a\left(\frac{x}{M}\right)^5$.

The value of a is ..... (Integer answer)

[31 Aug 2021 Shift 1]

Answer: 108

Solution:

Solution:

$$A_3B_2 \rightleftharpoons \underset{3S}{3A^{2+}(aq)} + \underset{2S}{2B^{3-}(aq)}$$

$$K_{sp} = [A^{2+}]^3[B^{3-}]^2$$

$$K_{sp} = (3S)^3(2S)^2 = 108S^5$$

$$AlsOS = \frac{x}{m}$$

$$K_{sp} = 108\left(\frac{x}{m}\right)^5$$

Given that, $K_{sp} = a\left(\frac{x}{m}\right)^5$

$\therefore$ a = 108

Question116

The pH of a solution obtained by mixing 50 mL of 1M HCl and 30 mL of 1M NaOH is $x \times 10^{-4}$. The value of x is ..... (Nearest integer)
[log 2.5 = 0.3979]
[31 Aug 2021 Shift 2]

Answer: 6021

Solution:

Solution:
Milliequivalents of $HCl(N_aV_a) = 50 \times 1 = 50$

Milliequivalents of $NaOH(N_bV_b) = 30 \times 1 = 30$

Since, $N_aV_a > N_bN_b$

and they neutralise each other

$$[H^+] = \frac{N_aV_a - N_bV_b}{V_a + V_b}$$

$$= \frac{50-30}{80} = 0.25 = 2.5 \times 10^{-1}$$

$$pH = -\log[H^+] = -\log(2.5 \times 10^{-1})$$

$$= 1 - 0.3979 = 0.6021$$

$$pH \times 10^4 = 0.6021 \times 10^4 = 6021$$

$\therefore$ x = 6021

Question117

The number of moles of NH_3, that must be added to 2L of 0.80M$AgNO_3$ in order to reduce the concentration of Ag^+ ions to 5.0×10^{-8}M ($K_{formation}$ for $[Ag(NH_3)_2]^+ = 1.0 \times 10^8$) is..... (Nearestinteger)
[Assume no volume change on adding NH_3]
[27 Aug 2021 Shift 1]

Answer: 4

Solution:

Solution:

Let moles added = a

$Ag^{+} + 2NH_3 \rightleftharpoons Ag(NH_3)_2^{+}$

t = 0

$0.8 \left(\frac{a}{2}\right)$

t = ∞

$5 \times 10^{-8}\left[\frac{a}{2} - 1.6\right]$ 0.8

$\frac{0.8}{5 \times 10^{-8}\left(\frac{a}{2} - 1.6\right)^2} = 10^8$

$\frac{a}{2} - 1.6 = 0.4$

⇒ a = 4.

Question118

The number of moles of NH_3, that must be added to 2L of 0.80M$AgNO_3$ in order to reduce the concentration of Ag^+ ions to 5.0×10^{-8}M ($K_{formation}$ for $[Ag(NH_3)_2]^+ = 1.0 \times 10^8$) is..... (Nearestinteger)
[Assume no volume change on adding NH_3]
[27 Aug 2021 Shift 1]

Answer: 4

Solution:

Solution:

Let moles added = a

$Ag^{+} + 2NH_3 \rightleftharpoons Ag(NH_3)_2^{+}$

t = 0

$0.8 \left(\frac{a}{2}\right)$

t = ∞

$5 \times 10^{-8}\left[\frac{a}{2} - 1.6\right]$ 0.8

$\frac{0.8}{5 \times 10^{-8}\left(\frac{a}{2} - 1.6\right)^2} = 10^8$

$\frac{a}{2} - 1.6 = 0.4$

⇒ a = 4.

Question119

The equilibrium constant K_c at 298K for the reaction A + B ⇌ C + D is 100 . Starting with an equimolar solution with concentrations of A, B, C and D all equal to 1M, the

equilibrium concentration of D is .....$\times 10^{-2}$M. (Nearestinteger)
[26 Aug 2021 Shift 2]

Answer: 182

Solution:

Solution:
$A + B \rightleftharpoons C + D$
Initially,
At equilibrium, 1− x, 1− x, 1+ x, 1+ x

$\therefore K_C = \left(\frac{1+x}{1-x}\right)^2$

$100 = \left(\frac{1+x}{1-x}\right)^2$

$\frac{1+x}{1-x} = 10$

$x = \frac{9}{11}$

Moles of $D = 1 + x$

$= 1 + \frac{9}{11} = \frac{20}{11}$

$= 1.818 = 181.8 \times 10^{-2} = 181.8 \times 10^{-2}$

$\cong 182 \times 10^{-2}M$

Question120

When 5.1g of solid NH_4HS is introduced into a two litre evacuated flask at 27°C, 20% of the solid decomposes into gaseous ammonia and hydrogen sulphide. The K_p for the reaction at 27°C is $x \times 10^{-2}$. The value of x is ..... (Integer answer)
[Given, R = 0.082L atm K^{-1} mol^{-1}]
[27 Aug 2021 Shift 2]

Answer: 6

Solution:

Solution:
51g of $NH_4HS = 1$ mol

5.1g of $NH_4HS = \frac{1}{51} \times 5.1 = 0.1$ mol

$NH_4HS(s) \rightleftharpoons NH_3(g) + H_2S(g)$

At t = 0, 0.1

At t = t, $0.1(1-\alpha)$ 0.1α 0.1α

It is given that dissociation is 20% from 100 moles.
∴ 20 moles get dissociated.

20% dissociation from $1\ \text{mol}\ \frac{20}{100} = 0.2$ moles get dissociated.

α = 0.2

$\because K_C = \frac{[NH_3][H_2S]}{[NH_4HS]} = \frac{[NH_3][H_2S]}{1}$ (Concentration of solid is assumed as 1)

$[NH_3] = \frac{\text{Number of moles of } NH_3(g)}{\text{Volume(in L)}} = \frac{0.1\alpha}{2}$

$[H_2S] = \frac{\text{Number of moles of } H_2S}{\text{Volume(in L)}} = \frac{0.1\alpha}{2}$

$K_C = 0.1\alpha 2 \times \frac{0.1\alpha}{2} = \frac{0.1 \times 0.2}{2} \times \frac{0.1 \times 0.2}{2} = 10^{-4}$

$\therefore K_p = K_C(RT)^{\Delta n}$

[Δn =change in the number of gaseous moles = 2]

$K_p = 10^{-4} \times (0.08 \times 300)^2$

$K_p = 10^{-4} \times (24)^2 = 0.06$

$x \times 10^{-2} = 0.06$

$\Rightarrow x = \frac{0.06}{10^{-2}} = 0.06 \times 10^2 = 6$

x = 6 is the answer.

Question121

The molar solubility of $Zn(OH)_2$ in 0.1M NaOH solution is $x \times 10^{-18}$M. The value of x is ..... (Nearest integer)
(Given; The solubility product of $Zn(OH)_2$ is 2×10^{-20}).
[1 Sep 2021 Shift 2]

Answer: 2

Solution:

Solution:

$Zn(OH)_2 \rightleftharpoons \underset{S}{Zn^{2+}} + \underset{2S}{2OH^-}$

Due to common-ion effect (presence of NaOH) the concentration of OH^- will be (2S + 0.1) ≈ 0.1

(∵ 0.1> > 2 S)

∴ Solubility of product,

$K_{sp} = (0.1)^2 \times S$

$2 \times 10^{-20} = 0.01 \times S$


$\Rightarrow S = \frac{2 \times 10^{-20}}{0.01} = 2 \times 10^{-18}$

∴ x = 2

Hence, answer is 2.

Question122

In the figure shown below reactant A (represented by square) is in equilibrium with product B (represented by circle). The equilibrium constant is:



[Jan. 09, 2020 (II)]

Options:

A. 4

B. 8

C. 1

D. 2

Answer: D

Solution:

Solution:
Equilibrium constant
$K_c = \frac{[B]}{[A]} = \frac{11}{6} \approx 2$

Question123

For the following Assertion and Reason, the correct option is:
Assertion: The pH of water increases with increase in temperature.
Reason: The dissociation of water into H^+ and OH^- is an exothermic reaction.
[Jan.08,2020(II)]

Options:

A. Both assertion and reason are true, and the reason is the correct explanation for the assertion.

B. Both assertion and reason are false.

C. Both assertion and reason are true, but the reason is not the correct explanation for the assertion.

D. Assertion is not true, but reason is true.

Answer: B

Solution:

Solution:
Temperature plays a significant role on pH measurements. As the temperature rises, molecular vibrations increase which results in greater ability of water to ionise and form more hydrogen ions.
As a result, the pH will drop. So assertion is incorrect. The dissociation of water molecules into ions is bond breaking and is therefore an endothermic process (energy must be absorbed to break the bonds). So reason is also incorrect.

Question124

Two solutions, A and B, each of 100L was made by dissolving 4g of N aOH and 9.8g of H_2SO_4 in water, respectively. The pH of the resultant solution obtained from mixing 40L of solution A and 10L of solution. B is _____.
[NV, Jan. 07, 2020 (I)]

Answer: 10.60

Solution:

Solution:

$$M_{H_2SO_4} = \frac{9.8}{98 \times 100} = 10^{-3}M$$

$$M_{NaOH} = \frac{4}{40 \times 100} = 10^{-3}M$$

After neutralisation $[OH^-]$ can be calculated as

$$[OH] = \frac{(40 \times 10^{-3}) - (2 \times 10^{-3} \times 10)}{50}$$

$$= \frac{20}{50} \times 10^{-3}$$

$$[OH^-] = \frac{2}{5} \times 10^{-3}$$

$$pOH = 3.397$$

$$pH = 14 - pOH$$

$$= 14 - 3.397 = 10.603$$

Question125

3g of acetic acid is added to 250mL of 0.1M H Cl and the solution made up to 500mL. To 20mL of this solution $\frac{1}{2}$ mL of 5M N aOH is added. The pH of the solution is _____.
[Given: pKa of acetic acid = 4.75, molar mass of acetic acid = 60g/mol, log 3 = 0.4771]
Neglect any changes in volume.
[NV, Jan. 07, 2020 (II)]

Answer: 5.22

Solution:

Solution:

$$\text{No. of moles} = \frac{\text{Mass}}{\text{Molar mass}}$$

$$3g\,CH_3COOH = \frac{3}{60}0.5mol = 50mmol$$

No. of millimoles = Molarity × Volume in mL

250mL of 0.1M HCl $= 250 \times 0.1 = 25\text{mmol}$
500mL solution $= 50\text{mmol } CH_3COOH$

20mL solution $= \frac{50}{500} \times 20 = 2\text{mmol } CH_3COOH$

500mL solution contains $= 25\text{mmol HCl}$

20mL solution contains $= \frac{25}{500} \times 20 = 1\text{mmol HCl}$

$\frac{1}{2}$mL of 5M NaOH $= \frac{1}{2} \times 5 = 2.5\text{mmol NaOH}$

$\underset{1}{HCl} + \underset{2.5}{NaOH} \longrightarrow \underset{1}{NaCl} + H_2O$

Remaining NaOH $= 2.5 - 1 = 1.5\text{mmol}$

CH_3COOH	+ NaOH (remaining)	$\rightarrow CH_3COONa$	+ Water
2	1.5	0	0
0.5	0	1.5	–

$pH = pK_a + \log \frac{1.5}{0.5} = 4.74 + \log 3$

$= 4.74 + 0.48 = 5.22$

Question126

The K_{sp} for the following dissociation is 1.6×10^{-5} $PbCl_2(s) \rightleftharpoons Pb^{2+}(aq) + 2Cl^-(aq)$ Which of the following choices is correct for a mixture of 300 mL 0.134 M $Pb(NO_3)_2$ and 100 mL 0.4 MNaCl?

[Jan. 09, 2020 (I)]

Options:

A. Not enough data provided

B. $Q < K_s$

C. $Q > K_{sp}$

D. $Q = K_{sp}$

Answer: C

Solution:

Solution:

$PbCl_2 \rightleftharpoons Pb^{2+}(aq) + 2Cl^-(aq)$

Given; $K_{sp} = 1.6 \times 10^{-5}$

$[Pb^{2+}] = \frac{300 \times 0.134}{400} = 0.1005$

$[Cl^-] = \frac{100 \times 0.4}{400} = 0.1$

$Q = [Pb^{2+}][Cl^-]^2 = 0.1005 \times (0.1)^2$

$= 1.005 \times 10^{-3}$

$Q > K_{sp}$

Question127

The solubility product of $Cr(OH)_3$ at 298K is 6.0×10^{-31}.
The concentration of hydroxide ions in a saturated solution of $Cr(OH)_3$ will be:
[Jan. 09,2020 (II)]

Options:

A. $(2.22 \times 10^{-31})^{1/4}$

B. $(18 \times 10^{-31})^{1/4}$

C. $(18 \times 10^{-31})^{1/2}$

D. $(4.86 \times 10^{-29})^{1/4}$


Answer: B

Solution:

Solution:
$Cr(OH)_3 \rightarrow \underset{s^{3+}}{Cr} + 3OH^-$
$K_{sp} = s \cdot (3s)^3$
$\Rightarrow 6 \times 10^{-31} = 27 \cdot s^4; s = \left(\frac{6}{27} \times 10^{-31}\right)^{1/4}$
$[OH^-] = 3s = 3 \times \left(\frac{6}{27} \times 10^{-31}\right)^{1/4}$
$= (18 \times 10^{-31})^{1/4} M$

Question128

The stoichiometry and solubility product of a salt with the solubility curve given below is, respectively:



[Jan. 08,2020 (I)]

Options:

A. X_2Y, $2 \times 10^{-9} M^3$

B. XY_2, $4 \times 10^{-9} M^3$

C. XY_2, $1 \times 10^{-9} M^3$

D. XY, $2 \times 10^{-6} M^3$

Answer: B

Solution:

Solution:
From the given curve,
if $[X] = 1mM$ then $[Y] = 2mM$
$\therefore$ Salt is XY_2
$K_{sp} = [X][Y]^2 = (10^{-3})(2 \times 10^{-3})^2 = 4 \times 10^{-9} M^3$

Question129

For the reaction
$Fe_2N(s) + \frac{3}{2}H_2(g) \rightleftharpoons 2Fe(s) + NH_3(g)$
[Sep. 06,2020(I)]

Options:

A. $K_c = K_p(RT)$

B. $K_c = K_p(RT)^{\frac{-1}{2}}$

C. $K_c = K_p(RT)^{\frac{1}{2}}$

D. $K_c = K_p(RT)^{\frac{3}{2}}$

Answer: C

Solution:

Solution:
$K_p = K_c(RT)^{\Delta n_g} = K_c(RT)^{1-3/2} = K_c(RT)^{-1/2}$
$\Rightarrow K_c = K_p(RT)^{1/2}$

Question130

The value of K c is 64 at 800K for the reaction $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$.
The value of K_c for the following reaction is:
$NH_3(g) \rightleftharpoons \frac{1}{2}N_2(g) + \frac{3}{2}H_2(g)$
[Sep. 06, 2020 (II)]

Options:

A. 1/64

B. 8

C. 1/4

D. 1/8

Answer: D

Solution:

Solution:

$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$; K_c

$2NH_3(g) \rightleftharpoons N_2(g) + 3H_2(g)$; $\frac{1}{K_c}$

For

$NH_3(g) \rightleftharpoons \frac{1}{2}N_2(g) + \frac{3}{2}H_2(g)$; $\frac{1}{K_c^{1/2}}$

$\frac{1}{K_c^{1/2}} = \frac{1}{(64)^{1/2}} = \frac{1}{8}$

Question131

For a reaction $X + Y \rightleftharpoons 2Z$, 1.0mol of X, 1.5mol of Y and 0.5 mol of Z were taken in a 1 L vessel and allowed to react. At equilibrium, the concentration of Z was 1.0mol L^{-1}.
The equilibrium constant of the reaction is $\frac{x}{15}$. The value of x is _____.
[NV, Sep. 05, 2020 (II)]

Answer: 16

Solution:

Solution:


Question132

For the equilibrium A ⇌ B, the variation of the rate of the forward (a) and reverse (b) reaction with time is given by:
[Sep. 04,2020(I)]


**For the equilibrium A ⇌ B, the variation of the rate of the forward (a) and reverse (b) reaction with time is given by:
[Sep. 04,2020(I)]**

Options:


A.




B.



C.



D.



Answer: B

Solution:

Solution:
At equilibrium, rate of forward reaction = Rate of backward reaction.

Question133

If the equilibrium constant for $A \rightleftharpoons B + C$ is $K_{eq}^{(1)}$ and that of $B + C \rightleftharpoons P$ is $K_{eq}^{(2)}$, the equilibrium constant for $A \rightleftharpoons P$ is:
[Sep. 04,2020(II)]

Options:

A. $K_{eq}^{(1)} / K_{eq}^{(2)}$

B. $K_{eq}^{(2)} - K_{eq}^{(t)}$

C. $K_{eq}^{(1)} + K_{eq}^{(2)}$

D. $K_{eq}^{(1)} K_{eq}^{(2)}$

Answer: D

Solution:

Solution:
$A \rightleftharpoons B + C; K_{eq}^{(1)}$ . . . (i)
$B + C \rightleftharpoons P; K_{eq}^{(2)}$ . . . (ii)
On adding equations (i) and (ii), we get
$A \rightleftharpoons P$
$K_{eq}(\text{ overall }) = K_{eq}^{(1)} \cdot K_{eq}^{(2)}$

Question134

The variation of equilibrium constant with temperature is given below:
Temperature Equilibrium Constant
$T_1 = 25^\circ C$ $K_1 = 10$
$T_2 = 100^\circ C$ $K_2 = 100$
The values of ΔH°, ΔG° at T_1 and ΔG° at T_2 (in kj mol^{-1}) respectively, are close to
[use $R = 8.314 J K^{-1} mol^{-1}$]
[Sep. 06, 2020 (I)]

Options:

A. 28.4, −7.14 and −5.71

B. 0.64, −7.14 and −5.71

C. 28.4, −5.71 and −14.29

D. 0.64, −5.71 and −14.29

Answer: C

Solution:

Solution:
$\Delta G^{\circ} = -RT \ln K$, $T_1 = 25^{\circ}C$, $K_1 = 10$

ΔG° at $T_1 = -8.314 \times 298 \times 2.303 \times \log 10 = -5.71 kJ/mol$

ΔG° at $T_2 = -8.314 \times 298 \times 373 \times 2.303 \times \log(100)$

$= -14.29 kJ/mol$

$\Delta G^{\circ} = \Delta H^{\circ} - T \Delta S^{\circ}$

$\Rightarrow -5.71 = \Delta H^{\circ} - 298(\Delta S^{\circ})$

$\Rightarrow -14.29 = \Delta H^{\circ} - 373(\Delta S^{\circ})$

$\Delta H^{\circ} = 28.4 kJ/mol$

Question135

Consider the following reaction:

$N_2O_4(g) \rightleftharpoons 2NO_2(g)$; $\Delta H^0 = +58kJ$

For each of the following cases ((i), (ii)), the direction in which the equilibrium shifts is:

(i) Temperature is decreases

(ii) Pressure is increased by adding N_2 at constant T .

[Sep .05,2020(I)]

Options:

A. (i) towards product, (ii) towards product

B. (i) towards reactant, (ii) towards product

C. (i) towards reactant, (ii) no change

D. (i) towards product, (ii) no change

Answer: C

Solution:

Solution:
(i) As reaction is endothermic (ΔH z = + ve) so on decrease in temperature equilibrium will shift towards reactant side.
(ii) On increase in pressure by adding inert gas (N_2) at same temperature, no shifting will take place. The equilibrium changes only if the added gas is a reactant or product involved in the reaction.

Question136

Arrange the following solutions in the decreasing order of pOH:

(A) 0.01M HCl

(B) 0.01M NaOH

(C) 0.01M CH_3COONa

(D) 0.01M NaCl

[Sep. 06, 2020 (I)]

Options:

A. (A) > (C) > (D) > (B)

B. (A) > (D) > (C) > (B)

C. (B) > (C) > (D) > (A)

D. (B) > (D) > (C) > (A)

Answer: B

Solution:

Solution:
0.01M HCl
$[H^+] = 10^{-2}$, $pH = -\log 10^{-2} = 2$
$pOH = 14 - 2 = 12$
(B) 0.01M NaOH
$[OH^-] = 10^{-2}$, $pOH = -\log[OH] = 2$
(C) 0.01M CH_3COONa

$$pH = 7 + \frac{1}{2}[pK_a + \log 0.01]$$

$pH > 7 \Rightarrow pOH < 7$
(D) 0.01M NaCl, $pH = 7$, $pOH = 7$
Decreasing order of pOH value is,
(A) > (D) > (C) > (B)

Question137

A soft drink was bottled with a partial pressure of CO_2 of 3 bar over the liquid at room temperature. The partial pressure of CO_2 over the solution approaches a value of 30 bar when 44g of CO_2 is dissolved in 1kg of water at room temperature. The approximate pH of the soft drink is _____ $\times 10^{-1}$.
(First dissociation constant of $H_2CO_3 = 4.0 \times 10^{-7}$; $\log 2 = 0.3$; density of the soft drink $= 1 gmL^{-1}$)
[NV, Sep. 05, 2020(I)]

Answer: 7

Solution:

Solution:

$$CO_2 + H_2O \longrightarrow H_2CO_3$$

30 bar ..... 1 mol/L

3 bar ..... 0.1 mol/L

$$H_2CO_3 \rightleftharpoons H^+ + HCO_3^-$$

$t=0$	0.1	0	0
at Equb.	$0.1(1-\alpha)$	0.1α	0.1α

$4.0 \times 10^{-7} = \frac{0.1\alpha^2}{1-\alpha}$

$\Rightarrow (1-\alpha) = 1$

$\alpha^2 = 4 \times 10^{-6} \Rightarrow \alpha = 2 \times 10^{-3}$

$=[H^+] = 2 \times 10^{-4}M$


$pH = 4 \times \log 2 = 3.7 = 37 \times 10^{-1}$

Question138


100mL of 0.1M H Cl is taken in a beaker and to it 100mL of 0.1M N aOH is added in steps of 2mL and the pH is continuously measured. Which of the following graphs correctly depicts the change in pH?
[Sep. 03, 2020 (II)]

Options:


A.




B.



C.



D.



Answer: C

Solution:

Solution:
At equivalence point pH is 7 and pH increases with addition of $NaOH$ so correct graph is (c).

Question139

If the solubility product of AB_2 is $3.20 \times 10^{-11} M^3$, then the solubility of AB_2 in pure water is _____$\times 10^{-4}$ mol L^{-1}.
[Assuming that neither kind of ion reacts with water]
[NV, Sep.06, 2020 (II)]

Answer: 2

Solution:

Solution:
$AB_2 \rightleftharpoons \underset{s}{A^{2+}}(aq) + \underset{2s}{2B^-}(aq)$
$K_{sp} = 4s^3 = 3.2 \times 10^{-11}$
$\Rightarrow s^3 = 8 \times 10^{-12}$
$\Rightarrow s = 2 \times 10^{-4}$

Question140

An acidic buffer is obtained on mixing :
[Sep. 03,2020 (I)]

Options:

A. 100mL of 0.1M CH_3COOH and 100mL of 0.1M NaOH

B. 100mL of 0.1M HCl and 200mL of 0.1M NaCl

C. 100mL of 0.1M CH_3COOH and 200mL of 0.1M NaOH

D. 100mL of 0.1M HCl and 200mL of 0.1M CH_3COONa

Answer: D

Solution:

Solution:

$$HCl + CH_3COONa \longrightarrow CH_3COOH + NaCl$$

Millimoles at start	10	20	0	0
Millimoles after reaction	0	10	10	10

Buffer solution contains CH_3COONa (10 millimole) and CH_3COOH (10 millimole) which is a acidic buffer.

Question141

For the following Assertion and Reason, the correct option is
Assertion (A): When Cu (II) and sulphide ions are mixed, they react together extremely quickly to give a solid.
Reason (R): The equilibrium constant of $Cu^{2+}(aq) + S^{2-}(aq) \rightleftharpoons CuS(s)$ is high because the solubility product is low.
[Sep. 02,2020(I)]

Options:

A. (A) is false and (R) is true

B. Both (A) and (R) are false

C. Both (A) and (R) are true but (R) is not the explanation for (A)

D. Both (A) and (R) are true and (R) is the explanation for (A)

Answer: C

Solution:

Solution:
Cu^{2+} ions get precipitated every quickely due to low K_{sp} value even at very low concentration of S^{2-} ion.
$CuS(s) \rightleftharpoons Cu^{2+} + S^{2-}$

$K_{sp} = [Cu^{2+}][S^{2-}]$

$Cu^{2+} + S^{2-} \rightleftharpoons CuS(s)$

$K_{eq} = \frac{1}{[Cu^{2+}][S^{2-}]} = \frac{1}{K_{sp}}$

Due to high value of K_{eq}, CuS precipitated easily.

Question142

For the equilibrium $2H_2O \rightleftharpoons H_3O^+ + OH^-$; the value of ΔG° at 298K is approximately: [Jan. 11, 2019 (II)]

Options:

A. $100 kJ\ mol^{-1}$

B. $-80 kJ\ mol^{-1}$

C. $80 kJ\ mol^{-1}$

D. $-100 kJ\ mol^{-1}$

Answer: C

Solution:

Solution:

$\Delta G = \Delta G^\circ + RT\ \ln Q$

At equilibrium; $\Delta G = 0$ and $Q = K_{eq}$

$\Rightarrow \Delta G^\circ = -2.303RT\ \log K_w$

$= -2.303 \times 8.314 \times 298 \times \log 10^{-14}$

$= 79.9 kJ/mol \approx 80 kJ/mol$

Question143

Which amongst the following is the strongest acid? [Jan. 9,2019 (I)]

Options:

A. $CHBr_3$

B. CHI_3


C. $CH(CN)_3$

D. $CHCl_3$

Answer: C

Solution:

Solution:
Due to the resonance stabilisation of the conjugate base, $CH(CN)_3$ is the strongest acid amongst the given compounds.
$CH(CN)_3 \rightleftharpoons H^+ + \overline{C}(CN)_3$



The conjugate bases of $CHBr_3$ and CHI_3 are stabilised by inductive effect of halogens. This is why, they are less stable. Also, the conjugate base of $CHCl_3$ involves backbonding between $2p$ and $3p$ orbitals.

Question144

If K_{50} of Ag_2CO_3 is 8×10^{-12}, the molar solubility of Ag_2CO_3 in 0.1M $AgNO_3$ is: [Jan. 12,2019(II)]

Options:

A. $8 \times 10^{-12}M$

B. $8 \times 10^{-11}M$

C. $8 \times 10^{-10}M$

D. $8 \times 10^{-13}M$

Answer: C

Solution:

Solution:
As $AgNO_3$ dissociates completely,
therefore in $0.1M\ AgNO_3$ solution, $[Ag^+] = 0.1M$
$AgNO_3 \longrightarrow \underset{0.1^+}{Ag} + NO_{0.1}^-$
$Ag_2CO_3 \rightleftharpoons {}_{0.1+2ss}Ag^+ + CO_3^{2-}$
$K_{sp} = [Ag^+]^2[CO_3^{2-}]$
$8 \times 10^{-12} = (0.1+2s)^2 \times s$
$0.01s = 8 \times 10^{-12};\ (0.1 + 2s \times 0.1)$
$s = 8 \times 10^{-10}M$

Question145

20mL of 0.1M H_2SO_4 solution is added to 30mL of 0.2 M NH_4OH solution. The pH of the resultant mixture is:
[pK_b of NH_4OH = 4.7]
[Jan. 9,2019 (I)]

Options:

A. 5.2

B. 9.0

C. 5.0

D. 9.4

Answer: B

Solution:

Solution:
m · mol of $H_2SO_4 = 20 \times 0.1 = 2$
m . mol of $NH_4OH = 30 \times 0.2 = 6$

$$H_2SO_4 + 2NH_4OH \rightarrow (NH_4)_2SO_4 + 2H_2O$$

Initial	2 m mol	6 m mol	0
Final	(2-2)	(6 – 2 × 2)	2 m mol

$[NH_4OH]_{left} = 2m\,mol$
$[(NH_4)_2SO_4] = 2m\,mol$
$[NH_4] = 2 \times 2 = 4m\,mol$
Total Volume $= 30 + 20 = 50\,mL$

$$pOH = pK_b + \log\left[\frac{Salt}{Base}\right]$$

$$= 4.7 + \log\frac{4/50}{2/50}$$

$$= 4.7 + \log 2 = 5$$

$pH = 14 - pOH$
$pH = 14 - 5 = 9$

Question146

A mixture of 100m mol of $Ca(OH)_2$ and 2g of sodium sulphate was dissolved in water and the volume was made up to 100mL. The mass of calcium sulphate formed and the concentration of OH^- in resulting solution, respectively, are: (Molar mass of $Ca(OH)_2$, Na_2SO_4 and $CaSO_4$ are 74,143 and 136gmol $^{-1}$, respectively; K_{sp} of $Ca(OH)_2$ is 5.5×10^{-6})
[Jan. 10,2019(I)]

Options:

A. 1.9g, 0.28mol L^{-1}

B. 13.6g, 0.28mol L^{-1}

C. 1.9g, 0.14mol L^{-1}

D. 13.6g, 0.14mol L^{-1}

Answer: A

Solution:

Solution:
$\text{N a}_2\text{SO}_4 + \text{Ca(OH)}_2 \rightarrow \text{CaSO}_4 + 2\text{N aOH}$
m mol of $\text{N a}_2\text{SO}_4 = \frac{2 \times 1000}{143} = 13.98 \text{m mol}$
m mol of CaSO_4 formed $= 13.98 \text{m mol}$
Mass of CaSO_4 formed $= 13.98 \times 10^{-3} \times 136 = 1.90 \text{g}$
m mol of N aOH $= 28 \text{m mol} \approx 0.028 \text{ mol}$
$\text{Ca(OH)}_2 \rightleftharpoons \underset{S}{\text{Ca}^{2+}} + \underset{2s+0.028}{2\text{OH}^-}$
Value of ' S ' will be negligible so
$[\text{OH}] = \frac{0.028 \text{ mol}}{0.1\text{L}} = 0.28 \text{ mol L}^{-1}$

Question147

Two solids dissociate as follows

$\text{A(s)} \rightarrow \text{B(g)} + \text{C(g)}; \text{K}_{\text{P}_1} = \text{xatm}^2$

$\text{D(s)} \rightarrow \text{C(g)} + \text{E (g)}; \text{K}_{\text{P}_2} = \text{yatm}^2$

The total pressure when both the solids dissociate simultaneously is: [Jan. 12, 2019 (I)]

Options:

A. $\sqrt{x+y}$ atm

B. $2(\sqrt{x+y})$ atm

C. $(x+y)$atm

D. $x^2 + y^2$ atm

Answer: B

Solution:

Solution:
$\text{A(s)} \rightarrow \text{B(g)} + \text{C(g)}$
$P_1 \quad P_1 + P_2$
$K_{P_1} = P_B \times P_C$
$P_1(P_1 + P_2) = x$
$K_{P_2} = P_C \times P_E$
$(P_1 + P_2)P_2 = y \quad ...(ii)$
Adding (i) and (ii)

$\therefore P_1(P_1+P_2)+P_2(P_1+P_2)=x+y$

$P_1^2+P_1P_2+P_2P_1+P_2^2=x+y$

$P_1^2+P_2^2+2P_1P_2=x+y$

$\Rightarrow \&(P_1+P_2)^2=x+y$

$\Rightarrow P_1+P_2=\sqrt{x+y}$

$\therefore$ Total pressure $(P_T)=P_C+P_B+P_E$

$(P_1+P_2)+P_1+P_2=2(P_1+P_2)$

$P_T=2(\sqrt{x+y})$

Question148

In a chemical reaction, $A+2B \overset{K}{\rightleftharpoons} 2C+D$, the initial concentration of B was 1.5 times of the concentration of A, but the equilibrium concentrations of A and B were found to be equal. The equilibrium constant (K) for the aforesaid chemical reaction is:
[Jan. 12, 2019 (I)]

Options:

A. 4

B. 16

C. $\frac{1}{4}$

D. 1

Answer: A

Solution:

Solution:

	A	+	2B	$\rightleftharpoons$	2C	+	D
$t=0$	2		3		0		0
t_{eq}	$2-x$		$3-2x$		$2x$		x

Given, $3-2x=2-x$

$\Rightarrow x=1$

$\therefore [C]=2, [D]=1, [A]=1, [B]=1$

$K_c=\left\{\frac{2^2\times 1}{1^2\times 1}\right\}=4$

Question149

Consider the reaction
$N_2(g)+3H_2(g) \rightleftharpoons 2NH_3(g)$
The equilibrium constant of the above reaction is K_p.If pure ammonia is left to dissociate, the partial pressure of ammonia at equilibrium is given by (Assume that

$P_{NH_3} << P_{total}$ at equilibrium)
[Jan. 11, 2019 (I)]

Options:

A. $\frac{3^{3/2}K_P^{1/2}P^2}{16}$

B. $\frac{K_P^{1/2}P^2}{16}$

C. $\frac{K_P^{1/2}P^2}{4}$

D. $\frac{3^{3/2}K_P^{1/2}P^2}{4}$

Answer: A

Solution:

Solution:

$2NH_3(g) \rightleftharpoons \underset{P_{NH_3}}{N_2(g)} + \underset{P_{N_2}}{3H_2(g)}, K = \frac{1}{K_P}$ $\quad\quad\quad

A. 1, $24.62 dm^3\ atm\ mol^{-1}$,
$606.0 dm^6 atm^2 mol^{-2}$

B. 1, $24.62 d\ m^3\ atm\ mol^{-1}$,
$1.65 \times 10^{-3} dm^{-6} atm^{-2} mol^2$

C. 1, $4.1 \times 10^{-2} dm^{-3} atm^{-1}$ mol,
$606 dm^6 atm^2 mol^2$

D. $24.62 dm^3\ atm\ mol^{-1}$,
$606.0 dm^6 atm^2 mol^{-2}$
$1.65 \times 10^{-3} dm^{-6} atm^{-2} mol^2$

Answer: B

Solution:

Solution:
$K_p = K_c (RT)^{\Delta n_g}$
$\Delta n_g =$ No. of gaseous moles of products - No. of gaseous moles of reactants $\frac{K_p}{K_c} = (RT)^{\Delta n_g}$
$N_2(g) + O_2(g) \rightleftharpoons 2NO(g), \Delta n_g = 0$
$\frac{K_p}{K_c} = (24.62 d\ m^3 atm mol^{-1})^0 = 1$
$N_2O_4(g) \rightleftharpoons 2NO_2(g), \Delta n_g = 1$
$\frac{K_p}{K_c} = 24.62 d\ m^3 atm mol^{-1}$
$N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g), \Delta n_g = -2$
$\frac{K_p}{K_c} = (24.62 d\ m^3 atm mol^{-1})^{-2}$
$= \frac{1}{(24.62 d\ m^3 atm mol^{-1})^2}$
$= 1.65 \times 10^{-3} d\ m^{-6} atm^{-2} mol^2$

Question151

5.1 g NH_4SH is introduced in 3.0L evacuated flask at 327°C, 30% of the solid NH_4SH decomposed to NH_3 and H_2S as gases. The K_p of the reaction at 327°C is (R = 0.082L atm $mol^{-1} K^{-1}$, molar mass of S = 32g mol^{-1}, molar mass of N = 14g mol^{-1}) [Jan. 10, 2019 (II)]

Options:

A. $0.242 \times 10^{-4} atm^2$

B. $1 \times 10^{-4} atm^2$

C. $4.9 \times 10^{-3} atm^2$

D. $0.242 atm^2$

Answer: D

Solution:

Solution:
Concerned reaction:
$NH_4SH \longrightarrow NH_3(g) + H_2S(g)$
$\text{Initial moles} = \frac{5.1}{51} = 0.1 mol$
Moles at equillibrium
$NH_4SH \longrightarrow NH_3(g) + H_2S(g)$
$0.1(1-0.3) \quad 0.1 \times 0.3 \quad 0.1 \times 0.3$
$\therefore K_c = [NH_3][H_2S] = \left(\frac{0.03}{3}\right)^2 = 10^{-4}$
$K_p = K_c(RT)^{\Delta n} g$
$= 10^{-4} \times (0.082 \times 600)^2 = 0.242 atm^2$

Question152

Consider the following reversible chemical reactions:

$A_2(g) + B_2(g) \overset{K_1}{\leftarrow} 2AB(g) \ldots\ldots (1)$

$6AB(g) \overset{K_2}{\rightleftharpoons} 3A_2(g) + 3B_2(g) \ldots\ldots (2)$

The relation between K_1 and K_2 is:

[Jan. 9,2019(II)]

Options:

A. $K_1 K_2 = \frac{1}{3}$

B. $K_2 = K_1^3$

C. $K_2 = K_1^{-3}$

D. $K_1 K_2 = 3$

Answer: C

Solution:

Solution:
Given:
$A_2 + B_2 \overset{K_1}{\rightleftharpoons} 2AB$
$\Rightarrow \quad 2AB \rightleftharpoons A_2 + B_2; K = \frac{1}{K_1}$
$6AB \overset{K_1}{\leftarrow} 3A_2 + 3B_2; K_2 = \left(\frac{1}{K_1}\right)^3$
The relation between K_1 and K_2 is $K_2 = K_1^{-3}$

Question153

The INCORRECT match in the following is: [April 12, 2019 (II)]

Options:

A. $\Delta G^0 < 0, K > 1$

B. $\Delta G^0 = 0, K = 1$

C. $\Delta G^0 > 0, K < 1$

D. $\Delta G^0 < 0, K < 1$

Answer: D

Solution:

Solution:
$\Delta G^\circ = -RT \ln K$
$\therefore$ If $K > 1$ then $\Delta G^\circ < 0$
If $K < 1$ then $\Delta G^\circ > 0$
If $K = 1$ then $\Delta G^\circ = 0$

Question154

In which one of the following equilibria, $K_p \neq K_c$? [April 12, 2019 (II)]

Options:

A. $2C(s) + O_2(g) \rightleftharpoons 2CO(g)$

B. $2HI(g) \rightleftharpoons H_2(g) + I_2(g)$

C. $NO_2(g) + SO_2(g) \rightleftharpoons NO(g) + SO_3(g)$

D. $2NO(g) \rightleftharpoons N_2(g) + O_2(g)$

Answer: A

Solution:

Solution:
We know that, $K_p = K_c . (RT)^{\Delta n_g}$
$\therefore$ If $\Delta n_g \neq 0$ then $K_p \neq K_c$
Now, $2C(s) + O_2(g) \rightleftharpoons 2CO(g)$
$\Delta n_g = +1$
$\Rightarrow K_p = K_c (RT)^1$
Hence, $K_p \neq K_c$

Question155

For the following reactions, equilibrium constants are given:

$S(s) + O_2(g) \rightleftharpoons SO_2(g)$; $K_1 = 10^{52}$

$2S(s) + 3O_2(g) \rightleftharpoons 2SO_3(g)$; $K_2 = 10^{129}$

The equilibrium constant for the reaction,

$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ is:

[April 8, 2019 (II)]

Options:

A. 10^{154}

B. 10^{181}

C. 10^{25}

D. 10^{77}

Answer: C

Solution:

Solution:
Given, $S + O_2 \rightleftharpoons SO_2$... (i); $K_1 = 10^{52}$
$2S + 3O_2 \rightleftharpoons 2SO_3$ ... (ii); $K_2 = 10^{129}$
$2SO_2 + O_2 \rightleftharpoons 2SO_3$ ... (iii) ; $K = ?$
To get equation (iii) follow (ii) −2 (i),
$2S + 3O_2 \rightarrow 2SO_3$ $K = 10^{129}$
$-(2S + 2O_2 \rightarrow 2SO_2$ $K = 10^{104})$
$O_2 \rightarrow 2SO_3 - 2SO_2$ $K = 10^{25}$
or $2SO_2 + O_2 \rightarrow 2SO_3 K = 10^{25}$

Question156

For the reaction,

$2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$

$\Delta H = -57.2 kJ\ mol^{-1}$ and $K_c = 1.7 \times 10^{16}$

Which of the following statement is INCORRECT?

[April 10, 2019 (II)]

Options:

A. The equilibrium constant is large suggestive of reaction going to completion and so no catalyst is required.

B. The equilibrium will shift in forward direction as the pressure increases.

C. The equilibrium constant decreases as the temperature increases.

D. The addition of inert gas at constant volume will not affect the equilibrium constant.

Answer: A

Solution:

Solution:
Equilibrium constant has no relation with catalyst. Catalyst only affects the rate of the reaction. Catalyst, V_2O_5 in the given reaction, is used to speed up the reaction.

Question157

Consider the following statements
(a) The pH of a mixture containing 400mL of 0.1M H_2SO_4 and 400mL of 0.1M N aOH will be approximately 1.3.
(b) Ionic product of water is temperature dependent.
(c) A monobasic acid with $K_a - 10^{-5}$ has a pH = 5. The degree of dissociation of this acid is 50%.
(d) The Le Chatelier's principle is not applicable to common-ion effect.
The correct statements are :
[April 10, 2019 (I)]

Options:

A. (a), (b), and (d)

B. (a), (b) and (c)

C. (b) and (c)

D. (a) and (b)

Answer: B

Solution:

Solution:

$$H_2SO_4 + NaOH \rightarrow NaHSO_4 + H_2O$$

Initial mol	0.04	0.04	0	0
mol at eqm.	0	0	0.04	0.04

$NaHSO_4 \rightarrow Na^+ + H^+ + SO_4^{2-}$

$[H^+] = \frac{0.04}{0.80} = 0.05M$; pH = 1.3

(b) Ionic product of water increases with increase in temperature because ionisation of water is endothermic.


(c) $HA \rightleftharpoons H^+ + A^-$

Initial	C	0	0
At eqm.	C (1 – α)	Cα	Cα

Given $pH = 5 \Rightarrow -\log(H^+) = 5$
$\therefore [H^+] = 10^{-5}$
As we know,
$K_a = \frac{C\alpha^2}{1-\alpha}$; $10^{-5} = \frac{C\alpha^2}{1-\alpha} = \frac{C\alpha \cdot \alpha}{(1-\alpha)}$
$10^{-5} = 10^{-5}\frac{\alpha}{1-\alpha}$; $\alpha = \frac{1}{2}$ i.e., 50%

Question158

In an acid base titration, 0.1M H Cl solution was added to the N aOH solution of unknown strength. Which of the following correctly shows the change of pH of the titration mixture in this experiment?



[April 9,2019 (II)]

Options:

A. (B)

B. (A)

C. (C)

D. (D)

Answer: B

Solution:

Solution:
Graph A and B, both represents the titration curve between strong acid and strong base, i.e., H Cl and N aOH but, the pH of N aOH is more than 7 and during the titration it decreases, so graph (A) is correct.

Question159

What is the molar solubility of $Al(OH)_3$ in 0.2M N aOH solution ?

Given that, solubility product of $Al(OH)_3 = 2.4 \times 10^{-24}$:

[April 12, 2019 (I)]

Options:

A. 3×10^{-19}

B. 12×10^{-21}

C. 3×10^{-22}

D. 12×10^{-23}

Answer: C

Solution:

Solution:
Let the solubility of $Al(OH)_3$ in $0.2M\ NaOH$ solution be s.
Then,
$Al(OH)_3 \rightleftharpoons \underset{s}{Al^{3+}} + \underset{3s}{3OH^-}$
and $\underset{0.2M}{NaOH} \rightleftharpoons \underset{0.2M}{Na^+} + \underset{0.2M}{OH^-}$
$[Al^{3+}] = s$ and $[OH^-] = 3s + 0.2 \approx 0.2$
$K_{sp} = 2.4 \times 10^{-24} = [Al^{3+}][OH]^3$
$2.4 \times 10^{-24} = s(0.2)^3$
$s = \frac{2.4 \times 10^{-24}}{8 \times 10^3} = 3 \times 10^{-22} mol/L$

Question160

The molar solubility of $Cd(OH)_2$ is $1.84 \times 10^{-5}M$ in water. The expected solubility of $Cd(OH)_2$ in a buffer solution $pH = 12$ is: [April 12, 2019 (II)]

Options:

A. $1.84 \times 10^{-9}M$

B. $\frac{2.49}{1.84} \times 10^{-9}M$

C. $6.23 \times 10^{-11}M$

D. $2.49 \times 10^{-10}M$

Answer: D

Solution:

Solution:
$Cd(OH)_2 \rightleftharpoons \underset{s}{Cd^{2+}} + \underset{2s}{2OH^-}$
At equilibrium, $K_{sp} = s(2s)^2 = 4s^3$
$\Rightarrow K_{sp} = 4 \times (1.84 \times 10^{-5})^{-3}$
Solubility in buffer solution having $pH = 12$
$[OH^-] = 10^{-2}$
$Cd(OH)_2 \rightleftharpoons Cd^{2+} + 2OH^-$

$2s' + 10^{-2} \approx 10^{-2}$

$\therefore K_{sp} = 4 \times (1.84 \times 10^{-5})^3 = s'(10^{-2})^2$

$\Rightarrow s' = \frac{24.9 \times 10^{-15}}{10^{-4}} = 2.49 \times 10^{-10} M$

Question161

The pH of a 0.02M NH_4Cl solution will be [given $K_b(NH_4OH) = 10^{-5}$ and log 2 = 0.301] [April 10, 2019 (II)]

Options:

A. 2.65

B. 4.35

C. 4.65

D. 5.35

Answer: D

Solution:

Solution:

$pH = 7 - \frac{1}{2}pK_b - \frac{1}{2}\log C$

$= 7 - \frac{5}{2} - \frac{1}{2}(\log 2 \times 10^{-2}) = 5.35$

$pH = 5.35$

Question162

If solubility product of $Zr_3(PO_4)_4$ is denoted by K_{sp} and its molar solubility is denoted by S, then which of the following relation between S and K_{sp} is correct? [April 8, 2019 (I)]

Options:

A. $S = \left(\frac{K_{gp}}{144}\right)^{1/6}$

B. $S = \left(\frac{K_{sp}}{6912}\right)^{1/7}$

C. $S = \left(\frac{K_{\phi}}{929}\right)^{1/9}$

D. $S = \left(\frac{K_{sp}}{216} \right)^{1/7}$

Answer: B

Solution:

Solution:
$Zr_3(PO_4)_4 \rightleftharpoons \underset{3S}{3Zr^{4+}} + \underset{4S}{3SO_4^{3-}}$
$K_{sp} = [Zr^{4+}]^3[PO_4^{3-}]^4 = (3S)^3(4S)^4$
$K_{sp} = 6912S^7$
$S = \left(\frac{K_{sp}}{6912} \right)^{1/7}$

Question163

At a certain temperature in a 5L vessel, 2 moles of carbon monoxide and 3 moles of chlorine were allowed to reach equilibrium according to the reaction, $CO + Cl_2 \rightleftharpoons COCl_2$ At equilibrium, if one mole of CO is present then equilibrium constant (K_c) for the reaction is:

[Online April 15, 2018 (II)]

Options:

A. 2.5

B. 4

C. 2

D. 3

Answer: A

Solution:

Solution:
Initially 2 moles of CO are present.
At equilibrium, 1 mole of CO is present
Hence, $2 - 1 = 1$ moles of CO has reacted.
l. mole of CO will react with 1 mole of Cl_2 to form 1 mole of $COCl_2$
$3 - 1 = 2$ moles of Cl_2 remains at equilibrium The equilibrium constant

$$K_c = \frac{[COCl_2]}{[CO][Cl_2]} = \frac{\frac{1mol}{5L}}{\frac{1mol}{5L} \times \frac{2mol}{5L}} = 2.5$$

Question164

In which of the following reactions, an increase in the volume of the container will favour the formation of products?

[Online April 15,2018(I)]

Options:

A. $4NH_3(g) + 5O_2(g) \rightleftharpoons 4NO(g) + 6H_2O(l)$

B. $2NO_2(g) \rightleftharpoons 2NO(g) + O_2(g)$

C. $3O_2 \rightleftharpoons 2O_3(g)$

D. $H_2(g) + I_2(g) \rightleftharpoons 2HI(g)$

Answer: B

Solution:

Solution:
Volume $\uparrow P \downarrow$, reaction proceeds in which direction where the number of moles of gases increases.
$2NO_2(g) \rightleftharpoons 2NO(g) + O_2(g) \quad \Delta n_g = (2+1) - 2 = 1$

Question165

The gas phase reaction $2NO_2(g) \longrightarrow N_2O_4(g)$ is an exothermic reaction. The decomposition of N_2O_4, in equilibrium mixutre of $NO_2(g)$ and $N_2O_4(g)$, can be increased by:
[Online April 16, 2018]

Options:

A. addition of an inert gas at constant pressure

B. lowering the temperature

C. increasing the pressure

D. addition of an inert gas at constant volume

Answer: C

Solution:

Solution:
Reaction at equilibirium
$N_2O_4 \rightleftharpoons 2NO_2$
According to Le chatelier's principle-
(a) addition of an inert gas at constant pressure will increase volume and equilibirium shifts towards more number of molecules.
(b) Decomposition of N_2O_4 will be endothermic, so reaction will move in forward reaction when temperature is increased.So, It is incorrect. It will not effect reaction (volume is constant)
(c) Increasing the pressure on a gas reaction shifts theposition of equilibirium towards thesidewith fewer molecules. So, it will move in backward direction which leads to formation of N_2O_4 from NO_2.

Question166

Following four solutions are prepared by mixing different volumes of N aOH and H Cl of different concentrations, pH of which one of them will be equal to 1 ? [Online April 15, 2018 (II)]

Options:

A. $55\text{mL}\ \frac{\text{M}}{10}\text{H Cl}\ + 45\text{mL}\ \frac{\text{M}}{10}\text{N aOH}$

B. $75\text{mL}\ \frac{\text{M}}{5}\text{H Cl}\ + 25\text{mL}\ \frac{\text{M}}{5}\text{N aOH}$

C. $100\text{mL}\ \frac{\text{M}}{10}\text{H Cl}\ + 100\text{mL}\ \frac{\text{M}}{10}\text{N aOH}$

D. $60\text{mL}\ \frac{\text{M}}{10}\text{H Cl}\ + 40\text{mL}\ \frac{\text{M}}{10}\text{N aOH}$

Answer: B

Solution:

Solution:

$75\text{mL}\ \frac{\text{M}}{5}\text{H Cl}\ + 25\text{mL}\ \frac{\text{M}}{5}\text{N aOH}$

$25\text{mL}\ \frac{\text{M}}{5}\text{N aOH}$ will neutralise $25\text{mL}\ \frac{\text{M}}{5}\text{H Cl}$

$75 - 25 = 50\text{mL}\ \frac{\text{M}}{5}\text{H Cl}$ will remain.

Total volume will be $75 + 25 = 100\text{mL}$

$50\text{mL}\ \frac{\text{M}}{5}\text{H Cl}$ is diluted to 100mL

$[\text{H}^+] = [\text{H Cl}] = \frac{\text{M}}{5} \times \frac{50}{100} = \frac{\text{M}}{10}$

$\text{pH} = -\log_{10}[\text{H}^+] = -\log_{10}\frac{\text{M}}{10} = 1$

Question167

Which of the following is a Lewis acid? [Online April 15, 2018 (I)]

Options:

A. PH_3

B. NF_3


C. NaH

D. $B(CH_3)_3$

Answer: D

Solution:

Solution:



Question168

Which of the following salts is the most basic in aqueous solution? [2018]

Options:

A. $Al(CN)_3$

B. CH_3COOK

C. $FeCl_3$

D. $Pb(CH_3COO)_2$

Answer: B

Solution:

Solution:
CH_3COOK is a salt of weak acid (CH_3COOH) and strong base (KOH).
$FeCl_3$ is a salt of weak base [$Fe(OH)_3$] and strong acid (HCl).
$Pb(CH_3COO)_2$, is a salt of weak base $Pb(OH)_2$ and weak acid (CH_3COOH)
$Al(CN)_3$ is a salt of weak base [$Al(OH)_3$] and weak acid (HCN).

Question169

An aqueous solution contains 0.10M H_2S and 0.20M HCl. If the equilibrium constants for the formation of HS from H_2S is 1.0×10^{-7} and that of S^{2-} from HS - ions is 1.2×10^{-13} then the concentration of S^{2-} ions in aqueous solution is : [2018]

Options:

A. 5×10^{8}

B. 3×10^{-20}

C. 6×10^{-21}

D. 5×10^{-19}

Answer: B

Solution:

Solution:

$\underset{\text{Atequb. } 0.10}{H_2S} \rightleftharpoons \underset{0.20}{2H^+} + \underset{?}{S^{2-}}, K_{a_1} \cdot K_{a_2} = K_{eq}$

$\therefore \frac{[H^+]^2[s^{2-}]}{[H_2S]} = 1 \times 10^{-7} \times 1.2 \times 10^{-13}$

$\frac{[0.2]^2[s^{2-}]}{[0.1]} = 1.2 \times 10^{-20}$

$[S^2] = 3 \times 10^{-20}$

Question170

An aqueous solution contains an unknown concentration of Ba^{2+}. When 50mL of a 1M solution of Na_2SO_4 is added, $BaSO_4$ just begins to precipitate. The final volume is 500 mL. The solubility product of $BaSO_4$ is 1×10^{-10}. What is the original concentration of Ba^{2+}?
[2018]

Options:

A. $5 \times 10^{-9}M$

B. $2 \times 10^{-9}M$

C. $1.1 \times 10^{-9}M$

D. $1.0 \times 10^{-10}M$

Answer: C

Solution:

Solution:

Concentration of SO_4^{2-} in $BaSO_4$ solution

$M_1V_1 = M_2V_2$

$1 \times 50 = M_2 \times 500$

$M_2 = \frac{1}{10}$

For just precipitation

Ionic product $= K_{sp}$

$[Ba^{2+}][SO_4^2] = K_{sp}(BaSO_4)$

$[Ba^{2+}] \times \frac{1}{10} = 10^{-10}$

$[Ba^{2+}] = 10^{-9}M$ in 500mL solution

Thus $[Ba^{2+}]$ in original solution

$(500 - 50 = 450mL)$

$\Rightarrow M_1 \times 450 = 10^{-9} \times 500$

[where $M_1 =$ Molarity of original solution]

$M_1 = \frac{500}{450} \times 10^{-9} = 1.11 \times 10^{-9}M$

Question171

The minimum volume of water required to dissolve 0.1g lead (II) chloride to get a saturated solution (K_{sP} of $PbCl_2$ = 3.2×10^{-8}; atomic mass of Pb = 207u) is: [Online April 15, 2018 (I)]

Options:

A. 1.798L

B. 0.36L

C. 17.95L

D. 0.18L

Answer: D

Solution:

Solution:

$(K_{sp})_{PbCl_2} = 3.2 \times 10^{-8} = 32 \times 10^{-9}$

$PbCl_2 \rightleftharpoons Pb^{2+} + \underset{s}{2Cl^-}$

$K_{sp} = [Pb^{2+}][Cl^-]^2$

$K_{sp} = 4s^3 = 32 \times 10^{-9}$

$s^3 = 8 \times 10^{-9}$

$s = 2 \times 10^{-3}M$

$\frac{W}{M \cdot W} \times \frac{1}{V_L} = 2 \times 10^{-3}$

$\frac{0.1}{278} \times \frac{1}{V_L} = 2 \times 10^{-3}$

$V_L = \frac{0.1 \times 1000}{278 \times 2} = 0.18L$

Question172

The following reaction occurs in the Blast Furnace where iron ore is reduced to iron metal :

$Fe_2O_3(s) + 3CO(g) \rightleftharpoons 2Fe(l) + 3CO_2(g)$

Using the Le Chatelier's principle, predict which one of the following will not disturb the equilibrium? [Online April 9, 2017]

Options:

A. Removal of CO

B. Removal of CO_2

C. Addition of CO_2

D. Addition of Fe_2O_3

Answer: D

Solution:

Solution:

Perturbation	Shifts reaction towards
Removal of CO	Left
Removal of CO_2	Right
Addition of CO_2	Left
Addition of Fe_2O_3	No change (This is a solid compound. Its concentration has noeffect on the equilibrium.)

Question173

50mL of 0.2M ammonia solution is treated with 25mL of 0.2M HCl. If pK_b of ammonia solution is 4.75, the pH of the mixture will be: [Online April 9, 2017]

Options:

A. 3.75

B. 4.75

C. 8.25

D. 9.25

Answer: D

Solution:

Solution:

$NH_3 + HCl \longrightarrow NH_4Cl$

moles of $HCl = 0.2M \times 25 \times 10^{-3}L = 0.005$ moles HCl (total consumed)

moles of $NH_3 = 0.2M \times 50 \times 10^{-3}L = 0.01 mol es HCl$

excess $NH_3 = 0.01 - 0.005 = 0.005 mol es$

1 mole ammonia $= 1 mol e NH_4Cl$

$0.005 NH_3 = 0.005 NH_4Cl$

Total volume $= V_{HCl} + V_{NH_3} = 25 + 50 = 75mL$ $[NH_3] = [NH_4Cl] = \frac{0.005 mol e}{75 \times 10^{-3}L} = 0.066M$

$pOH = pK_b + \log \frac{[NH_4Cl]}{[NH_3]}$

$$pOH = 4.75 + \log \frac{[0.066]}{[0.066]}$$
$$pOH = 4.75$$
$$pH = 14 - pOH \Rightarrow pH = 9.25$$

Question174

pK_a of a weak acid (HA) and pK_b of a weak base (BOH) are 3.2 and 3.4, respectively. The pH of their salt (AB) solution is
[2017]

Options:

A. 7.2

B. 6.9

C. 7.0

D. 1.0

Answer: B

Solution:

Solution:
The salt (AB) given is a salt of weak acid and weak base. Hence the pH can be calculated by the following formula

$$\therefore pH = 7 + \frac{1}{2}pK_a - \frac{1}{2}pK_b$$
$$= 7 + \frac{1}{2}(3.2) - \frac{1}{2}(3.4) = 6.9$$

Question175

Addition of sodium hydroxide solution to a weak acid (HA) results in a buffer of pH . . If ionisation constant of H A is 10^{-5}, the ratio of salt to acid concentration in the buffer solution will be:
[Online April 8, 2017]

Options:

A. 4 : 5

B. 1 : 10

C. 10 : 1

D. 5 : 4

Answer: C

Solution:

Solution:

$HA \rightleftharpoons H^+ + A^-$

(Unionized, weak acid and common ion effect)

$HA + NaOH \longrightarrow NaA + H_2O$

$NaA \longrightarrow Na^+ + A^-$ (ionized)

$K_a = \frac{[H^+][A^-]}{[HA]}$

Given, $pH = 6, [H^+] = 1 \times 10^{-6}$

$[H^+] = \frac{K_a[\text{ Acid }]}{[\text{ Salt }]}$

$\frac{[\text{ Salt }]}{[\text{ Acid }]} = \frac{K_a}{[H^+]} = \frac{10^{-5}}{10^{-6}} = 10:1$

Question176

The equilibrium constant at 298K for a reaction $A + B \rightleftharpoons C + D$ is 100 . If the initial concentration of all the four species were 1 M each, then equilibrium concentration of D (in mol L^{-1}) will be:

[2016]

Options:

A. 1.818

B. 1.182

C. 0.182

D. 0.818

Answer: A

Solution:

Solution:

Given,	$A + B \rightleftharpoons C + D$			
No. of moles initially	1	1	1	1
At eqm.	$1-x$	$1-x$	$1+x$	$1+x$

Given, $A + B \rightleftharpoons C + D$

$K_c = \left(\frac{1+a}{1-a}\right)^2 = 100; \ \frac{1+a}{1-a} = 10$

On solving; $a = 0.81$

$[D]_{At_{Oq}} = 1 + a = 1 + 0.81 = 1.81$

Question177

A solid XY kept in an evacuated sealed container undergoes decomposition to form a mixture of gases X and Y at temperature T . The equilibrium pressure is 10 bar in the vessel. K_p for this reaction is:

[Online April 10,2016]

Options:

A. 25

B. 100

C. 10


D. 5

Answer: A

Solution:

Solution:
$X\,Y\,(s) \rightleftharpoons X\,(g) + Y\,(g)$
$\quad\quad\quad

Solution:



At $T < 1200K$, carbon will reduce $MO(s)$ to $M(s)$ hence, chemical reaction $MO(s) + C(s) \longrightarrow M(s) + CO(g)$ is spontaneous.

Question179

For the reaction, $A(g) + B(g) \rightarrow C(g) + D(g)$, ΔH° and ΔS° are, respectively, $-29.8 kJ\ mol^{-1}$ and $-0.100 kJ\ K^{-1} mol^{-1}$ at 298K. The equilibrium constant for the reaction at 298K is : [Online April 9, 2016]

Options:

A. 1.0×10^{-10}

B. 10

C. 1

D. 1.0×10^{10}

Answer: C

Solution:

Solution:
Given $\Delta H^\circ = -29.8 kJ\ mol^{-1}$
$\Delta S^\circ = -1.00 kJ\ K^{-1}$
From the equation
$\Delta G^\circ = \Delta H^\circ - T\ \Delta S^\circ = -29.8 - (298 \times -0.100)$
$= -29.8 + 29.8 = 0$
Now, $\Delta G^\circ = -2.303 RT\ \log K_{eq}$
$0 = -2.303 RT\ \log K_{eq}$
$\therefore K_{eq} = 1.$

Question180

Gaseous N_2O_4 dissociates into gaseous NO_2 according to the reaction $[N_2O_4(g) \rightleftharpoons 2NO_2(g)]$

At 300K and 1 atm pressure, the degree of dissociation of N_2O_4 is 0.2. If one mole of N_2O_4 gas is contained in a vessel, then the density of the equilibrium mixture is : [Online April 10, 2015]

Options:

A. 1.56g/L

B. 6.22g/L

C. 3.11gL

D. 4.56g/L

Answer: C

Solution:

Solution:

$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

t = 0	1	0
t = eqm	1 − α	2α

$N_2O_4(g) \rightleftharpoons 2NO_2(g)$

Where α = degree of dissociation.

∴ Mol. wt. of mixture

$$= \frac{(1-\alpha) \times M_{N_2O_4} + 2\alpha \times M_{NO_2}}{(1-\alpha+2\alpha)}$$

$$= \frac{(1-0.2)92 + 2 \times 0.2 \times 46}{(1+0.2)} = 76.66$$

Now, as per ideal gas equation,

$PV = nRT$

$PM_{mix} = dRT$

$$\therefore \ d = \frac{PM_{mix}}{RT} = \frac{1 \times 76.66}{0.0821 \times 300} = 3.11 g/L$$

Question181

The standard Gibbs energy change at 300K for the reaction $2A \rightleftharpoons B + C$ is 2494.2J . At a given time, the composition of the reaction mixture is [A] = $\frac{1}{2}$, [B] = 2 and [C] = $\frac{1}{2}$. The reaction proceeds in the : [R = 8.314J / K / mol , e = 2.718] [2015]

Options:

A. forward direction because $Q < K_c$

B. reverse direction because $Q < K_c$

C. forward direction because $Q > K_c$

D. reverse direction because $Q > K_c$

Answer: D

Solution:

Solution:
$\Delta G^\circ = 2494.2J$
$2A \rightleftharpoons B + C.$
$[A] = \frac{1}{2}, [B] = 2, [C] = \frac{1}{2}$
$Q = \frac{[B][C]}{[A]^2} = \frac{2 \times 1/2}{\left(\frac{1}{2}\right)^2} = 4$
$\Delta G^\circ = -2.303RT \log K_c$
$2494.2J = -2.303 \times (8.314J/K/mol) \times (300K) \log K_c$
$\Rightarrow \log K_c = -\frac{2494.2J}{2.303 \times 8.314J/K/mol \times 300K}$
$\Rightarrow \log K_c = -0.4341; K_c = 0.37; Q > K_c$

Question182

The following reaction is performed at 298K .
$2NO(g) + O_2(g) \rightleftharpoons 2NO_2(g)$
The standard free energy of formation of NO(g) is 86.6 KJ/mol at 298K .
What is the standard free energy of formation of $NO_2(g)$ at 298K ?($K_p = 1.6 \times 10^{12}$)
[2015]

Options:

A. $86600 - \frac{\ln(1.6 \times 10^{12})}{R(298)}$

B. $0.5[2 \times 86,600 - R(298) \ln(1.6 \times 10^{12})]$

C. $R(298) \ln(1.6 \times 10^{12}) - 86600$

D. $86600 + R(298) \ln(1.6 \times 10^{12})$

Answer: B

Solution:

Solution:
$\Delta G_{rex^n}^\circ = 2\Delta G_f^\circ(NO_2) - 2\Delta G_f^\circ(NO) - \Delta G_f^\circ(O_2)$
$2\Delta G_f^\circ(NO_2) = \Delta G_{rex^n}^\circ + 2\Delta G_f^\circ(NO) + \Delta G_f^\circ(O_2)$
$\because \quad \Delta G = \Delta G^\circ + RT \ln K_p$
At equilibrium,
$\Delta G = 0, Q = K_p; \Delta G^\circ = -R \ln K_p$
$\Delta G_j^\circ(O_2) = 0$
$\therefore \Delta G_f^\circ(NO_2) = \frac{1}{2}[2 \times 86600 - R(298) \ln(1.6 \times 10^{12})]$

Question183

The increase of pressure on ice ⇌ water system at constant temperature will lead to [Online April 11, 2015]

Options:

A. a decrease in the entropy of the system

B. an increase in the Gibb's energy of the system

C. no effect on the equilibrium

D. a shift of the equilibrium in the forward direction

Answer: D

Solution:

Solution:
Volume of ice is greater than that of water. The direction in which the reaction will proceed can be predicted by applying Le-Chatelier's principle

Pressure $\propto \frac{1}{\text{Volume}}$

So equilibrium, will shift forward.

Question184

For the reaction $SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$, if $K_p = K_c(RT)^x$ where the symbols have usual meaning then the value of x is (assuming ideality): [2014]

Options:

A. -1

B. $-\frac{1}{2}$

C. $\frac{1}{2}$

D. 1

Answer: B

Solution:

Solution:

$SO_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons SO_3(g)$

$K_p = K_c(RT)^x$

where $x = \Delta n_g =$ number of gaseous moles in product − number of gaseous moles in reactants

$= 1 - \left(1 + \frac{1}{2}\right) = 1 - \frac{3}{2} = -\frac{1}{2}$

Question185

At a certain temperature, only 50%H I is dissociated into H_2 and I_2 at equilibrium.

The equilibrium constant is:
[Online April 9, 2014]

Options:

A. 1.0

B. 3.0

C. 0.5

D. 0.25

Answer: D

Solution:

Solution:

$$2HI \rightleftharpoons H_2 + I_2$$

At t = 0	c	0	0
At eqm.	c−cα	$\frac{c\alpha}{2}$	$\frac{c\alpha}{2}$

$2HI \rightleftharpoons H_2 + I_2$

At eqm. c-cα $\frac{c\alpha}{2}$ $\frac{c\alpha}{2}$

Now, $K_c = \frac{\left(\frac{c\alpha}{2}\right)\left(\frac{c\alpha}{2}\right)}{(c - c\alpha)^2}$

$K_C = \frac{\alpha}{4(1-\alpha)^2}$; $K_c = \frac{0.5}{4(1-0.5)^2}$

$K_0 = 0.25$

Question186

For the decomposition of the compound, represented as
$NH_2COONH_4(s) \rightleftharpoons 2NH_3(g) + CO_2(g)$ the $K_p = 2.9 \times 10^{-5} atm^3$.
If the reaction is started with 1mol of the compound, the total pressure at equilibrium would be:
[Online April 19, 2014]

Options:

A. 1.94×10^{-2} atm

B. 5.82×10^{-2}atm

C. 7.66×10^{-2}atm

D. 38.8×10^{-2}atm

Answer: B

Solution:

Solution:
$NH_2COONH_4(s) \rightleftharpoons 2NH_3(g) + CO_2(g)$

$K_p = \frac{(p_{NH_3})^2 \times (p_{Co_2})}{P_{NH_2COONH_4}(s)} = (p_{NH_3})^2 \times (p_{CO_2})$

As evident by the reaction, NH_3 and CO_2 are formed in molar ratio of $2:1$.

Thus if P is the total pressure of the system at equilibrium, then

$p_{NH_3} = \frac{2}{3} \times p \quad p_{CO_2} = \frac{1}{3} \times P$

$K_p = \left(\frac{2P}{3}\right)^2 \times \frac{P}{3} = \frac{4P^3}{27}$

Given, $K_p = 2.9 \times 10^{-5}$

$\therefore 2.9 \times 10^{-5} = \frac{4P^3}{27}$

$P^3 = \frac{2.9 \times 10^{-5} \times 27}{4}$

$P = \left(\frac{2.9 \times 10^{-5} \times 27}{4}\right)^{1/3} = 5.82 \times 10^{-2}$atm

---

Question187

What happens when an inert gas is added to an equilibrium keeping volume unchanged?
[Online April 12, 2014]

Options:

A. More product will form

B. Less product will form

C. More reactant will form

D. Equilibrium will remain unchanged

Answer: D

Solution:

Solution:
On adding inert gas at constant volume the total pressure of the system is increased, but the partial pressure of each reactant and product remains the same. Hence no effect on the state of equilibrium.

Question188

The conjugate base of hydrazoic acid is:
[Online April 12, 2014]

Options:

A. N^{-3}

B. N_3^-

C. N_2^-

D. HN_3^-

Answer: B

Solution:

Solution:
$\underset{\text{Hydrazoic acid}}{N_3H} \rightleftharpoons N_3^- + H^+$
i.e, conjugate base of hydrazoic acid is N_3.

Question189

Assuming that the degree of hydrolysis is small, the pH of 0.1M solution of sodium acetate ($K_a = 1.0 \times 10^{-5}$) will be:
[Online April 11, 2014]

Options:

A. 5.0

B. 6.0

C. 8.0

D. 9.0

Answer: D

Solution:

Solution:
Sodium acetate is a salt of strong base and weak acid.
$\therefore pH = 7 + \frac{1}{2}pK_a + \frac{1}{2}\log c$ where $pK_a = -\log K_a$
$= 7 + \frac{5}{2} - \frac{1}{2}$
$= 9.0$
$= -\log 10^{-5} = 5$
$\log c = \log 10^{-1} = -1$

Question190

In some solutions, the concentration of H_3O^+ remains constant even when small amounts of strong acid or strong base are added to them. These solutions are known as:
[Online April 11, 2014]

Options:

A. Ideal solutions

B. Colloidal solutions

C. True solutions

D. Buffer solutions

Answer: D

Solution:

Solution:
Solutions which resist the change in the value of pH when small amount of acid or base is added to them are known as buffers.

Question191

Consider the following equilibrium
$AgCl \downarrow +2NH_3 \rightleftharpoons [Ag(NH_3)_2]^+ + Cl^-$
White precipitate of AgCl appears on adding which of the following?
[Online April 11,2014]

Options:

A. NH_3

B. aqueous NaCl

C. aqueous HNO_3

D. aqueous NH_4Cl

Answer: C

Solution:

Solution:
$2HNO_3(aq) + [Ag(NH_3)_2]^+ + Cl^- \longrightarrow AgCl(s) \downarrow +2NH_4 + 2NO_3^-$
When nitric acid is added to amine solution, solution is made acidic and the complex ion dissociates and liberate silver ion to recombine with chloride ion. This is the conformatory test for silver in group 1.

Question192

Zirconium phosphate [$Zr_3(PO_4)_4$] dissociates into three zirconium cations of charge +4 and four phosphate anions of charge −3. If molar solubility of zirconium phosphate is denoted by S and its solubility product by K_{sp} then which of the following relationship between S and K_{sp} is correct?
[Online April 19,2014]

Options:

A. $S = \{K_{sp}/(6912)^{1/7}\}$

B. $S = \{K_{sp}/144\}^{1/7}$

C. $S = \{K_{sp}/6912\}^{1/7}$

D. $S = \{K_{sp}/6912\}^{7}$

Answer: C

Solution:

Solution:

$$[Zr_3(PO_4)_4] \rightleftharpoons \underset{3s}{3Zr^{4+}} + \underset{4s}{4PO_4^{3-}}$$

$K_{sp} = (3s)^3(4s)^4$

$= 27s^3 \times 256s^4$

$= 6912s^7$

$\therefore s = \left(\frac{K_{sp}}{6912}\right)^{1/7}$

Question193

In reaction A + 2B ⇌ 2C + D, initial concentration of B was 1.5 times of [A], but at equilibrium the concentrations of A and B became equal. The equilibrium constant for the reaction is :
[Online April 9, 2013]

Options:

A. 8

B. 4

C. 12

D. 6

Answer: B

Solution:

Solution:

	A	+	2B	$\rightleftharpoons$	2C	+	D
At t = 0	a		$1.5a$		0		0
At eqns.	$(a-x)$		$(1.5a-2x)$		$2x$		x

Hence, $K_c = \frac{(2x)^2 \times x}{(a-x)(1.5a-2x)^2}$

Given, at equilibrium

$\therefore \ (a-x) = (1.5a-2x)$

$\therefore \ a = 2x$

On solving $K_c = 4$

Question194

(1) $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$, K_1

(2) $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$, K_2

(3) $H_2(g) + \frac{1}{2}O_2(g) \rightleftharpoons H_2O(g)$, K_3

The equation for the equilibrium constant of the reaction $2NH_3(g) + \frac{5}{2}O_2(g) \rightleftharpoons 2NO(g) + 3H_2O(g)$, ($K_4$) in ter of K_1, K_2 and K_3 is : [Online April 23, 2013]

Options:

A. $\frac{K_1 \cdot K_2}{K_3}$

B. $\frac{K_1 \cdot K_3^2}{K_2}$

C. $K_1K_2K_3$

D. $\frac{K_2 \cdot K_3^3}{K_1}$

Answer: D

Solution:

Solution:

To calculate the value of K_4 in the given equation we should apply:

eqn. (2) $+$cqn . (3) $\times 3$ -eqn. (1)

hence $K_4 = \frac{K_2K_3^3}{K_1}$

Question195

The ratio $\frac{K_p}{K_c}$ for the reaction is:

[Online April 25, 2013]

Options:

A. $\frac{1}{\sqrt{RT}}$

B. $(RT)^{1/2}$

C. RT

D. 1

Answer: A

Solution:

Solution:
$K_p = K_c(RT)^{\Delta n_g}$
For the reaction
$CO(g) + \frac{1}{2}O_2(g) \rightleftharpoons CO_2(g)$
$\Delta n_g = 1 - \left(1 + \frac{1}{2}\right) = -\frac{1}{2}$
$\therefore K_p = \frac{K_c}{\sqrt{RT}}; \quad \frac{K_p}{K_c} = \frac{I}{\sqrt{RT}}$

Question196

How many litres of water must be added to 1 litre an aqueous solution of H Cl with a pH of 1 to create an aqueous solution with pH of 2 ?
[2013]

Options:

A. 0.1L

B. 0.9L

C. 2.0L

D. 9.0L

Answer: D

Solution:

Solution:
$\because pH = 1; H^+ = 10^{-1} = 0.1M$
$pH = 2; H^+ = 10^{-2} = 0.01M$
$\therefore M_1 = 0.1 \quad V_1 = 1$
$M_2 = 0.01, V_2 = ?$
From
$M_1V_1 = M_2V_2$
$0.1 \times 1 = 0.01 \times V_2$
$V_2 = 10L$
$\therefore$ Volume of water added $= 10 - 1 = 9L$

Question197

N aOH is a strong base. What will be pH of 5.0×10^{-2}M N aOH solution ? (log 2 = 0.3) [Online April 22, 2013]

Options:

A. 14.00

B. 13.70

C. 13.00

D. 12.70

Answer: D

Solution:

Solution:
Given $[OH^-] = 5 \times 10^{-2}$
$\therefore pOH = -\log 5 \times 10^{-2}$
$= -\log 5 + 2 \log 10 = 1.30$
$\because pH + pOH = 14$
$\because pH = 14 - pOH$
$= 14 - 1.30 = 12.70$

Question198

Equimolar solutions of the following compounds are prepared separately in water. Which will have the lowest pH value? [Online April 23, 2013]

Options:

A. $BeCl_2$

B. $SrCl_2$

C. $CaCl_2$

D. $MgCl_2$

Answer: A

Solution:

Solution:
Metal halide on hydrolysis with water form corresponding hydroxides.
The basic strength of hydroxide increases as we move down in a group. This is because of the increase in size which results in decrease of ionization energy which weakens the strength of $M-O$ bonds in MOH and thus increases the basic

strength.

$\underset{\text{Amphoteric}}{Be(OH)_2}$ $\underset{\text{Weak base}}{Mg(OH)_2}$

$Ca(OH)_2$ $\underset{\text{Strong base}}{Sr(OH)_2}$ $Ba(OH)_2$

Hence, $Be(OH)_2$ will have lowest pH.

Question199

What is the pH of a 10^{-4}M OH^- solution at 330K, if K_w at 330K is $10^{-13.6}$? [Online April 23, 2013]

Options:

A. 4

B. 9.0

C. 10

D. 9.6

Answer: D

Solution:

Solution:
Given at 330K

$K_w = 10^{-13.6}$

i.e. $pK_w = pH + pOH$

$\because pOH = -\log[OH^-]$

$13.6 = pH + pOH$

$pOH = -\log 10^{-4}$

$pOH = 4$

$\therefore pH = 13.6 - 4$

$= 9.6$

Question200

Which one of the following arrangements represents the correct order of the proton affinity of the given species: [Online April 25, 2013]

Options:

A. $I^- < F^- < HS^- < NH_2^-$

B. $HS^- < NH_2^- < F^- < I^-$

C. $F^- < I^- < NH_2^- < HS$

D. $NH_2^- < HS^- < I^- < F^-$

Answer: A

Solution:

Solution:
The species with the greatest proton affinity will be the strongest base, and its conjugate acid will be the weakest acid. The weakest acid will have the smallest value of K_a. Since HI is a stronger acid than HF which is a stronger acid than H_2S, a partial order of proton affinity is
$I^- < F^- < HS^-$
Since NH_3 is a very weak acid, NH_2^- must be a very strong base. Therefore the correct order of proton affinity is
$I^- < F^- < HS^- < NH_2$

Question201

Values of dissociation constant, K_0 are given as follows:

Acid	K_a
HCN	6.2×10^{-10}
HF	7.2×10^{-4}
HNO_2	4.0×10^{-4}

Correct order of increasing base strength of the base CN^- F^- and NO_2^- will be: [Online April 22, 2013]

Options:

A. $F < CN^- < NO_2^-$

B. $NO_2^- < CN^- < F^-$

C. $F < NO_2^- < CN^-$

D. $NO_2^- < F^- < CN^-$

Answer: C

Solution:

Solution:
Higher the value of K_a lower will be the value of pK_a i.e. higher will be the acidic nature. Further since CN^-, F^- and NO_2^- are conjugate base of the acids HCN, HF and HNO_2 respectively hence the correct order of base strength will be
$F^- < NO_2^- < CN$
(∵ stronger the acid weaker will be its conjugate base)

Question202

What would be the pH of a solution obtained by mixing 5g of acetic acid and 7.5g of sodium acetate and making the volume equal to 500mL ?
($K_a = 1.75 \times 10^{-5}$, $pK_a = 4.76$)
[Online April 25, 2013]

Options:

A. pH = 4.70

B. pH < 4.70

C. pH of solution will be equal to pH of acetic acid

D. 4.76 < pH < 5.0

Answer: D

Solution:

Solution:
Concentration of CH_3COOH is computed as under.

$$\text{conc.} = 5g \text{ in } 500mL$$
$$= 10g/L \quad [\text{ Mol. wt. of } CH_3COOH = 60$$
$$[CH_3COOH] = \frac{10}{60}M ; \frac{1}{6}M$$

concentration of CH_3COONa is computed as under.

conc. = 7.5g in 500mL
= 15g/L

$$[CH_3COOH] = \frac{15}{18}M$$
$$pK_a = -\log K_a$$
$$= \log(1.8 \times 10 - 5) = 4.7447$$
$$pH = pK_a + \log \frac{[salt]}{[\text{ acid }]}$$
$$= 4.744 \log \frac{15/82}{1/6}$$
$$= 4.7447 + \log 1.097$$
$$= 4.7447 + 0.0402$$
$$= 4.78$$

Question203

Solid $Ba(NO_3)_2$ is gradually dissolved in a 1.0×10^{-4}M Na_2CO_3 solution. At which concentration of Ba^{2+}, precipitate of $BaCO_3$ begins to form ?(K_{sp} for $BaCO_3$ $= 5.1 \times 10^{-9}$)
[Online April 9, 2013]

Options:

A. 5.1×10^{-5}M

B. 7.1×10^{-8}M

C. 4.1×10^{-5}M

D. 8.1×10^{-7}M

Answer: A

Solution:

Solution:
Conc. of $Na_2CO_3 = 1.0 \times 10^{-4}$M
$\therefore [CO_3^{2}] = 1.0 \times 10^{-4}$M i.e. $s = 1.0 \times 10^{-4}$M
At equilibrium
$[Ba^{2+}][CO_3^{2-}] = K_{sp}$ of $BaCO_3$

$$[Ba^{2+}] = \frac{K_{sp}}{[CO_3^{2-}]} = \frac{5.1 \times 10^{-9}}{1.0 \times 10^{-4}}$$

$= 5.1 \times 10^{-5}$M

Question204

Which one of the following arrangements represents the correct order of solubilities of sparingly soluble salts Hg_2Cl_2, $Cr_2(SO_4)_3$, $BaSO_4$ and $CrCl_3$ respectively? [Online April 22, 2013]

Options:

A. $BaSO_4 > Hg_2Cl_2 > Cr_2(SO_4)_3 > CrCl_3$

B. $BaSO_4 > Hg_2Cl_2 > CrCl_3 > Cr_2(SO_4)_3$

C. $BaSO_4 > CrCl_3 > Hg_2Cl_2 > Cr_2(SO_4)_3$

D. $Hg_2Cl_2 > BaSO_4 > CrCl_3 > Cr_2(SO_4)_3$

Answer: B

Solution:

Solution:
$Cr_2(SO_4)_3 \Longrightarrow 2Cr^{3+} + 3SO_4^{2-}$
$K_{sp} = (2s)^2(3s)^3 = 4s^2 \times 27s^3 = 108s^5$

$$s = \left(\frac{K_{sp}}{108}\right)^{1/5}$$

$Hg_2Cl_2 \rightleftharpoons 2Hg^{2+} + \underset{2s}{2Cl^-}$
$K_{sp} = (2s)^2 \times (2s)^2 = 16s^4$

$$s = \left(\frac{K_{sp}}{16}\right)^{1/4}$$

$BaSO_4 \rightleftharpoons \underset{s}{Ba^{2+}} + \underset{s}{SO_4^{2-}}$
$K_{sp} = s^2$
$s = \sqrt{K_{sp}}$
$CrCl_3 \Longrightarrow \underset{s}{Cr^{3+}} + \underset{3s}{3Cl^-}$
$K_{sp} = s \times (3s)^3 = 27s^4$

$s = \left(\frac{K_{sp}}{27}\right)^{1/4}$

Hence the correct order of solubilities of salts is

$\sqrt{K_{sp}} > \left(\frac{K_{sp}}{16}\right)^{1/4} > \left(\frac{K_{sp}}{27}\right)^{1/4} > \left(\frac{K_{sp}}{108}\right)^{1/5}$

Question205

K_1, K_2 and K_3 are the equilibrium constants of the following reactions (I), (II) and (III) respectively:

(I) $N_2 + 2O_2 \rightleftharpoons 2NO_2$

(II) $2NO_2 \rightleftharpoons N_2 + 2O_2$

(III) $NO_2 \rightleftharpoons \frac{1}{2}N_2 + O_2$

The correct relation from the following is

[Online May 7, 2012]

Options:

A. $K_1 = \frac{1}{K_2} = \frac{1}{K_3}$

B. $K_1 = \frac{1}{K_2} = \frac{1}{(K_3)^2}$

C. $K_1 = \sqrt{K_2} = K_3$

D. $K_1 = \frac{1}{K_2} = K_3$

Answer: B

Solution:

Solution:

(I) $N_2 + 2O_2 \overset{K_1}{\rightleftharpoons} 2NO_2$

$K_1 = \frac{[NO_2]^2}{[N_2][O_2]^2}$ . . . (i)

(II) $2NO_2 \overset{K_2}{\rightleftharpoons} N_2 + 2O_2$

$K_2 = \frac{[N_2][O_2]^2}{[NO_2]^2}$ . . . (ii)

(III) NO_2 ef tarrows $\overset{K_3}{\frac{1}{2}}N_2 + O_2$

$K_3 = \frac{[N_2]^{1/2}[O_2]}{[NO_2]}$

$\therefore (K_3)^2 = \frac{[N_2][O_2]^2}{[NO_2]^2}$ . . . (iii)

∴ From equation (i), (ii) and (iii)

$K_1 = \frac{1}{K_2} = \frac{1}{(K_3)^2}$

Question206

8 mol of $AB_3(g)$ are introduced into a 1.0d m^3 vessel. If it dissociates as $2AB_3(g) \rightleftharpoons A_2(g) + 3B_2(g)$. At equilibrium, 2 mol of A_2 are found to be present. The equilibrium constant of this reaction is
[Online May 12, 2012]

Options:

A. 2

B. 3

C. 27

D. 36

Answer: C

Solution:

Solution:

$$2AB_3(g) \rightleftharpoons A_2(g) + 3B_2(g)$$

at $t = 0$	8	0	0
at eqm.	$(8 - 2 \times 2)$	2	3×2
	$= 4$	2	6
molar cones.	$4/V$	$2/V$	$6/V$

$2AB_3(g) \leftrightarrows A_2(g) + 3B_2(g)$

now $K_c = \frac{[A_2][B_2]^3}{[AB_3]^2} = \frac{2/1 \times [6/1]^3}{[4/1]^2} = 27$

Question207

The value of K_p for the equilibrium reaction $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ is 2 . The percentage dissociation of $N_2O_4(g)$ at a pressure of 0.5 atm is
[Online May 19, 2012]

Options:

A. 25

B. 88

C. 50

D. 71

Answer: D

Solution:

Solution:

$$N_2O_4(g) \rightleftharpoons 2NO_2(g)$$

Initial moles	1	0
Moles at eqm.	$(1 - \alpha)$	2α

(α = degree of dissociation)

Total number of moles at eqm.
$= (1 - \alpha) + 2\alpha$
$= (1 + \alpha)$

$$p_{N_2O_4} = \frac{(1-\alpha)}{a(1+\alpha)} \times P$$

$$p_{NO_2} = \frac{2\alpha}{a(1+\alpha)} \times P$$

$$K_p = \frac{(p_{NO_2})^2}{P_{N_2O_4}} = \frac{\left(\frac{2\alpha}{(1+\alpha)} \times P\right)^2}{\left(\frac{1-\alpha}{1+\alpha}\right) \times P}$$

$$K_p = \frac{4\alpha^2 P}{\frac{(1+\alpha)^2(1-\alpha)}{1+\alpha}}; K_p = \frac{4\alpha^2 P}{(1+\alpha)(1-\alpha)}; = \frac{4\alpha^2 P}{1-\alpha^2}$$

Given, $K_p = 2$, $P = 0.5$atm

$$\therefore K_p = \frac{4\alpha^2 P}{1-\alpha^2}; 2 = \frac{4\alpha^2 \times 0.5}{1-\alpha^2}$$

$\alpha = 0.707 \approx 0.71$
∴ Percentage dissociation
$= 0.71 \times 100 = 71$

Question208

One mole of $O_2(g)$ and two moles of $SO_2(g)$ were heated in a closed vessel of one-litre capacity at 1098K . At equilibrium 1.6 moles of $SO_3(g)$ were found. The equilibrium constant K_c of the reaction would be
[Online May 26, 2012]

Options:

A. 30

B. 40

C. 80

D. 60

Answer: C

Solution:

Solution:

$$O_2 + 2SO_2 \rightleftharpoons 2SO_3$$

At t = 0	1	2	0
At eqm.	$(1 - \alpha)$	$2(1 - \alpha)$	2α

Given at equilibrium,
$2\alpha = 1.6$

$\alpha = 0.8$

$K_c = \frac{(2\alpha)^2}{(1-\alpha)(2-2\alpha)^2} = \frac{(0.8)^2}{(1-0.8)(1-0.8)^2} = \frac{0.64}{0.002}$

$K_c = 80$

Question209

The pH of a 0.1 molar solution of the acid H Q is 3. The value of the ionization constant, K_a of the acid is :

[2012]

Options:

A. 3×10^{-1}

B. 1×10^{-3}

C. 1×10^{-5}

D. 1×10^{-7}

Answer: C

Solution:

Solution:

$H^+ = C\alpha$; $\alpha = \frac{[H^+]}{C}$

or $\alpha = \frac{10^{-3}}{0.1} = 10^{-2}$

$K_a = C\alpha^2 = 0.1 \times 10^{-2} \times 10^{-2} = 10^{-5}$

Question210

Given

(i) $HCN(aq) + H_2O(I) \rightleftharpoons H_3O^+(aq) + CN^-(aq)$

$K_a = 6.2 \times 10^{-10}$

(ii) $CN^-(aq) + H_2O(I) \rightleftharpoons HCN(aq) + OH^-(aq)$

$K_b = 1.6 \times 10^{-5}$

These equilibria show the following order of the relative base strength, [Online May 12, 2012]

Options:

A. $OH^- > H_2O > CN^-$

B. $OH^- > CN^- > H_2O$

C. $H_2O > CN^- > OH^-$

D. $CN^{->}H_2O > OH^-$

Answer: B

Solution:

Solution:
The more is the value of equilibrium constant, the more is the completion of reaction or more is the concentration of products i.e. the order of relative strength would be
$OH^- > CN^- > H_2O$

Question211

The solubility (in mol L^{-1}) of AgCl ($K_{sp} = 1.0 \times 10^{-10}$) in a 0.1 M KCl solution will be [Online May 7, 2012]

Options:

A. 1.0×10^{-9}

B. 1.0×10^{-10}

C. 1.0×10^{-5}

D. 1.0×10^{-11}

Answer: A

Solution:

Solution:
Let solubility of $AgCl = s mol/L$
$AgCl \rightleftharpoons Ag^+ + Cl^-$
i.e., $K_{sp(AgCl)} = s \times s$
$KCl \longrightarrow \underset{0.1}{K^+} + Cl^-$
$[Cl^-]$ from $KCl = 0.1M$
Total $[Cl^-]$ in solution $= s + 0.1$
$K_{cp}(AgCl) = [Ag^+][Cl^-] = s(s + 0.1)$
$1.0 \times 10^{-10} = s(s + 0.1)$
$1.0 \times 10^{-10} = s^2 + 0.1s$
$1.0 \times 10^{-10} = 0.1s$ (as $s^2 \ll < 1$)
$s = 1.0 \times 10^{-9} mol/L$

Question212

If K_{sp} of CaF_2 at 25°C is 1.7×10^{-10}, the combination amongst the following which gives a precipitate of CaF_2 is

[Online May 19, 2012]

Options:

A. 1×10^{-2}M Ca^{2+} and 1×10^{-3}M F^-

B. 1×10^{-4}M Ca^{2+} and 1×10^{-4}M F^-

C. 1×10^{-2}M Ca^{2+} and 1×10^{-5}M F

D. 1×10^{-3}M Ca^{2+} and 1×10^{-5}M F^-

Answer: A

Solution:

Solution:
When ionic product i.e. the product of the concentration of ions in the solution exceeds the value of solubility product, formation of precitpiate occurs.
$CaF_2 \rightleftharpoons Ca^{2+} + 2F^-$
Ionic product $= [Ca^{2+}][F^-]^2$
when, $[Ca^{2+}] = 1 \times 10^{-2}$M
$[F^-]^2 = (1 \times 10^{-3})^2$M
$= 1 \times 10^{-6}$M
$\therefore [Ca^{2+}][F^-]^2 = (1 \times 10^{-2})(1 \times 10^{-6}) = 1 \times 10^{-8}$
In this case,
Ionic product $(1 \times 10^{-8})>$
solubility product (1.7×10^{-10})

Question213

The solubility of PbI_2 at 25°C is $0.7gL^{-1}$. The solubility product of PbI_2 at this temperature is (molar mass of $PbI_2 = 461.2gmol^{-1}$)

[Online May 26, 2012]

Options:

A. 1.40×10^{-9}

B. 0.14×10^{-9}

C. 140×10^{-9}

D. 14.0×10^{-9}

Answer: D

Solution:

Solution:
$PbI_2 \rightleftharpoons \underset{s}{Pb^{2+}} + \underset{2s}{2I^-}$
$K_{cp} = s \times (2s)^2 = 4s^3$

$= 4 \times \left(\frac{0.7}{461.2} \right)^3 = 14.0 \times 10^{-9}$

Question214

An acid HA ionises as
$HA \rightleftharpoons H^+ + A^-$
The pH of 1.0M solution is 5. Its dissociation constant would be : [2011RS]

Options:

A. 5

B. 5×10^{-8}

C. 1×10^{-5}

D. 1×10^{-10}

Answer: D

Solution:

Solution:
pH = 5 means
$[H^+] = 10^{-5}$

	HA $\rightleftharpoons$	H^+ +	A^-
$t = 0$	c	0	0
t_{eq}	$c(1-\alpha)$	$c\alpha$	$c\alpha$

$K_a = \frac{[H^+][A^-]}{[HA]} = \frac{(c\alpha)^2}{c(1-\alpha)} = \frac{[H^+]^2}{c - [H^+]}$

But, $[H^+] << c$
$\therefore K_a = [H^+]^2 = (10^{-5})^2 = 10^{-10}$

Question215

The K_{sp} for $Cr(OH)_3$ is 1.6×10^{-30}. The solubility of this compound in water is : [2011 RS]

Options:

A. $4\sqrt{1.6 \times 10^{-30}}$

B. $4\sqrt{1.6 \times 10^{-30}/27}$

C. $1.6 \times 10^{-30/27}$

D. $2\sqrt{1.6 \times 10^{-30}}$

Answer: B

Solution:

Solution:

$Cr(OH)_3(s) \Longrightarrow Cr^{3+}(aq) + 3OH^-$ (aq.)

$27s^4 = K_{sp}$

$s = \left(\frac{K_{sp}}{27}\right)^{1/4} = \left(\frac{1.6 \times 10^{-30}}{27}\right)^{1/4}$

Question216

In aqueous solution the ionization constants for carbonic acid are $K_L = 4.2 \times 10^{-7}$ and $K_2 = 4.8 \times 10^{-11}$ Select the correct statement for a saturated 0.034M solution of the carbonic acid. [2010]

Options:

A. The concentration of CO_3^{2-} is 0.034M .

B. The concentration of CO_3^{2-} is greater than that of HCO_3^-

C. The concentrations of H^+ and HCO_3^- are approximately equal.

D. The concentration of H^+ is double that of CO_3^{2-}.

Answer: C

Solution:

Solution:

$H_2CO_3 \rightleftharpoons H^+ + HCO_3^- K_1 = 4.2 \times 10^{-7}$

$HCO_3^- \rightleftharpoons H^+ + CO_3^{2-}\ K_2 = 4.8 \times 10^{-11}$

Second dissociation constant (K_2) is much smaller than the first one (K_1). Just a small fraction of total HCO_3^- formed will undergo second stage of ionization. Hence in saturated solution

$[H^+] \gg > [CO_3^{2-}]$; $[CO_3^{2-}] \neq 0.034M$

$[HCO_3^{2-}] > > [CO_3^{2-}]$ and $[H^+] \approx [HCO_3^-]$

Question217

Solubility product of silver bromide is 5.0×10^{-13}. The quantity of potassium bromide (molar mass taken as 120g mol^{-1}) to be added to 1 litre of 0.05M solution of silver nitrate to start the precipitation of AgBr is [2010]

Options:

A. 1.2×10^{-10}g

B. 1.2×10^{-9}g

C. 6.2×10^{-5}g

D. 5.0×10^{8}g

Answer: B

Solution:

Solution:
$AgBr \rightleftharpoons Ag^{+} + Br^{-}$
$K_{sp} = [Ag^{+}][Br^{-}]$
For precipitation to occur
Ionic product > Solubility product
$[Br^{-}] = \frac{K_{sp}}{[Ag^{+}]} = \frac{5 \times 10^{-13}}{0.05} = 10^{-11}$
i.e., precipitation just starts when 10^{-11} moles of KBr is added to $1\ L\ AgNO_3$ solution
$\therefore$ Number of moles of Br^{-} needed from $KBr = 10^{-11}$
$\therefore$ Mass of $KBr = 10^{-11} \times 120 = 1.2 \times 10^{-9}$g

---

Question218

At 25°C, the solubility product of $Mg(OH)_2$ is 1.0×10^{-11}. At which pH , will Mg^{2+} ions start precipitating in the form of $Mg(OH)_2$ from a solution of 0.001M Mg^{2+} ions?
[2010]

Options:

A. 9

B. 10

C. 11

D. 8

Answer: B

Solution:

Solution:
$Mg(OH)_2 \rightleftharpoons Mg^{2+} + 2OH^{-}$
$K_{sp} = [Mg^{2+}][OH^{-}]^2$
$1.0 \times 10^{-11} = 10^{-3} \times [OH^{-}]^2$
$[OH^{-}] = \sqrt{\frac{10^{-11}}{10^{-3}}} = 10^{-4}$
$\therefore\ pOH = 4$
$\therefore\ pH + pOH = 14$
$\therefore\ pH = 10$

Question219

Three reactions involving $H_2PO_4^-$ are given below:

(i) $H_3PO_4 + H_2O \rightarrow H_3O^+ + H_2PO_4^-$ [2010]

(ii) $H_2PO_4^- + H_2O \rightarrow HPO_4^{2-} + H_3O^+$

(iii) $H_2^2PO_4^- + OH^- \rightarrow H_3PO_4 + O^2$

In which of the above does $H_2PO_4^-$ act as an acid ?

[2010]

Options:

A. (ii) only

B. (i) and (ii)

C. (iii) only

D. (i) only

Answer: A

Solution:

Solution:

(i) $\underset{acid_1}{H_3PO_4} + \underset{base_2}{H_2O} \rightarrow \underset{acid_2}{H_3O^+} + \underset{base_1}{H_2PO_4^-}$

(ii) $\underset{acid_1}{H_2PO_4^-} + \underset{base_2}{H_2O} \rightarrow \underset{base_1}{HPO_4^{2-}} + \underset{acid_2}{H_3O^+}$

(iii) $\underset{base_1}{H_2PO_4^-} + \underset{acid_2}{OH^-} \rightarrow \underset{acid_1}{H_3PO_4} + \underset{base_2}{O^{2-}}$

Hence only in (ii) reaction, $H_2PO_4^-$ is acting as an acid.

Question220

Solid $Ba(NO_3)_2$ is gradually dissolved in a 1.0×10^{-4}M Na_2CO_3 solution. At what concentration of Ba^{2+} will a precipitate begin to form? (K_{xp} for for $BaCO_3 = 5.1 \times 10^{-9}$)

[2009]

Options:

A. 5.1×10^{-5}M

B. 8.1×10^{-8}M

C. 8.1×10^{-7}M

D. 4.1×10^{-5}M

Answer: A

Solution:

Solution:

$$\underset{1\times10^{-4}M}{Na_2CO_3} \longrightarrow \underset{1\times10^{-4}M}{2Na^{+}} + \underset{1\times10^{-4}M}{CO_3^{2-}}$$

$$K_{sp(BaCO_3)} = [Ba^{2+}][CO_3^{2-}]$$

$$[Ba^{2+}] = \frac{5.1\times10^{-9}}{1\times10^{-4}} = 5.1\times10^{-5}M$$

Question221

The equilibrium constants K_{p_1} and K_{p_2} for the reactions $X \rightleftharpoons 2Y$ and $Z \rightleftharpoons P+Q$, respectively are in the ratio of 1 : 9. If the degree of dissociation of X and Z be equal, then the ratio of total pressures at these equilibria is
[2008]

Options:

A. 1 : 36

B. 1 : 1

C. 1 : 3

D. 1 : 9

Answer: A

Solution:

Solution:
Let the initial moles of X be 'a' and that of Z be ' b ' then for the given reactions.

$$X \rightleftharpoons 2Y$$

Initial	a moles	0
At eqm. (moles)	$a(1-\alpha)$	$2a\alpha$

Total no. of moles $= a(1-\alpha)+2a\alpha$
$= a - a\alpha + 2a\alpha$
$= a(1+\alpha)$

$$K_{p_1} = \frac{(n_y)^2}{n_x} \times \left(\frac{P_{T_1}}{\sum n}\right)^{\Delta n}$$

Now $K_{p_2} = \frac{n_Q \times n_P}{n_z} \times \left[\frac{P_{T_2}}{\sum n}\right]^{dn}$

or $K_{p_2} = \frac{(b\alpha)(b\alpha)\cdot P_{T_2}}{[b(1-\alpha)][b(1+\alpha)]}$

or $\frac{K_{p_1}}{K_{p_2}} = \frac{4\alpha^2 \cdot P_{T_1}}{(1-\alpha^2)} \times \frac{(1-\alpha)^2}{P_{T_2}\cdot\alpha^2} = \frac{4P_{T_1}}{P_{T_2}}$

or $\frac{4P_{T_1}}{P_{T_2}} = \frac{1}{9}$ $\left[\because \frac{K_{P_1}}{K_{P_2}} = \frac{1}{9} \text{ given }\right]$

or $\frac{P_{T_1}}{P_{T_2}} = \frac{1}{36}$ or 1 : 36

Question222

For the following threereactions (i), (ii) and (iii), equilibrium constants are given:
(i) $CO(g) + H_2O(g) \rightleftharpoons CO_2(g) + H_2(g)$; K_1
(ii) $CH_4(g) + H_2O(g) \rightleftharpoons CO(g) + 3H_2(g)$; K_2
(iii) $CH_4(g) + 2H_2O(g) \rightleftharpoons CO_2(g) + 4H_2(g)$; K_3
Which of the following is correct?
[2008]

Options:

A. $K_1\sqrt{K_2} = K_3$

B. $K_2K_3 = K_1$

C. $K_3 = K_1K_2$

D. $K_3 \cdot K_2^3 = K_1^2$

Answer: C

Solution:

Solution:
Reaction (iii) can be obtained by adding reactions (i) and (ii) therefore $K_3 = K_1 . K_2$

Question223

Four species are listed below:
i. HCO_3^-
ii. H_3O^+
iii. HSO_4^-
iv. HSO_3F
Which one of the following is the correct sequence of their acid strength?
[2008]

Options:

A. iv < ii < iii < i

B. ii < iii < i < iv

C. i < iii < ii < iv

D. iii < 1 < iv < ii

Answer: C

Solution:

Solution:
The correct order of acidic strength of the given species is
$\underset{(iv)}{HSO_3F} > \underset{(ii)}{H_3O^+} > \underset{(iii)}{HSO_4^-} > \underset{(i)}{HCO_3^-}$
or (i) < (iii) < (ii) < (iv)

Question224

The pK_a of a weak acid, H A, is 4.80. The pK_b of a weak base, BOH , is 4.78. The pH of an aqueous solution of the corresponding salt, BA, will be [2008]

Options:

A. 9.58

B. 4.79

C. 7.01

D. 9.22

Answer: C

Solution:

Solution:
In aqueous solution, BA(salt) hydrolyses to give
$BA + H_2O \rightleftharpoons \underset{\text{base}}{BOH} + \underset{\text{acid}}{HA}$
Now pH is given by
$pH = \frac{1}{2}pK_W + \frac{1}{2}pK_a - \frac{1}{2}pK_b$
Substituting given values, we get
$pH = \frac{1}{2}(14 + 4.80 - 4.78) = 7.01$

Question225

The first and second dissociation constants of an acid H_2A are 1.0×10^{-5} and 5.0×10^{-10} respectively. The overall dissociation constant of the acid will be [2007]

Options:

A. 0.2×10^{5}

B. 5.0×10^{-5}

C. 5.0×10^{15}

D. 5.0×10^{-15}.

Answer: D

Solution:

Solution:

$H_2A \rightleftharpoons H^+ + HA^-$

$\therefore K_1 = 1.0 \times 10^{-5} = \frac{[H^+][HA^-]}{[H_2A]}$

$HA^- \rightleftharpoons H^+ + A^-$

$\therefore\ K_2 = 5.0 \times 10^{-10} = \frac{[H^+][A^-]}{[HA^-]}$

For the reaction,

$H_2A \rightleftharpoons 2H^+ + A^{2-}$

$K = \frac{[H^+]^2[A^{2-}]}{[H_2A]} = K_1 \times K_2$

$= (1.0 \times 10^{-5}) \times (5 \times 10^{-10}) = 5 \times 10^{-15}$

Question226

The pK_a of a weak acid (HA) is 4.5. The pOH of an aqueous buffer solution of HA in which 50% of the acid is ionized is [2007]

Options:

A. 7.0

B. 4.5

C. 2.5

D. 9.5

Answer: D

Solution:

Solution:

For acidic buffer, $pH = pK_a + \log\left[\frac{\text{salt}}{\text{acid}}\right]$ $pH = 4.5 + \log\frac{[\text{salt}]}{[\text{acid}]}$

As HA is 50% ionized so [salt] = [acid]

$\therefore pH = 4.5$

$\because pH + pOH = 14$

$pOH = 14 - pH = 14 - 4.5 = 9.5$

Question227

In a saturated solution of the sparingly soluble strong electrolyte $AgIO_3$ (molecular mass = 283) the equilibrium which sets in is $AgIO_3(s) \rightleftharpoons Ag^+(aq) + IO_3^-(aq)$. If the solubility product constant K_{sp} of $AgIO_3$ at a given temperature is 1.0×10^{-8}, what is the mass of $AgIO_3$ contained in 100mL of its saturated solution?
[2007]

Options:

A. 1.0×10^{-4}g

B. 28.3×10^{-2}g

C. 2.83×10^{-3}g

D. 1.0×10^{-7}g.

Answer: C

Solution:

Solution:
Let s = solubility $AgIO_3 \rightleftharpoons Ag^+ + IO_3^-$
$K_{sp} = [Ag^+][IO_3^s] = s \times s = s^2$
Given $K_{5p} = 1 \times 10^{-8}$
$\therefore \ s = \sqrt{K_{sp}} = \sqrt{1 \times 10^{-8}}$
$= 1.0 \times 10^{-4}$mol $/$ L $= 1.0 \times 10^{-4} \times 283g/$L
($\because$ Molecular mass of $AgIO_3 = 283$)
$= \frac{1.0 \times 10^{-4} \times 283 \times 100}{1000}g/$100mL
$= 2.83 \times 10^{-3}$g$/$100mL

Question228

Phosphorus pentachloride dissociates as follows, in a closed reaction vessel
$PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$
If total pressure at equilibrium of the reaction mixture is P and degree of dissociation of PCl_5 is x, the partial pressure of PCl_3 will be
[2006]

Options:

A. $\left(\frac{x}{x-1}\right)P$

B. $\left(\frac{x}{1-x}\right)P$

C. $\left(\frac{x}{x+1}\right)P$

D. $\left(\frac{2x}{1-x}\right)P$

Answer: C

Solution:

Solution:

$$\underset{1-x}{PCl_5(g)} \rightleftharpoons \underset{x}{PCl_3(g)} + \underset{x}{Cl_2(g)}$$

Total moles after dissociation $1-x+x+x=1+x$

p_{PCl_3} = Mole fraction of $PCl_3 \times$ Total pressure

$= \left(\frac{x}{1+x}\right)P$

Question229

The equilibrium constant for the reaction $SO_3(g) \rightleftharpoons SO_2(g) + \frac{1}{2}O_2(g)$ is $K_c = 4.9 \times 10^{-2}$. The value of K_c for the reaction $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g)$ will be [2006]

Options:

A. 9.8×10^{-2}

B. 4.9×10^{-2}

C. 416

D. 2.40×10^{-3}

Answer: C

Solution:

Solution:

$$SO_3(g) \rightleftharpoons SO_2(g) + \frac{1}{2}O_2(g)$$

$$K_c = \frac{[SO_2][O_2]^{1/2}}{[SO_3]} = 4.9 \times 10^{-2}$$

On taking the square of the above reaction

$$\frac{[SO_2]^2[O_2]}{[SO_3]^2} = 24.01 \times 10^{-4}$$

Now for $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3$

$$K_c' = \frac{[SO_3]^2}{[SO_2]^2[O_2]} = \frac{1}{24.01 \times 10^{-4}} = 416$$

Question230

For the reaction :
$2N\,O_2(g) \rightleftharpoons 2N\,O(g) + O_2(g)$
$(K_c = 1.8 \times 10^{-6}$ at $184^\circ C)(R = 0.0831 kJ/(mol\ .\ K))$
When K_p and K_c are compared at $184^\circ C$, it is found that
[2005]

Options:

A. Whether K_p is greater than, less than or equal to K_c depends upon the total gas pressure

B. $K_p = K_c$

C. K_p is less than K_c

D. K_p is greater than K_c

Answer: D

Solution:

Solution:
For the reaction:-
$2N\,O_2(g) \rightleftharpoons 2N\,O(g) + O_2(g)$
Given $K_c = 1.8 \times 10^{-6}$ at $184^\circ C$
$R = 0.0831 kJ/mol\ .\ K$
$K_p = K_c(RT)^{\Delta n}$
$K_p = 1.8 \times 10^{-6} \times 0.0831 \times 457$
$= 6.836 \times 10^{-6}$
Hence it is clear that $K_p > K_c$

Question231

An amount of solid $N\,H_4H\,S$ is placed in a flask already containing ammonia gas at a certain temperature and 0.50 atm pressure. Ammonium hydrogen sulphide decomposes to yield $N\,H_3$ and N_2S gases in the flask. When the decomposition reaction reaches equilibrium, the total pressure in the flask rises to 0.84 atm? The equilibrium constant for $N\,H_4H\,S$ decomposition at this temperature is
[2005]

Options:

A. 0.11

B. 0.17

C. 0.18

D. 0.30

Answer: A

Solution:

Solution:

$$NH_4HS(s) \rightleftharpoons NH_3(s) + H_2S(g)$$

At	$t = 0$	0.5 atm	0 atm
At eqm.		$0.5 + x$ atm	x atm.

Then $0.5 + x + x = 2x + 0.5 = 0.84$ (given)

$x = 0.17$ atm.

$p_{NH_3} = 0.5 + 0.17 = 0.67$atm

$p_{H_2S} = 0.17$atm

$K = p_{NH_3} \times p_{H_2S} = 0.67 \times 0.17\text{atm}^2$

$= 0.1139 = 0.11$

Question232

The exothermic formation of ClF_3 is represented by the equation:

$Cl_2(g) + 3F_2(g) \rightleftharpoons 2ClF_3(g); \quad \Delta H = -329kJ$

Which of the following will increase the quantity of ClF_3 in an equilibrium mixture of Cl_2, F_2 and ClF_3 ?

[2005]

Options:

A. Adding F_2

B. Increasing the volume of the container

C. Removing Cl_2

D. Increasing the temperature

Answer: A

Solution:

Solution:
The reaction given is an exothermic reaction thus accordingly to Le-Chatalier's principle lowering of temperature, addition of F_2 and or Cl_2 favour the forward direction and in hence the production of ClF_3.

Question233

Hydrogen ion concentration in mol / L in a solution of pH = 5.4 will be: [2005]

Options:

A. 3.98×10^{-6}

B. 3.68×10^{-6}

C. 3.88×10^{6}

D. 3.98×10^{8}

Answer: A

Solution:

Solution:
$\text{pH} = -\log[H^+] = \log \frac{1}{[H^+]}$
$5.4 = \log \frac{1}{[H^+]}$
On solving, $[H^+] = 3.98 \times 10^{-6}$

Question234

What is the conjugate base of OH^-? [2005]

Options:

A. O^{2-}

B. O^-

C. H_2O

D. O_2

Answer: A

Solution:

Solution:
Conjugate acid-base pair differ by only one proton.
$OH^- \longrightarrow H^+ + O^{2-}$
Conjugate base of OH is O^2

Question235

The solubility product of a salt having general formula MX_2, in water, is :4×10^{-12}. The concentration of M^{2+} ions in the aqueous solution of the salt is
[2005]

Options:

A. 4.0×10^{-10}M

B. 1.6×10^{-4}M

C. 1.0×10^{-4}M

D. 2.0×10^{-6}M

Answer: C

Solution:

Solution:
$MX_2 \rightleftharpoons \underset{s}{M^{2+}} + \underset{2s}{2s^{2+}}$
Where s is the solubility of MX_2
then $K_{sp} = 4s^3$
$4 \times 10^{-12} = 4s^3$
or $s = 1 \times 10^{-4}$

Question236

The equilibrium constant (K_c) for the reaction $N_2(g) + O_2(g) \rightarrow 2NO(g)$ at temperature T is 4×10^{-4}. The value of K_c for the reaction $NO(g) \rightarrow \frac{1}{2}N_2(g) + \frac{1}{2}O_2(g)$ at the same temperature is:
[2004, 2012]

Options:

A. 0.02

B. 2.5×10^2

C. 4×10^{-4}

D. 50.0

Answer: D

Solution:

Solution:
For the reaction
$N_2 + O_2 \rightarrow 2NO \quad Kc = 4 \times 10^{-4}$

Hence for the reaction
$NO \longrightarrow \frac{1}{2}N_2 + \frac{1}{2}O_2 \quad K_c' = \frac{1}{\sqrt{K_c}} = \frac{1}{\sqrt{4 \times 10^{-4}}} = 50$

Question237

What is the equilibrium expression for the reaction $P_4(s) + 5O_2(g) \rightleftharpoons P_4O_{10}(s)$?
[2004]

Options:

A. $K_c = [O_2]^5$

B. $K_c = [P_4O_{10}] / 5[P_4][O_2]$

C. $K_c = [P_4O_{10}] / [P_4][O_2]^5$

D. $K_c = 1 / [O_2]^5$

Answer: D

Solution:

Solution:
For $P_4(s) + 5O_2(g) \rightleftharpoons P_4O_{10}(s)$
$K_c = \frac{1}{(O_2)^5}$
Solids have concentration unity.

Question238

For the reaction, $CO(g) + Cl_2(g) \rightleftharpoons COCl_2(g)$ the K_p / K_c is equal to
[2004]

Options:

A. $\sqrt{RT}$

B. RT

C. $1 / RT$

D. 1.0

Answer: C

Solution:

Solution:

$K_p = K_c(RT)^{\Delta n}$;
Here $\Delta n = 1 - 2 = -1$
$\therefore \frac{K_p}{K_c} = \frac{1}{RT}$

Question239

The conjugate base of $H_2PO_4^-$ is
[2004]

Options:

A. H_3PO_4

B. P_2O_5

C. PO_4^{3-}

D. HPO_4^{2-}

Answer: D

Solution:

Solution:
$\underset{\text{Acid}}{H_2PO_4^-} \xrightarrow{-H^+} \underset{\text{Conjugate base}}{HPO_4^{2-}}$
Conjugate acid-base differs by H^+.

Question240

The molar solubility (in mol L^{-1}) of a sparingly soluble salt MX_4 is 's'. The corresponding solubility product is K_{sp° 's' is given in term of K_{sp} by the relation:
[2004]

Options:

A. $s = (256K_{sp})^{1/5}$

B. $s = (128K_{sp})^{1/4}$

C. $s = (K_{sp}/128)^{1/4}$

D. $s = (K_{sp}/256)^{1/5}$

Answer: D

Solution:

Solution:

$MX_4 \rightleftharpoons M_S^{4+} + \underset{4S^-}{4X}$

$K_{sp} = [s][4s]^4 = 256s^5$

$\therefore s = \left(\frac{K_{sp}}{256}\right)^{1/5}$

Question241

For the reaction equilibrium $N_2O_4(g) \rightleftharpoons 2NO_2(g)$ the concentrations of N_2O_4 and NO_2 at equilibrium are 4.8×10^{-2} and $1.2 \times 10^{-2} mol\ L^{-1}$ respectively. The value of K_c for the reaction is [2003]

Options:

A. $3 \times 10^{-1} mol\ L^{-1}$

B. $3 \times 10^{-3} mol\ L^{-1}$

C. $3 \times 10^{3} mol\ L^{-1}$

D. $3.3 \times 10^{2} mol\ L^{-1}$

Answer: B

Solution:

Solution:

$K_c = \frac{[NO_2]^2}{[N_2O_4]} = \frac{[1.2 \times 10^{-2}]^2}{[4.8 \times 10^{-2}]}$

$= 3 \times 10^{-3} mol/L$

Question242

Consider the reaction equilibrium $2SO_2(g) + O_2(g) \rightleftharpoons 2SO_3(g);\ \Delta H^\circ = -198kJ$ On the basis of Le Chatelier's principle, the condition favourable for the forward reaction is [2003]

Options:

A. increasing temperature as well as pressure

B. lowering the temperature and increasing the pressure

C. any value of temperature and pressure

D. lowering of temperature as well as pressure

Answer: B

Solution:

Solution:
Due to exothermic nature of reaction low or optimum temperature will be required. Since 3 moles are changing to 2 moles, therefore high pressure will be required.

Question243

Which one of the following statements is not true?
[2003]

Options:

A. $pH + pOH = 14$ for all aqueous solutions

B. The pH of $1 \times 10^{8} M$ HCl is 8

C. 96,500 coulombs of electricity when passed through a $CuSO_4$ solution deposits 1 gram equivalent of copper at the cathode

D. The conjugate base of $H_2PO_4^-$ is HPO_4^{2-}

Answer: B

Solution:

Solution:
pH of an acidic solution should be less than 7 . The reason is that from H_2O, $[H^+] = 10^{-7} M$ which cannot be neglected in comparison to $10^{-8} M$. The pH can be calculated as.
From acid, $[H^+] = 10^{-8} M$
From H_2O, $[H^+] = 10^{-7} M$
$\therefore$ Total $[H^+] = 10^{-8} + 10^{-7}$
$= 10^{-8}(1 + 10) = 11 \times 10^{-8}$
$\therefore pH = -\log[H^+] = -\log 11 \times 10^{-8}$
$= -[\log 11 + 8 \log 10]$
$= -[1.0414 - 8] = 6.9586$

Question244

When rain is accompanied by a thunderstorm, the collected rain water will have a pH value
[2003]

Options:

A. slightly higher than that when the thunderstorm is not there

B. uninfluenced by occurrence of thunderstorm

C. that depends on the amount of dust in air

D. slightly lower than that of rain water without thunderstorm.

Answer: D

Solution:

Solution:
The rain water after thunderstorm contains dissolved acid and therefore the pH is less than rain water without thunderstorm.

Question245

The solubility in water of a sparingly soluble salt AB_2 is 1.0×10^{-5}mol L^{-1}. Its solubility product will be
[2003]

Options:

A. 4×10^{-10}

B. 1×10^{-15}

C. 1×10^{-10}

D. 4×10^{-15}

Answer: D

Solution:

Solution:
$AB_2 \rightleftharpoons A^{2+} + 2B^-$
$[A] = 1.0 \times 10^{-5}$, $[B] = [2.0 \times 10^{-5}]$
$K_{gp} = [B][A] = [2 \times 10^{-5}]^2[1.0 \times 10^{-5}] = 4 \times 10^{-15}$

Question246

For the reaction $CO(g) + (1/2)O_2(g) \rightleftharpoons CO_2(g)$, K_p/K_c is
[2002]

Options:

A. RT

B. $(RT)^{-1}$

C. $(RT)^{-1/2}$

D. $(RT)^{1/2}$

Answer: C

Solution:

Solution:
$K_p = K_c(RT)^{\Delta n}$;

$\Delta n = 1 - \left(1 + \frac{1}{2}\right) = 1 - \frac{3}{2} = -\frac{1}{2}$

$\therefore \frac{K_p}{K_c} = (RT)^{-1/2}$

Question247

Change in volume of the system does not alter which of the following equilibria? [2002]

Options:

A. $N_2(g) + O_2(g) \rightleftharpoons 2NO(g)$

B. $PCl_5(g) \rightleftharpoons PCl_3(g) + Cl_2(g)$

C. $N_2(g) + 3H_2(g) \rightleftharpoons 2NH_3(g)$

D. $SO_2Cl_2(g) \rightleftharpoons SO_2(g) + Cl_2(g)$

Answer: A

Solution:

Solution:
In reaction (a) the ratio of number of moles of reactants to products is same i.e. $2:2$, hence change in volume will not alter the number of moles.

Question248

Species acting as both Bronsted acid and base is [2002]

Options:

A. HSO_4^-

B. Na_2CO_3

C. NH_3

D. OH

Answer: A

Solution:

Solution:
$(HSO_4)^-$ can accept and donate a proton
$(HSO_4)^- + H^+ \rightarrow H_2SO_4$ (acting as base)
$(HSO_4) - H^+ \rightarrow SO_4^{2-}$. (acting as acid)

Question249

Let the solubility of an aqueous solution of $Mg(OH)_2$, be x, then its K_{sp} is [2002]

Options:

A. $4x^3$

B. $108x^5$

C. $27x^4$

D. 9x.

Answer: A

Solution:

Solution:
$Mg(OH)_2 \rightarrow [\underset{x}{Mg^{2+}}] + 2[\underset{2x}{OH^-}]$
$K_{sp} = [Mg][OH]^2 = [x][2x]^2 = x.4x^2 = 4x^3$

Question250

1M NaCl and 1M HCl are present in an aqueous solution. The solution is [2002]

Options:

A. not a buffer solution with pH < 7

B. not a buffer solution with pH > 7

C. a buffer solution with pH < 7

D. a buffer solution with pH > 7.

Answer: A

Solution:

Solution:

A buffer is a solution of weak acid and its salt with strong base and vice versa. HCl is strong acid and $NaCl$ is its salt with strong base. pH is less than 7 due to HCl.
