

The Arithmetic of Order: A Unified Origin for the Standard Model's Gauge Symmetries, Hypercomplex Numbers, and Codewords

Faysal El Khettabi
Ensemble AIs
faysal.el.khettabi@gmail.com

July 15, 2025

Abstract

This report details the Arithmetic of Order (AoO) as a discrete arithmetic framework that provides a unified origin for the Standard Model's gauge symmetries, hypercomplex numbers, and optimal error-correcting codes. The AoO posits that physical and mathematical structures emerge from the ordered combinatorial progression $1 \rightarrow n \rightarrow n + 1$ and its associated powerset hierarchy, $\mathcal{P}(\Omega_n)$, without assuming a pre-existing continuum. We show how this framework, realized through Farey sequences under the modular group $SL(2, \mathbb{Z})$, lifts naturally to its universal central extension, the braid group B_3 . This topological extension provides a direct mechanism where the three Reidemeister moves generate the gauge groups $U(1)$, $SU(2)$, and $SU(3)$. The discussion section contextualizes these results, highlighting the framework's power in unifying the language of physical transformations (hypercomplex numbers) with principles of information stability (codewords), and notes the convergence of our conclusions with parallel topological models of physics.

Keywords: gauge symmetry, Reidemeister moves, braid group, arithmetic foundations, hypercomplex numbers, Golay code, $SL(2, \mathbb{Z})$, Standard Model, Veldkamp Space

1 Introduction

A primary goal of fundamental physics is to understand the logical origins of the laws of nature. The Standard Model of particle physics is remarkably successful, yet it does not explain the genesis of its foundational gauge symmetries: $U(1)$, $SU(2)$, and $SU(3)$. This work presents the Arithmetic of Order (AoO) as a candidate for this explanation.

The AoO challenges the traditional reliance on continuum mathematics, which posits that finite physical systems require infinite constructs for their description [3]. Instead, the AoO proposes that the structure of physics is an inevitable consequence of a simple, finitistic, and constructive process based on ordered distinctions. This paper outlines the results of this framework, showing how its core principles give rise to the topology of gauge interactions, and discusses the broader implications of its unifying power.

2 Core Principles of the Discrete Arithmetic Framework

2.1 Powerset Combinatorics and Finite Fields

The AoO begins with an ordered set of n distinguishable degrees of freedom, $\Omega_n = \{1, 2, \dots, n\}$. The full space of all possible system configurations is its powerset, $\mathcal{P}(\Omega_n)$, which contains 2^n unique subsets [2]. Each subset corresponds to a specific configuration, represented empirically by a characteristic function ($\chi_S : \Omega_n \rightarrow \{0, 1\}$).

This powerset, when equipped with the symmetric difference operation (Δ), is isomorphic to the n -dimensional vector space over the finite field of two elements, $\mathbb{F}_2^n$. This provides a native algebraic language for system states based on binary logic (bitwise XOR).

2.2 Emergence of Hypercomplex Numbers and Codewords

From this single powerset engine, two critical structures emerge deterministically:

- **Hypercomplex Numbers:** The 2^n elements of $\mathcal{P}(\Omega_n)$ directly correspond to the 2^n basis units of a hypercomplex algebra [2]. For $n = 3$, the $2^3 = 8$ elements of the powerset $\mathcal{P}(\Omega_3)$ provide a natural basis for the octonions $\{1, e_1, \dots, e_7\}$ [8]. The multiplication rules are defined by the bitwise XOR operation on the binary representations of the basis units.
- **Optimal Codewords:** When specific, finite constraints (e.g., linearity, self-duality) are applied as a "sieve" to the powerset, exceptionally stable structures emerge. The premier example at $n=24$ is the extended binary Golay code (G_{24}), a structure known for its unique error-correcting properties and its deep connection to other fundamental objects like the Leech Lattice [6, 3].

In summary, the structured powerset $\mathcal{P}(\Omega_n)$ not only provides the state space of discrete systems, but also underlies the emergence of algebraic and informational tools previously seen as unrelated—namely, hypercomplex units and optimal codes.

3 Results: From Arithmetic to Gauge Theory

3.1 Arithmetic Realization: Farey Sequence and $SL(2, \mathbb{Z})$

The AoO finds a concrete realization in the Farey sequence hierarchy, $\mathcal{F}_n$. The refinement of the sequence via the mediant operation, where the mediant of $\frac{a}{b}$ and $\frac{c}{d}$ is defined as $\frac{a+c}{b+d}$, is a construction governed coherently by the modular group $SL(2, \mathbb{Z})$ [4]. This establishes a direct link between the arithmetic progression of the AoO and the algebraic symmetries of the modular group.

3.2 Topological Extension: The Braid Group B_3

The framework's key topological step relies on a well-established mathematical fact: the braid group on 3 strands, B_3 , is the universal central extension of the modular group $PSL(2, \mathbb{Z})$ [7]. This provides a rigorous, non-arbitrary pathway to lift the arithmetic process into the topological domain of braids. The braid relation, $\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2$, is precisely the third Reidemeister move.

3.3 Finite Geometries and Veldkamp Spaces

The powerset $\mathcal{P}(\Omega_n)$, under symmetric difference, forms the finite projective geometry $PG(n-1, 2)$. Its point-line incidence structure naturally encodes the combinatorial logic underlying the AoO.

Formally, a *configuration* $\mathcal{C} = (\mathcal{P}, \mathcal{L}, I)$ consists of a finite set of points $\mathcal{P}$, a finite set of lines $\mathcal{L}$, and an incidence relation $I \subseteq \mathcal{P} \times \mathcal{L}$ indicating which points lie on which lines.

A *geometric hyperplane* of $\mathcal{C}$ is a proper subset $H \subsetneq \mathcal{P}$ such that any line of $\mathcal{L}$ is either entirely contained in H or intersects H in exactly one point.

If $\mathcal{C}$ possesses geometric hyperplanes, its *Veldkamp space* $\mathcal{V}(\mathcal{C})$ is defined as follows [5]:

- Each point of $\mathcal{V}(\mathcal{C})$ is a geometric hyperplane of $\mathcal{C}$.
- Each line of $\mathcal{V}(\mathcal{C})$ is the triple $\{H', H'', \overline{H' \Delta H''}\}$, where H' and H'' are distinct hyperplanes, Δ denotes symmetric difference, and the overline denotes the complement in $\mathcal{P}$.

This symmetric difference–complement rule mirrors the braid relation of B_3 : the minimal nontrivial closure occurs when distinctions braid into a stable trio, paralleling the Reidemeister III slide relation $\sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2$. Thus, the AoO's combinatorial hierarchy naturally generates the hyperplane logic that encodes topological phase coherence and the emergence of the Standard Model gauge groups.

3.4 Physical Interpretation: Reidemeister Moves and Gauge Groups

Following the interpretation advanced by Christoph Schiller [1], the three Reidemeister moves correspond directly to the gauge interactions. They represent the fundamental ways to deform a topological structure while preserving its essential identity. The mapping is as follows:

- **Twist (Move I)** generates the $U(1)$ group of electromagnetism.

- **Poke (Move II)** generates the $\mathbf{SU}(2)$ group of the weak interaction.
- **Slide (Move III)** generates the $\mathbf{SU}(3)$ group of the strong interaction.

In summary, this section establishes a formal chain of logic: the AoO's discrete arithmetic, realized via Farey sequences, is governed by $SL(2, \mathbb{Z})$, which lifts naturally to the Braid Group B_3 , whose combinatorial and topological relations in turn map directly to the gauge groups of the Standard Model.

4 Discussion

4.1 A Unified Origin for Symmetries and Stability

The ability of the AoO to derive the language of physical transformations (hypercomplex numbers) and the principles of information stability (optimal codes) from the same combinatorial foundation is a powerful indicator of its potential as a unifying theory. It suggests that the geometric symmetries of physics and the rules ensuring the existence of stable, coherent structures are not independent features of our universe, but are two facets of the same underlying arithmetic of order [3].

4.2 Convergence with Topological Models

The conclusion that the three Reidemeister moves generate the three gauge groups is not unique to this framework. Schiller's "Strand Tangle Model" arrives at the same result from a different foundation—one based on physical postulates about unobservable Planck-scale strands [1]. This convergence from two distinct starting points (AoO's abstract arithmetic vs. Schiller's physical strands) provides powerful, independent corroboration for the idea itself. It suggests that the relationship between topology and gauge physics is a deep and fundamental one, worthy of intense investigation.

4.3 Limitations and Open Questions

While the AoO framework suggests compelling origins for gauge symmetries, it remains to be shown whether the full Lagrangian dynamics of the Standard Model—including the Higgs mechanism and fermion masses—can be derived within this framework. Additionally, the physical context for the lifting from $SL(2, \mathbb{Z})$ to B_3 must be further formalized. These questions provide fertile ground for future research.

5 Conclusion

The Arithmetic of Order demonstrates that the gauge hierarchy of the Standard Model need not be an arbitrary postulate, but can be seen as the inevitable result of a finitistic, arithmetic process that is lifted to its minimal topological extension. By unifying the origins of geometric symmetries, information-theoretic stability, and combinatorial geometry, and by converging with conclusions from independent physical models, the AoO presents a compelling, testable, and potentially revolutionary foundation for understanding the logical structure of our universe.

A Appendix: Farey Sequence Generation

The Farey sequence $\mathcal{F}_n$ is generated recursively. To obtain $\mathcal{F}_{n+1}$ from $\mathcal{F}_n$, one inserts the median $\frac{a+c}{b+d}$ between any two adjacent fractions $\frac{a}{b}$ and $\frac{c}{d}$ in $\mathcal{F}_n$ for which $b+d = n+1$.

Example for $n = 3$:

- $\mathcal{F}_1 = \{\frac{0}{1}, \frac{1}{1}\}$
- $\mathcal{F}_2 = \{\frac{0}{1}, \frac{1}{2}, \frac{1}{1}\}$ (insert $\frac{0+1}{1+1} = \frac{1}{2}$)
- $\mathcal{F}_3 = \{\frac{0}{1}, \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, \frac{1}{1}\}$ (insert $\frac{0+1}{1+2} = \frac{1}{3}$ and $\frac{1+1}{2+1} = \frac{2}{3}$)

References

- [1] C. Schiller, “On the relation between the three Reidemeister moves and the three gauge groups,” arXiv:2312.14173v1 [physics.gen-ph] (Dec 2023).
- [2] Faysal El Khettabi, “A Comprehensive Modern Mathematical Foundation for Hypercomplex Numbers with Recollection of Sir William Rowan Hamilton, John T. Graves, and Arthur Cayley,” viXra:2505.0054 (May 2025).
- [3] Faysal El Khettabi, “The Arithmetic of Order: A Finitistic Foundation for Mathematics, Emergent Structures, and Intelligent Systems,” viXra:2505.0059 (May 2025).
- [4] Faysal El Khettabi, “Physics as Quantized Measurement: The Farey Sequence as a Realization of the Arithmetic of Order,” viXra:2507.0068 (July 2025).
- [5] M. Saniga et al., “Cayley–Dickson Algebras and Finite Geometry,” arXiv:1405.6888.
- [6] J. H. Conway and N. J. A. Sloane, “Sphere Packings, Lattices and Groups,” 3rd ed., Springer, 1998.
- [7] L. H. Kauffman, “Knots and Physics,” 4th ed., World Scientific, 2013.
- [8] J. Baez, “The Octonions,” Bull. Amer. Math. Soc., vol. 39, no. 2, pp. 145-205, 2002, arXiv:math/0105155.