






# Draft information

version: draft-0.0

20xx

## Requests to coauthors

- Should I include more networks? Easy to get some more from other published studies. Could improve the manuscript but every new data set has to be cleaned independently. This could take around 2 days per network. 
- Should I increase the resolution of the bioclimatic variables? Currently working with 2.5' (~5km). Increasing to 30" (~1km) might improve the analysis but we're talking about working with ~25GB+ of climatic data instead of the 1GB used now. Unsure the benefits would outweigh the costs. 
- Audrey: Could you please fill in the gaps in the rationale behind filtering out occurrences with potential low quality coordinates (page ??, line ??)
- I chose, rather arbitrarily, a mean absolute error of 10% for the sensitivity analysis. Does this choice sound appropriate to you? 
- Should I remove the species overall generalism and the size of the grinellian niche from the model to simplify things. Both contribute little to the model and the story.  For the former, seems to be redundant once the number of possible interacting species has been included in the model. For the later, there is little biological reasons (what I could gather from the lit.) to justify their inclusion in the model. What you think? If not, at the very least I think I would move panels a and c in Fig. 5 to the supplementary info.
- What you think would be the best way to end the paper? Thinking about what could be highlighted in the conclusion. 

## Done

- Downloaded network data from the Web of Life and cleaned it. I didn't anticipated data needed SO MUCH work! I estimated three days to do this but took roughly 3 weeks to do it properly and there is still some work to do. Quite a mess...

- Downloaded bioclimatic and habitat data
- Downloaded species occurrences. Wasn't straightforward as there is a lot of species and lots of restrictions with the GBIF API. About a week getting this to work.
- Calculated environmental niche size and the environmental suitability of species in their communities
- Modelled the normalised degree of species as a proxy of eltonian niche size using a bayesian framework

## To-do

- Manually fix the remaining species names that are incorrect and were not picked up by the automatic procedure. Doing 5-10 per day, so should be done in a couple months
- Fix references. There are lots of mistakes re. author, years and publication venue particularly for those from which data was downloaded.
- Finish off discussion with a punchy conclusion.
- Fix the WAIC table. Currently shows the model without guild as unknown.
- Add Lupe's feedback to the intro. Need to better define stress and Eltonian/Grinellian niche and don't need to say realised Eltonian niche when I really just mean number of partners

# Environmental stress affects niche breadth in plant-pollinator communities

*E. Fernando Cagua*<sup>1</sup> (*efc29@uclive.ac.nz*)

*Audrey Lustig*<sup>2</sup> (*audrey.lustig@canterbury.ac.nz*)

*Jason M. Tylianakis*<sup>1</sup> (*jason.tylianakis@canterbury.ac.nz*)

*Daniel B. Stouffer*<sup>1</sup> (*daniel.stouffer@canterbury.ac.nz*)

<sup>1</sup> Centre for Integrative Ecology, School of Biological Sciences, University of

Canterbury, Private Bag 4800, Christchurch 8041, New Zealand

<sup>2</sup> Geospatial Research Institute, University of Canterbury, Private Bag 4800,

Christchurch 8041, New Zealand

**Running title:** XX

**Keywords:** XX

**Type of article:** XX

**Number of words:** 151 in abstract; 5,025 in main text.

**Number of displays:** 6 figures; 1 tables; 0 text boxes.

**Number of references:** 86

**Author for correspondence:** E. Fernando Cagua (+64 20 4026 8153).

**Data accessibility:** Data supporting the results will be accessible in an appropriate data repository after publication. The DOI will be included here.

**Author contributions:** XX


## 16 Abstract

17 Evidence that the environment influences the interaction between species  
18 is rapidly accumulating. However, how it happens is currently unclear as  
19 environmental gradients appear to have contrasting or non-linear effects on  
20 the species' trophic niche breadth depending on the environmental variable.  
21 Here, we explore the relationship between the stresses imposed by the envi-  
22 ronment, instead of environmental gradients directly, and niche breadth using  
23 a global dataset of plant-pollinator interactions. We found that environmental  
24 stress plays a significant role in determining the number of partners a species  
25 interacts with, but this role is highly variable across species. In particular,  
26 when faced with environmental stress, species that have a large number of  
27 interactions are more likely to focus on a smaller number of, presumably  
28 higher-quality, interactions. Contrastingly, the specialists that can cope  
29 with increased stress are more likely to broaden their niche and engage in  
30 opportunistic interactions, effectively behaving as facultative generalists.

## 31 Introduction

32 Species interactions are known to vary widely across space and time. There  
33 are multiple examples of species that interact tightly in an ecological com-  
34 munity or a particular season but not in another. It is crucial to understand  
35 why and how this turnover of interactions occur because the network of  
36 interactions of an ecological community is known to underpin ecosystem  
37 function and stability. Despite much progress in the last decades, we are  
38 still unable to predict species interactions, partially because we still have not  
39 accurately identified the factors that drive species to interact under certain  
40 conditions and not under another.






41 There are two main processes involved in whether two species in a community  
42 interact or not. The first is an “Eltonian” niche process that is underpinned  
43 by the traits of the interacting species. Compared to other kinds of trophic  
44 interactions, mutualistic interactions, and plant-pollination interactions,  
45 in particular, are relatively generalised. While a large proportion of the  
46 interactions appear to be opportunistic, a fraction of interactions shows a  
47 strong signature of deep co-evolutionary history (Hutchinson, Cagua, and   
48 Stouffer 2017). In contrast to opportunistic interactions with low trait-  
49 matching, the more co-evolved interactions with optimally matched traits are  
50 more persistent and presumably lead to a more substantial per-capita benefit  
51 for the interacting species. The second is a neutral process underpinned by  
52 the encounter rate of the interacting species. All else being equal, when the  
53 encounter rate of interacting species is high, there is a higher probability to  
54 interact than if the rate were lower, for example, due to species rarity.

55 Perhaps, because it operates at scales larger than those of the Eltonian niche  
56 concept, the abiotic environment is one of the factors that is commonly  
57 overlooked by community ecologists when explaining interaction turnover  
58 (but see Gravel et al. 2018). The most evident—and best-studied—way


59 through which the environment can affect the structure of ecological networks  
60 is by way of patterns of community composition. At the heart of biogeography  
61 is the idea that the abiotic environment can determine the “Grinnellian” niche  
62 of a species, this is whether the species can have a positive growth-rate when  
63 rare in a particular location. For two species to interact, they first need to  
64 co-occur in the same community, which can only happen if the environmental  
65 conditions of the community fall within the Grinnellian niche of both species.

66 Although less understood, the environment can also affect the interactions  
67 themselves, after community composition has been established (Tylianakis  
68 and Morris 2017). For instance, some studies show that trophic interactions,  
69 like predation (McKinnon et al. 2010; Vucic-Pestic et al. 2011) and herbivory  
70 (Baskett and Schemske 2018), can **increase** with temperature but might  
71 decrease with precipitation (Pires et al. 2016), and both plant-plant (Bertness  
72 and Callaway 1994) and plant-pollinator (Tur et al. 2016) interactions  
73 can **switch from facilitative to competitive along an elevation/temperature**  
74 gradient. Contrastingly, other studies have shown either no effect or non-  
75 linear effects of temperature or precipitation on plant-pollinator interactions  
76 (Devoto, Medan, and Montaldo 2005; Gravel et al. 2018).

 77 One way to summarise the seemingly contradictory evidence of whether the  
78 environment can influence whether two species interact or not is by moving  
79 from gradients of temperature or precipitation to gradients of environmental  
80 stress. Previous research is so far equivocal  taken together, suggest that  
81 environmental stress **can have two contrasting effects on species'** Eltonian-  
82 niche (Tylianakis and Morris 2017). On the one hand, it is possible that   
83 when species are under environmental stress, they might be “pressured” to  
84 focus on partners with which they are best adapted to interact. For instance,  
85 Hoiss et al. (2012) found increased phylogenetic clustering between plants  
86 and pollinators at higher altitudes; while Peralta et al. (2015) found that  
87 parasitoids in plantation forest, where environmental stress was higher than

88 in native forests, were constrained to interact with hosts they were best  
89 adapted to attack. Similarly, Lavandero and Tylianakis (2013) found that  
90 environmental stress due to higher temperature reduced the breadth of the  
91 Eltonian niche of parasitoids.

92 On the other hand, it is also possible that when species are under envi-  
93 ronmental stress, they are forced to be more flexible in their interactions  
94 as higher environmental stress is likely to be reflected in greater energetic  
95 or reproductive costs and therefore they might not be able to sustain en-  
96 counter rates with their preferred partners at sufficient levels. In line with  
97 this hypothesis, Hoiss, Krauss, and Steffan-Dewenter (2015) found that the  
98 specialisation of plant-pollinator networks decreased both with elevation  
99 and after extreme drought events. Likewise, Pellissier et al. (2010) found a  
100 positive relationship between niche breadth and environmental stress: disk-  
101 or bowl-shaped blossoms (which allow a large number of potential pollinator  
102 species to access pollen and nectar rewards) dominated at high altitude  
103 flower communities.

104  We analyse data on plant-pollinator interactions globally and calculate the  
105 environmental suitability of all species present at two or more communities  
106 ~~to~~. Our main aim is to leverage that information to test the two competing  
107 hypotheses that relate environmental stress and Eltonian niche breadth  
108 and whether there is a pattern within and across species. Specialist species  
109 can become “facultative” generalists to reduce their vulnerability to the  
110 absence of preferred partners (for example, when variations in climate de-  
111 couple phenologies; Benadi et al. 2014). We therefore also expect that  
112 as environmental stress increases species with a relatively small number of  
113 partners are more likely to broaden their Eltonian niche. Species with a  
114 large number of partners, on the other hand, should have a larger pool of  
115 available partners and might, therefore, be more likely to narrow their niche  
116 under environmental stress. Importantly, when testing these hypotheses, we

control for the potential effects of the environment in community composition and the size of the species fundamental niche, both from an Eltonian (interactions) and Grinnellian (environment) perspective.

## Methods

We retrieved plant-pollinator networks from the Web of Life database (Fortuna, Ortega, and Bascompte 2014). This database contains datasets originating from 57 studies published in the primary literature between 106 and 2016 (Kaiser-Bunbury et al. 2010, 2014; Bartomeus, Vilà, and Santamaría 2008; Arroyo, Primack, and Armesto 1982; Hattersley-Smith 1985; Barrett and Helenurm 1987; Dicks, Corbet, and Pywell 2002; Dupont, Hansen, and Olesen 2003; Elberling and Olesen 1999; Herrera 1988; Hocking 1968; Inouye and Pyke 1988; Kato et al. 1990; Medan et al. 2002; Memmott 1999; Motten 1986; Olesen, Eskildsen, and Venkatasamy 2002; Ollerton 2003; Petanidou and Vokou 1993; McMullen 106AD; Mosquin 1967; Clements and Long 1923; Kevan 1970; Small 1976; Montero 2005; Stald, Valido, and Olesen 2003; Ingversen 2006; Bundgaard 2003; Bek 2006; KATo 2000; Dupont and Olesen 2009; Lundgren and Olesen 2005; Percival 1974; Philipp et al. 2006; Primack 1983; Ramirez 1989; Ramirez and Brito 1992; Schemske et al. 1978; Smith-Ramírez et al. 2005; Vázquez 2002; Robertson 1929; Ferreira Canela 2006; Kohler 2011; Lasprilla and others 2003; Sabatino 2010; Lara 2006; Las-Casas, Azevedo Júnior, and Dias Filho 2012; del Coro Arizmendi and Ornelas 1990; Abreu and Vieira 2004; Vizentin-Bugoni et al. 2016; Bezerra, Machado, and Mello 2009; INouE et al. 1990; KATo, Matsumoto, and Kato 1993; Yamazaki and Kato 2003; Gutierrez Z, Rojas-Nossa, and Stiles 2004; Kato and Miura 1993; Kakutani et al. 1990). Calculating the environmental stress of species in their community and their Eltonian niche breadth requires that the scientific names used to identify them are valid and unambiguous. This is because valid scientific names are necessary to



145 both accurately match species across studies and to obtain other information  
146 from biological databases. However, a significant proportion of the data  
147 obtained contain spelling mistakes, taxonomic synonyms, or non-standard  
148 abbreviations. Therefore, our first step was to validate and correct scientific  
149 names across studies to ensure that we matching of species across studies  
150 was as accurate as possible.

151 Second, we calculated the suitability of the environment for a species in a  
152 particular community as a proxy of environmental stress. We assume that the  
153 environmental stress a species experience in a particular location is inversely  
154 related to the suitability of the average environmental conditions in that  
155 place. Our third and final step was to relate the environmental suitability  
156 to the relative number of partners a species has in a community, as a proxy  
157 for Eltonian niche breadth. To explore this relationship within and across  
158 species, we used a multilevel Bayesian model in which we controlled for the  
159 potential effects of the environment on co-occurrence.

## 160 **Plant-pollinator interactions**

161 All together, interaction data from the included studies included 11,231  
162 unique organism names. From these 1,166 were present in more than one  
163 study. From the total number of organisms, 159 were identified to the  
164 subspecies or variety level, 6,759 to the species level, 1,755 to the genus level,  
165 whereas the remaining 2,558 were unidentified. As the species level was the  
166 most common taxonomic rank available in our interaction datasets, in all  
167 further analysis, we grouped together subspecies or varieties within the same  
168 species.

169 We were able to confirm the validity of 5,263 of the scientific names used  
170 to identify organisms (roughly 76%). We assessed the validity of a name by  
171 querying the Global Names Resolver database (<https://resolver.globalnames>).

org) which includes data from 98 taxonomic sources. We accessed this database using the function `gnr_resolve` from the R package `taxize` 0.9.6 (Chamberlain and Szocs 2013; S. Chamberlain, Szoezs, et al. 2019).

From the remaining 1,655 names we were unable to validate, we were able to identify and correct 726 that contained spelling mistakes. These spelling mistakes were corrected automatically by fuzzy matching the canonical names in our data sources with those in the Global Names Resolver database. However, on rare occasions, the fuzzy matching algorithm can suggest a scientific name that has a similar spelling, but that corresponds to an organism in a different taxonomic group, often a separate kingdom. To address this potential problem, we checked the taxonomic hierarchy of suggested names and confirmed that it matched our expected taxon. We retrieved all taxonomic hierarchies from the National Center for Biotechnology Information taxonomic database (<https://www.ncbi.nlm.nih.gov/taxonomy>).

As species names are constantly changing, we subsequently checked for possible synonyms of the canonical names in our data sources. Using data from the Integrated Taxonomic Information System database (<http://www.itis.gov>), we found synonyms and alternative names for 611 species.

Finding these alternative names was required for two main reasons. First, because we wanted to be able to identify the cases in which the same species might have been recorded with different names in various data sources. This can occur not only when the canonical name has been changed but also when there are widely used orthographic variants. Second, because retrieving occurrence data is often only possible using the latest accepted/valid name for a particular species.

All together, from the 1,655 names we were unable to validate, it was not possible to automatically correct or find synonyms 332 of them. We then manually consulted multiple online databases, chiefly Wikispecies (<https://species.wikimedia.org/>), and looked for canonical names that both,




Figure 1: Worldwide distribution of pollination communities included in this study

resembled the unvalidated names and matched the geographic and taxonomic expectations. In this fashion, we were able to further correct 25 names. Most manual corrections were made on names that have been abbreviated or had more than two spelling mistakes. A complete list of manual name corrections can be seen in Table S1.

This cleaning process allowed us to match further 270 names across data sources and, by doing so, identify another 72 species that were present in more than one study. The process also allowed us to identify problematic data sources in which some names were included as both plants and pollinators. These data sources were removed from further analysis. In seven of our data sources interaction data was recorded at multiple points in time. When this was the case, we combined interaction data into one single interaction network.

The cleaning process resulted on a total of 2,555 plants and 8,406 pollinator species distributed across 73 locations around the globe (Figure 1, 2).

## Species suitability

Our next step was to determine the habitat suitability of the species as a proxy of the environmental stress they experience in their community. As we aim to compare the trophic niche for different suitability levels, we only do this for species that were present in at least two communities. To calculate




Figure 2: Frequency distribution of the number of locations in which a species is present. The most common pollinator species was *Apis mellifera*, which was sampled on 42 locations, while the most common plant species was *Trifolium repens*, which was sampled on 11 locations.

the suitability of a species in a particular location, we used a (Grinnellian) niche-factor analysis (Hirzel et al. 2002; Broennimann et al. 2012). This approach is based on the probability density function of species distribution in an environmental variable space. In a nutshell, habitats (characterised by a collection of environmental variables) in which the species occurs most often are deemed to be suitable for the species than habitats in which the species has never been observed. This approach to estimating the habitat suitability requires two critical pieces of information. First, we require information about the occurrences of the species of interest. Second, we require information about the environmental conditions for all the locations in which the species occurs.

We retrieved occurrences from the Global Biodiversity Information Facility (GBIF; <https://www.gbif.org>) using the R package `rgbif` 0.9.6 (Chamberlain and Boettiger 2017; S. Chamberlain, Barve, et al. 2019). Specifically, for each species, we only requested occurrences for which the coordinates of the observation were available and that had no known geospatial issue in the GBIF database. Roughly, we downloaded 38.1 million occurrences for the 986 species we were interested on. These occurrences, however, contain observations of mixed quality. Therefore, we followed (ref?) and applied a

240 series of simple filters and geographic heuristics to remove those of lower  
 241 quality. Specifically, we removed all occurrences with (i) a coordinate  
 242 uncertainty larger than 100km; (ii) those recorded prior to 1945 (ref? o  
 243 why?); (iii) those in which the number of individuals registered was either  
 244 zero or more than 100 as such large counts are often erroneous (ref? or  
 245 why?); and (iv) those occurrences in which the “basis of record” was not a  
 246 human observation or a preserved specimen (ref? why?). We then used the  
 247 R package **CoordinateCleaner** 0.9.6 (Zizka et al. 2019) and land mass and  
 248 country data from Natural Earth (<https://www.naturalearthdata.com>) with  
 249 a 1:10,000,000 scale to further identify and remove problematic occurrences.  
 250 We removed occurrences for which their coordinates (v) fell outside the  
 251 borders of the country where they were recorded; (vi) those around a country  
 252 capital or the centroid of the country and province centroids; (vii) those  
 253 around a biodiversity institution; and (viii) those located within oceans.  
 254 Thorough this cleaning process, we removed with 7.2 million occurrences  
 255 distributed accross 916 species. Finally, we integrated the occurrences from  
 256 our plant-pollinator communities to the cleaned occurrences retrieved from  
 257 GBIF.

258 We retrieved environmental data from WorldClim V2.0, which includes 19  
 259 bioclimatic variables commonly used in species distribution modelling (Fick  
 260 and Hijmans 2017). We then complemented data obtained from WorldClim  
 261 with data from Envirem (Title and Bemmels 2017), which includes 16 extra  
 262 bioclimatic and two topographic variables. The additional set of variables  
 263 from Envirem are relevant to ecological or physiological processes and as  
 264 such, have the potential to improve our suitability estimation (Title and  
 265 Bemmels 2018). We obtained all environmental data as rasters composed  
 266 by cells of 2.5 arc-minutes. We chose this resolution because it provides  
 267 a reasonable match to the locational accuracy of the species occurrences  
 268 found in GBIF, particularly those that come preserved specimens in museum

269 collections.

270 After obtaining information about species occurrence and the environment,  
271 we then merged these two datasets such that the location of each occur-  
272 rence was characterised by a vector with details of our 37 bioclimatic and  
273 topographic variables. If a species had more than one occurrence records  
274 within one of the cells of the bioclimatic raster, we only included one of  
275 the occurrence records. We did this to avoid giving more weight to areas  
276 with a high number of occurrences, a common scenario in occurrence records  
277 collected opportunistically as the ones we use here. In this step we removed  
278 85.2% of the occurrences which resulted in a total of 4.6 million occurrences  
279 used in our niche analysis.

280 A common issue of terrestrial bioclimatic datasets is that the boundaries of  
281 the cells with information do not precisely match the landmass boundaries.  
282 The result of this mismatch is that not all environmental variables was not  
283 available for 3,315 of the raster cells with occurrences (0.8% of the total).  
284 As expected, the vast majority of these problematic cells were close to the  
285 shore. To address this issue, we calculated the average value of environmental  
286 variables within an 5km buffer of the centre of the cell where the variable  
287 was missing and used it to approximate the value of the variable in that  
288 cell. Using this procedure, we were able to fill environmental variables for  
289 89.4% of the cells where they were missing. To fill the remaining 350 cells, we  
290 repeated the aforementioned procedure but instead using a 10km buffer. We  
291 removed from further analysis occurrences located within the 135 cells for  
292 which we were unable to fill environmental variables.

293 Next, we calculate the probability density function of the species distribution  
294 in environmental space. To determine the environmental space, we use  
295 the first two components from a principal component analysis of the 37  
296 bioclimatic variables associated with the species occurrences. Specifically  
297 we use the `dudi.pca` function from the R package `ade4` 1.7.13 (Dray and

298 Dufour 2007) and center and scale all bioclimatic variables to have a mean  
 299 of 0 and a unit standard deviation. We then determine the position of  
 300 species occurrences in the environmental space and estimate their bivariate  
 301 probability density function. We use a kernel method to estimate this density  
 302 and normalise it such that it ranges between zero and one. Specifically, to  
 303 calculate the probability density function we use `ecospat.grid.clim.dyn`  
 304 from the R package `ecospat` 3.0 (Broennimann, Di Cola, and Guisan 2018)  
 305 with a grid resolution of 200. We then determine the location in the environ-  
 306 mental space of the plant-pollinator communities using the function `suprow`  
 307 from `ade4`. The normalised density at that particular location corresponds  
 308 to our suitability metric, which we calculate using the R package `raster`  
 309 2.8.19 (Hijmans 2019). We use the kernel density method in the niche-factor  
 310 analysis (Broennimann et al. 2012) rather than the distance from the mode  
 311 (Hirzel et al. 2002), as it has been proposed earlier, as it has been shown to  
 312 reduce the procedure’s sensitivity to sampling effort and the resolution of  
 313 the environmental space.

314 We used a sensitivity analysis to determine the minimum number of occur-  
 315 rences that are necessary to have robust environmental suitability values  
 316 in our communities. For that we used the species with most occurrences  
 317 available, *Archilochus colubris*, and calculated the mean absolute error of the  
 318 suitability values obtained with one thousand subsamples from the 74,791  
 319 occurrences available from GBIF.

## 320 **Data analysis**

321 We then used a set of bayesian multilevel models to evaluate the impact of  
 322 environmental suitability on the size of the realised Eltonian trophic niche.  
 323 We measure the size of the Eltonian niche using the normalised degree of  
 324 species, this is, the number of species it interacts with given the number  
 325 of species in the opposite guild (Martín González, Dalsgaard, and Olesen

2010). The normalised degree was modelled using a logit link function, and a binomial distribution in which the number of species interacts with is the number of successes, and the number of species in the opposite guild is the number of trials. We are aware that whether species interacts or not is not a Bernoulli process as species interactions are not strictly independent from each other. However, a binomial distribution allows us to account for the differences in species richness across communities indirectly. Importantly, however, results are qualitatively similar when we model species degree directly using a Poisson distribution and a logarithmic link function (Supp info).

We evaluate four models to assess the relative importance of suitability. A first model, our baseline model, included five population-level predictors and two grouping levels, species and the community. The population-level predictors in the baseline model, commonly called fixed effects, were the habitat suitability, the species guild (plant or a pollinator), a metric of the species generalism, and its number of known possible partners. We included the overall level of generalism as one would expect species with a large number of interactions globally to be more likely to interact with a large number of species locally. Concretely, we estimated an overall species level of generalism as the total number of interacting species across communities in our dataset. We also included the number of known possible partners as a predictor in our models as it allows us to control for the environmental effects on species co-occurrence. We calculate this metric by determining the number of partners with which the species is known to interact in any other community. Controlling for the number of potential partners makes our model a particularly stringent test of our environmental stress hypotheses. Often the potential and the actual number of partners is the same or very close to each other, especially for rare species present only in a few communities.



355 As we were particularly interested in understanding whether the effect of  
356 habitat suitability is conditional on the species guild (plant or pollinator), we,  
357 therefore, included guild and its interaction with suitability in the model. We  
358 allowed the intercept of degree and slope of the suitability-degree relationship  
359 to vary among species, this allowed us to investigate two questions. First,  
360 it allows us to inspect the extent to which suitability is a population or  
361 a group level effect. Second, by investigating the correlation between the  
362 intercept and the slope as a model parameter, it allowed us to inspect the  
363 extent by which species with a small or large number of interactions respond  
364 to increasing levels of environmental stress. To account for unmeasured  
365 differences between communities, like sampling effort, sampling method, or  
366 diversity, we calculated an intercept for each community in our study. To  
367 facilitate model interpretation and convergence, we scaled all continuous  
368 variables to have a mean of zero and a unit standard deviation.

369 We compared this baseline model with three alternative models in which  
370 we remove one predictor at a time. To quantify the difference between  
371 models, in terms of their expected out-of-sample performance, we use the  
372 Wanatabe-Akaike information criterion (WAIC). All models were fitted under  
373 a bayesian framework using the R package `brms` 2.8.0 (Bürkner 2017, 2018)  
374 as an interface for Stan (Carpenter et al. 2017). For each model, we used  
375 four Markov chains of 4,000 iterations each; half of the iterations were used  
376 for warmup. We used weakly informative priors for all model parameters.  
377 Specifically we used normal priors of mean zero and standard deviation ten  
378 for the population-level effects and the intercepts, a half-Cauchy prior with  
379 a location of zero and a scale of two for the standard deviations, and, when  
380 applicable, an LKJ-correlation prior with parameter  $\zeta = 1$  for the correlation  
381 matrix between group-level parameters.




Figure 3: Sensitivity analysis of environmental suitability error. The number of independent occurrences retrieved from GBIF is inversely related to the error of environmental suitability for our plant-pollinator networks. The sensitivity analysis was performed by subsampling occurrences of *Archilochus colubris* the species in our dataset with the largest number of occurrences in GBIF, which was recorded in two of our communities.

## Results

After performing our sensitivity analysis, we found that we need roughly 26, 18 independent occurrences for each community for which we calculated a suitability value in order to obtain a mean absolute error below 0.1 (Fig. 3). We therefore removed from further analyses species for which we did not have enough occurrences to obtain robust estimates. When inspecting the suitability values of the analysed species we found that some communities have larger overall suitability values. However, most communities included species for which habitat suitability was low and species for which it was high (Fig. 4).

We found that our models performed relatively well. The bayesian R-squared for our baseline model was 0.91, which indicates our models were able to capture a large proportion of the variability on the data. Overall we found only weak evidence of a relationship between the environmental niche size and the normalised degree (Figure 5a). This relationship was stronger for plants than for animals. Similarly environmental suitability does not show a consistent pattern across species. Indeed, when looking at the population




Figure 4: Median habitat suitability of communities in our dataset. Each row represents a different community and horizontal lines represent span the 2.5 and 97.5 quantiles.

level effects, suitability virtually no relationship with normalised degree, neither for plants or pollinators (Figure 5b). However, suitability is still an important predictor as the WAIC difference between our baseline model and that that did not include suitability was  $574 \pm 101$ . Both among plants and pollinators for some species there is a strong negative relationship between suitability and normalised degree and some others with a strong positive relationship (Figure 6a). As expected, we found a positive relationship between a species generalism and normalised degree 5c). However our model comparisons based on WAIC shows that including generalism in the model adds relatively little information once we have accounted for the other variables in the model. Finally, we found a strong and positive relationship between the number of possible interactions and the number of realised interactions in the community.

The group level variation among communities was larger than that among species. The standard deviation (in the parameters scale) of the community intercepts was 1.02 [0.85, 1.23] while the standard deviation of the species




Figure 5: Conditional effects of predictors in our baseline model. The predicted values of the number of interacting species are based on a hypothetical community with 76 plants and 33 pollinators. These values correspond to the median number of species in each guild respectively. In each panel we condition on the mean values of all other predictors in the model, mean values for each predictor are indicated with a vertical dashed line. For model fitting, we scaled all predictors to have mean of zero and unit variance. However, with the exception of environmental niche size, here we show the unscaled predictors to facilitate interpretation. To illustrate the uncertainty around the fitted estimates we plot the fits of 100 independent draws from the posterior distribution. The thick lines indicate the mean values of the response distribution. As there was no interaction between guild and generality or the number of possible interactions, for these two predictors we only show conditional the conditional effect of pollinators.

Table 1: Comparison in out of sample predictive power of the baseline model (bold) and their alternatives. We rank models by their expected log predictive density based on their leave-one-out cross-validation information criterion (LOO). The standard error of the LOO difference provides rough guidance to the uncertainty of the model ranking. We also show the Wanatabe-Akaike information criterion (WAIC) of each model for comparison.

predictors	WAIC	SE
<b>Suit. + Gen. + Pot. + Env.</b>	6,167	135
Suit. + Pot. + Env.	6,157	138
Suit. + Gen. + Pot.	6,170	135
Suit. + Gen. + Env.	7,737	281
Gen. + Pot. + Env.	6,740	183
unknown	6,163	136

intercept was 0.5 [0.44, 0.57], and that of the species' suitability slope was 0.34 [0.28, 0.4] (95% credible intervals shown within square brackets). Interestingly, the slope the relationship between suitability and normalised degree was negatively correlated with the species' intercept in the model (Figure 6b). The mean correlation coefficient was -0.51 [-0.69, -0.3].

## Discussion

We found that environmental stress, which we measured using habitat suitability, was an important predictor for the number of partners a species interacts with. However, this relationship was highly variable among species. So much so that, at the population level, there is no discernible pattern between stress and the number of partners. Interestingly, within species, there was a significant correlation between the number of partners a species has and the stress-partners relationship itself. Specifically, species with a large number of partners in low-stress communities were more likely to have a negative relationship and hence reduce the number of partners as stress increases. Contrastingly, species in our datasets with a small number of partners in low-stress communities were more likely to have a larger number of partners in more stressful communities.






Figure 6: Species level effects of suitability. (a) Conditional effect of suitability for individual species. To facilitate visualization we show only species for which there is suitability information in at least six communities (10 plants and 33 pollinators). As in the previous figure, fitted values assume a hypothetical community of median size. In each panel we highlight two species for which the relationship between environmental suitability and normalised degree was particularly strong. (b) The correlation between the species' intercept and the species' slope of suitability was negatively correlated.

433 Previous research supported the idea that the differences in network structure  
434 along an environmental gradient were due solely to changes to community  
435 composition. In other words, it showed that the differences between two  
436 hypothetical communities with the same species were stochastic for the most  
437 part. Here, by abstracting with multiple environmental gradients into a  
438 single gradient of environmental stress, we show that this might not be  
439 the case. Instead, the environment can also have a direct impact on the  
440 number of partners a species interacts with and, therefore, influence network  
441 structure. Our results still show that the primary channel through which  
442 environment affects interaction probabilities is determining which species  
443 are present or absent from a particular community. However, they also  
444 show that, for a large number of species, it can have a substantial effect on  
445 their realised Eltonian component of the niche. Indeed, as evidenced by the  
446 predictive power of competing models, habitat suitability had a substantial  
447 contribution to predicting the number of possible partners in the community.



448 Although we cannot tell with certainty how much of the effect of environmen-  
449 tal stress can be attributed to either niche or neutral processes, the patterns  
450 we observe are likely more related to the niche. On the one hand, it has been  
451 shown that environmental models, such as our niche factor analysis, do a  
452 poor job predicting species abundances across sites (Pearce and Ferrier 2001;  
453 Sagarin, Gaines, and Gaylord 2006). It would be, therefore, somewhat naive  
454 to assume that high habitat suitability translates directly to high species  
455 abundances. On the other hand, even if species abundances are correlated  
456 with suitability, we show that within a community there is a wide range  
457 of suitabilities, even for the limited number of species we included in our  
458 analyses. Therefore, we might expect little or no overall effect of neutral  
459 processes on our response variable, as the abundance of some interaction  
460 partners might increase and decrease for others. Furthermore, the fact that  
461 the effects of environmental stress do not depend on whether the species is



462 a plant or a pollinator (for which there can be niche differences based on  
463 individual behaviour) emphasises that co-evolutionary history, which in turn  
464 shape niche processes driven by, might be driving the patterns we observe.

465 One of our most interesting results was the negative correlation between  
466 the species' intercept and the slope of the habitat suitability in our model.  
467 Specifically, species with a small intercept, this is, species that were likely  
468 to have an above-average number of partners in a community with little  
469 environmental stress, were more likely to interact with a smaller number of  
470 partners as stress increased. At least for these species, the hypothesis that  
471 species focus on partners with which they are best adapted to interact as  
472 environmental conditions deteriorate seems to hold weight. Contrastingly,  
473 species with a below-average number of partners in a low-stress environment  
474 were more likely to interact with a large number of partners as stress increased.  
475 At first glance, these species behave as facultative generalists when the  
476 environment worsens. This is certainly true for this group of species in  
477 our dataset. However, there are some nuances to this observation once we  
478 take into account the biases inherent to community-level interaction data.  
479 Specifically, we have no information about species for which environmental  
480 conditions would, in principle, allow for a positive growth rate but are absent  
481 from our datasets. A ramification of this bias is that an undetermined  
482 number of species on this second group might also be more likely to focus  
483 on partners with higher trait matching as conditions deteriorate. However,  
484 we are not able to detect that trend because these species are not recorded  
485 in community composition or interaction data. Presumably, because their  
486 preferred partners are unavailable and as such, they are absent from the  
487 community or too rare to be observed. Note that the correlation between  
488 species' intercept and the slope is unlikely to be just a byproduct of our  
489 modelling framework, as, in principle, there is nothing preventing species  
490 with an above-average number of partners to interact with even more species



491 as environmental stress increases.

492 These observations have implications for understanding the assembly of  
493 ecological networks and how it might be impacted by climate change, which  
494 currently threatens to increase the environmental stress of species within its  
495 range. Previous research suggests that flexibility of interactions in ecological  
496 communities might be a rather ubiquitous phenomenon whenever environ-  
497 mental conditions change in space or time (CaraDonna et al. 2017). This  
498 flexibility has been shown to confer robustness and stability to the loss of  
499 interaction partners (Kaiser-Bunbury et al. 2010). The current paradigm of  
500 interaction turnover suggests first that trait matching is a relatively unim-  
501 portant driver of interaction turnover for generalist species (CaraDonna et  
502 al. 2017). The fact that species with a high number of interactions were  
503 more likely to interact with fewer species as environmental stress increased  
504 indicates that trait matching might still play a role in determining the in-  
505 teractions of generalist species. Second, current research suggests that the  
506 rewiring of the interaction network is primarily driven by generalist species  
507 (Burkle, Marlin, and Knight 2013). We, however, demonstrate that at least  
508 a fraction of the species that appear to be specialist in their communities  
509 might be more flexible, if not more, than generalist species and, therefore,  
510 might have a more significant role in network persistence than previously  
511 expected. Contrastingly, specialists that are highly constrained to interact  
512 with partners of high trait-matching might be more vulnerable to increased  
513 climate change-induced environmental stress and habitat degradation. Taken  
514 together, the fact that most species appear to be flexible interaction partners,  
515 combined with the fact that those that are not flexible are less likely to per-  
516 sist, might explain why specialised communities that are highly constrained  
517 by trait matching (like some plant-hummingbird networks; Vizentin-Bugoni,  
518 Maruyama, and Sazima 2014; Maruyama et al. 2014) might be relatively  
519 rare.



520 In conclusion, we show that...

## 521 Acknowledgements

522 We thank Warwick Allen and Guadalupe Peralta for feedback and valuable  
523 discussions. EFC acknowledges the support from the University of Can-  
524 terbury Doctoral Scholarship and a New Zealand International Doctoral  
525 Research Scholarship administered by New Zealand Education. DBS and  
526 JMT acknowledge the support of Rutherford Discovery Fellowships (RDF-13-  
527 UOC-003 and RDF-UOC-1002) and the Marsden Fund Council (UOC-1705),  
528 administered by the Royal Society of New Zealand Te Apārangi.

## 529 References

- 530 Abreu, Carlos R M, and Milene F Vieira. 2004. “Os beija-flores e seus  
531 recursos florais em um fragmento florestal de Viçosa, sudeste brasileiro.”  
532 *Lundiana* 5 (2): 129–34.
- 533 Arroyo, Mary T. Kalin, Richard Primack, and Juan Armesto. 1982. “Com-  
534 munity sTUDIES IN POLLINATION ECOLOGY IN THE HIGH TEMPER-  
535 ATE ANDES OF CENTRAL CHILE. I. POLLINATION MECHANISMS  
536 AND ALTITUDINAL VARIATION.” *American Journal of Botany* 69 (1):  
537 82–97. <https://doi.org/10.1002/j.1537-2197.1982.tb13237.x>.
- 538 Barrett, Spencer C. H., and Kaius Helenurm. 1987. “The Reproductive Biol-  
539 ogy of Boreal Forest Herbs. I. Breeding Systems and Pollination.” *Canadian*  
540 *Journal of Botany* 65 (10): 2036–46. <https://doi.org/10.1139/b87-278>.
- 541 Bartomeus, Ignasi, Montserrat Vilà, and Luís Santamaría. 2008. “Contrast-  
542 ing Effects of Invasive Plants in Plant-Pollinator Networks.” *Oecologia* 155  
543 (4): 761–70. <https://doi.org/10.1007/s00442-007-0946-1>.

544 Baskett, Carina A., and Douglas W. Schemske. 2018. “Latitudinal Patterns  
545 of Herbivore Pressure in a Temperate Herb Support the Biotic Interactions  
546 Hypothesis.” Edited by Vojtech Novotny. *Ecology Letters* 21 (4): 578–87.  
547 <https://doi.org/10.1111/ele.12925>.

548 Bek, S. 2006. “A Pollination Network from a Danish Forest Meadow.” MSc  
549 Thesis, Aarhus, Denmark: Aarhus University.

550 Benadi, Gita, Thomas Hovestadt, Hans-Joachim Poethke, and Nico Blüthgen.  
551 2014. “Specialization and Phenological Synchrony of Plant-Pollinator Inter-  
552 actions Along an Altitudinal Gradient.” Edited by Thomas Ings. *Journal of*  
553 *Animal Ecology* 83 (3): 639–50. <https://doi.org/10.1111/1365-2656.12158>.

554 Bertness, Mark D., and Ragan Callaway. 1994. “Positive Interactions  
555 in Communities.” *Trends in Ecology & Evolution* 9 (5): 191–93. [https://doi.org/10.1016/0169-5347\(94\)90088-4](https://doi.org/10.1016/0169-5347(94)90088-4).

557 Bezerra, Elisângela L.S., Isabel C. Machado, and Marco A. R. Mello. 2009.  
558 “Pollination Networks of Oil-Flowers: A Tiny World Within the Smallest of  
559 All Worlds: Pollination Networks of Oil-Flowers.” *Journal of Animal Ecology*  
560 78 (5): 1096–1101. <https://doi.org/10.1111/j.1365-2656.2009.01567.x>.

561 Broennimann, Olivier, Valeria Di Cola, and Antoine Guisan. 2018. *Ecospat:*  
562 *Spatial Ecology Miscellaneous Methods*.

563 Broennimann, Olivier, Matthew C. Fitzpatrick, Peter B. Pearman, Blaise  
564 Petitpierre, Loïc Pellissier, Nigel G. Yoccoz, Wilfried Thuiller, et al. 2012.  
565 “Measuring Ecological Niche Overlap from Occurrence and Spatial Environ-  
566 mental Data: Measuring Niche Overlap.” *Global Ecology and Biogeography*  
567 21 (4): 481–97. <https://doi.org/10.1111/j.1466-8238.2011.00698.x>.

568 Bundgaard, M. 2003. “Tidslig Og Rumlig Variation I et Plante-  
569 Bestøvernetv\’a Erk.” MSc Thesis, Aarhus, Denmark: Aarhus University.

570 Burkle, L. A., J. C. Marlin, and T. M. Knight. 2013. “Plant-Pollinator

571 Interactions over 120 Years: Loss of Species, Co-Occurrence, and Function.”  
572 *Science* 339 (6127): 1611–5. <https://doi.org/10.1126/science.1232728>.

573 Bürkner, Paul-Christian. 2017. “Brms: An R Package for Bayesian Multilevel  
574 Models Using Stan.” *Journal of Statistical Software* 80 (1). [https://doi.org/](https://doi.org/10.18637/jss.v080.i01)  
575 10.18637/jss.v080.i01.

576 ———. 2018. “Advanced Bayesian Multilevel Modeling with the R Package  
577 Brms.” *The R Journal* 10 (1): 395. <https://doi.org/10.32614/RJ-2018-017>.

578 CaraDonna, Paul J., William K. Petry, Ross M. Brennan, James L. Cun-  
579 ningham, Judith L. Bronstein, Nickolas M. Waser, and Nathan J. Sanders.  
580 2017. “Interaction Rewiring and the Rapid Turnover of Plant-Pollinator  
581 Networks.” Edited by Ferenc Jordan. *Ecology Letters* 20 (3): 385–94.  
582 <https://doi.org/10.1111/ele.12740>.

583 Carpenter, Bob, Andrew Gelman, Matthew D. Hoffman, Daniel Lee, Ben  
584 Goodrich, Michael Betancourt, Marcus Brubaker, Jiqiang Guo, Peter Li,  
585 and Allen Riddell. 2017. “Stan : A Probabilistic Programming Language.”  
586 *Journal of Statistical Software* 76 (1). <https://doi.org/10.18637/jss.v076.i01>.

587 Chamberlain, Scott A, and Carl Boettiger. 2017. “R Python, and Ruby  
588 Clients for GBIF Species Occurrence Data.” Preprint. PeerJ Preprints.  
589 <https://doi.org/10.7287/peerj.preprints.3304v1>.

590 Chamberlain, Scott, Vijay Barve, Dan Mcglinn, Damiano Oldoni, Peter  
591 Desmet, Laurens Geffert, and Karthik Ram. 2019. *Rgbif: Interface to the*  
592 *Global Biodiversity Information Facility API*.

593 Chamberlain, Scott, and Eduard Szocs. 2013. “Taxize - Taxonomic Search  
594 and Retrieval in R.” *F1000Research*.

595 Chamberlain, Scott, Eduard Szocs, Zachary Foster, Zebulun Arendsee, Carl  
596 Boettiger, Karthik Ram, Ignasi Bartomeus, et al. 2019. *Taxize: Taxonomic*  
597 *Information from Around the Web*.

598 Clements, Frederic Edward, and Frances Louise Long. 1923. *Experimental*  
 599 *Pollination: An Outline of the Ecology of Flowers and Insects*. 336. Carnegie  
 600 Institution of Washington.

601 del Coro Arizmendi, Ma., and Juan Francisco Ornelas. 1990. "Hummingbirds  
 602 and Their Floral Resources in a Tropical Dry Forest in Mexico." *Biotropica*  
 603 22 (2): 172. <https://doi.org/10.2307/2388410>.

604 Devoto, Mariano, Diego Medan, and Norberto H. Montaldo. 2005. "Patterns  
 605 of Interaction Between Plants and Pollinators Along an Environmental  
 606 Gradient." *Oikos* 109 (3): 461–72. [https://doi.org/10.1111/j.0030-1299.2005.](https://doi.org/10.1111/j.0030-1299.2005.13712.x)  
 607 13712.x.

608 Dicks, L. V., S. A. Corbet, and R. F. Pywell. 2002. "Compartmentalization  
 609 in Plant-Insect Flower Visitor Webs." *Journal of Animal Ecology* 71 (1):  
 610 32–43. <https://doi.org/10.1046/j.0021-8790.2001.00572.x>.

611 Dray, Stéphane, and Anne-Béatrice Dufour. 2007. "The Ade4 Package:  
 612 Implementing the Duality Diagram for Ecologists." *Journal of Statistical*  
 613 *Software* 22 (4). <https://doi.org/10.18637/jss.v022.i04>.

614 Dupont, Yoko L., Dennis M. Hansen, and Jens M. Olesen. 2003. "Structure  
 615 of a Plant-Flower-Visitor Network in the High-Altitude Sub-Alpine Desert  
 616 of Tenerife, Canary Islands." *Ecography* 26 (3): 301–10. [https://doi.org/10.](https://doi.org/10.1034/j.1600-0587.2003.03443.x)  
 617 1034/j.1600-0587.2003.03443.x.

618 Dupont, Yoko L., and Jens M. Olesen. 2009. "Ecological Modules and Roles  
 619 of Species in Heathland Plant-Insect Flower Visitor Networks." *Journal of*  
 620 *Animal Ecology* 78 (2): 346–53. [https://doi.org/10.1111/j.1365-2656.2008.](https://doi.org/10.1111/j.1365-2656.2008.01501.x)  
 621 01501.x.

622 Elberling, Heidi, and Jens M. Olesen. 1999. "The Structure of a High  
 623 Latitude Plant-Flower Visitor System: The Dominance of Flies." *Ecography*  
 624 22 (3): 314–23. <https://doi.org/10.1111/j.1600-0587.1999.tb00507.x>.

625 Ferreira Canela, Maria Bernadete. 2006. “Interações Entre Plantas E Beija-  
626 Flores Numa Comunidade de Floresta Atlantica Montana Em Itatiaia, Rio  
627 de Janeiro.” PhD Thesis, Campinas, Brazil: Universidade Estadual de  
628 Campinas.

629 Fick, Stephen E., and Robert J. Hijmans. 2017. “WorldClim 2: New 1-Km  
630 Spatial Resolution Climate Surfaces for Global Land Areas: NEW CLI-  
631 MATE SURFACES FOR GLOBAL LAND AREAS.” *International Journal*  
632 *of Climatology* 37 (12): 4302–15. <https://doi.org/10.1002/joc.5086>.

633 Fortuna, Miguel A., Raul Ortega, and Jordi Bascompte. 2014. “The Web of  
634 Life.” *arXiv:1403.2575 [Q-Bio]*, March. <http://arxiv.org/abs/1403.2575>.

635 Gravel, Dominique, Benjamin Baiser, Jennifer A Dune, Jens-Peter Kopelke,  
636 Neo D Martinez, Tommi Nyman, Timothee Poisot, et al. 2018. “Bringing  
637 Elton and Grinnell Together: A Quantitative Framework to Represent the  
638 Biogeography of Ecological Interaction Networks.” *Ecography* 41: 1–15.  
639 <https://doi.org/10.1111/ecog.04006>.

640 Gutierrez Z, Aquiles, Sandra Victoria Rojas-Nossa, and Gary F Stiles. 2004.  
641 “DINÁMICA ANUAL DE LA INTERACCIÓN COLIBRÍ-FLOR EN ECO-  
642 SISTEMAS ALTOANDINOS.” *Ornitologia Neotropical* 15: 205–13.

643 Hattersley-Smith, G. 1985. “Botanical Studies in the Lake Hazen Region,  
644 Northern Ellesmere Island, Northwest Territories, Canada, by James H.  
645 Soper and John M. Powell.” *ARCTIC* 38 (4): 346. [https://doi.org/10.14430/](https://doi.org/10.14430/arctic2421)  
646 [arctic2421](https://doi.org/10.14430/arctic2421).

647 Herrera, Javier. 1988. “Pollination Relationships in Southern Spanish  
648 Mediterranean Shrublands.” *The Journal of Ecology* 76 (1): 274. <https://doi.org/10.2307/2260469>.

650 Hijmans, Robert J. 2019. *Raster: Geographic Data Analysis and Modeling*.

651 Hirzel, A H, J Hausser, D Chessel, and N Perrin. 2002. “ECOLOGICAL-

652 NICHE FACTOR ANALYSIS: HOW TO COMPUTE HABITAT-  
653 SUITABILITY MAPS WITHOUT ABSENCE DATA?" 83 (7): 10.

654 Hocking, Brian. 1968. "Insect-Flower Associations in the High Arctic with  
655 Special Reference to Nectar." *Oikos* 19 (2): 359. [https://doi.org/10.2307/](https://doi.org/10.2307/3565022)  
656 3565022.

657 Hoiss, Bernhard, Jochen Krauss, Simon G. Potts, Stuart Roberts, and  
658 Ingolf Steffan-Dewenter. 2012. "Altitude Acts as an Environmental Filter  
659 on Phylogenetic Composition, Traits and Diversity in Bee Communities."  
660 *Proceedings of the Royal Society B: Biological Sciences* 279 (1746): 4447–56.  
661 <https://doi.org/10.1098/rspb.2012.1581>.

662 Hoiss, Bernhard, Jochen Krauss, and Ingolf Steffan-Dewenter. 2015. "Inter-  
663 active Effects of Elevation, Species Richness and Extreme Climatic Events  
664 on Plant-Pollinator Networks." *Global Change Biology* 21 (11): 4086–97.  
665 <https://doi.org/10.1111/gcb.12968>.

666 Hutchinson, Matthew C., Edgar Fernando Cagua, and Daniel B. Stouffer.  
667 2017. "Cophylogenetic Signal Is Detectable in Pollination Interactions Across  
668 Ecological Scales." *Ecology* 98 (10): 2640–52. [https://doi.org/10.1002/ecy.](https://doi.org/10.1002/ecy.1955)  
669 1955.

670 Ingversen, TT. 2006. "Plant-Pollinator Interactions on Jamaica and Do-  
671 minica: The Centrality, Asymmetry and Modularity of Networks." MSc  
672 Thesis, Aarhus, Denmark: University of Aarhus.

673 INouE, Tamiji, Makoto KATo, Takehiko KAKuTANi, Takeshi SuKA, and  
674 Takao It. 1990. "Of Kibune, Kyoto: An Overview of the Flowering Phenology  
675 and the Seasonal Pattern of Insect Visits'," 88.

676 Inouye, David W., and Graham H. Pyke. 1988. "Pollination Biology in the  
677 Snowy Mountains of Australia: Comparisons with Montane Colorado, USA."  
678 *Austral Ecology* 13 (2): 191–205. [https://doi.org/10.1111/j.1442-9993.1988.](https://doi.org/10.1111/j.1442-9993.1988.tb00968.x)  
679 tb00968.x.

680 Kaiser-Bunbury, Christopher N., Stefanie Muff, Jane Memmott, Christine  
681 B. Müller, and Amedeo Caflisch. 2010. “The Robustness of Pollination  
682 Networks to the Loss of Species and Interactions: A Quantitative Approach  
683 Incorporating Pollinator Behaviour.” *Ecology Letters* 13 (4): 442–52. <https://doi.org/10.1111/j.1461-0248.2009.01437.x>.  
684

685 Kaiser-Bunbury, Christopher N., Diego P. Vázquez, Martina Stang, and  
686 Jaboury Ghazoul. 2014. “Determinants of the Microstructure of PlantPollina-  
687 tor Networks.” *Ecology* 95 (12): 3314–24. <https://doi.org/10.1890/14-0024.1>.

688 Kakutani, Takehiko, Tamiji Inoue, Makoto Kato, and Hidekuji Ichihashi.  
689 1990. “Insect-Flower Relationship in the Campus of Kyoto University, Kyoto  
690 : An Overview of the Flowering Phenology and the Seasonal Pattern of Insect  
691 Visits.” *Contributions from the Biological Laboratory, Kyoto University* 27  
692 (4): 465–522.

693 KATo, Makoto. 2000. “Anthophilous Insect Community and Plant-Pollinator  
694 Interactions on Amami Islands in the Ryukyu Archipelago, Japan,” 101.

695 Kato, Makoto, Takehiko Kakutani, Tamiji Inoue, and Takao Itino. 1990.  
696 “Insect-Flower Relationship in the Primary Beech Forest of Ashu, Kyoto: An  
697 Overview of the Flowering Phenology and the Seasonal Pattern of Insect  
698 Visits,” 68.

699 KATo, Makoto, Masamichi Matsumoto, and Toru Kato. 1993. “Flowering  
700 Phenology and Anthophilous Insect Community in the Cool-Temperate  
701 Subalpine Forests and Meadows at Mt. Kushigata in the Central Part of  
702 Japan,” 58.

703 Kato, Makoto, and Reiichi Miura. 1993. “Flowering Phenology and An-  
704 thophilous Insect Community at a Threatened Natural Lowland Marsh at  
705 Nakaikemi in Tsuruga, Japan.” *Contributions from the Biological Laboratory,*  
706 *Kyoto University* 29 (1): 1.

707 Kevan, PG. 1970. “High Arctic Insect-Flower Relations: The Interrelation-



ships of Arthropods and Flowers at Lake Hazen, Ellesmere Island, Northwest Territories, Canada.” PhD Thesis, Edmonton, Canada: University of Alberta.

Kohler, Glaucio Ubiratan. 2011. “Redes de Interação Planta Beija-Flor Em Um Gradiente Altitudinal de Floresta Atlântica No Sul Do Brasil.” MSc Thesis, Curitiba, Brasil: Universidade Federal do Paraná.

Lara, Carlos. 2006. “Temporal Dynamics of Flower Use by Hummingbirds in a Highland Temperate Forest in Mexico.” *Ecoscience* 13 (1): 23–29. [https://doi.org/10.2980/1195-6860\(2006\)13\[23:TDOFUB\]2.0.CO;2](https://doi.org/10.2980/1195-6860(2006)13[23:TDOFUB]2.0.CO;2).

Las-Casas, Fmg, Sm Azevedo Júnior, and Mm Dias Filho. 2012. “The Community of Hummingbirds (Aves: Trochilidae) and the Assemblage of Flowers in a Caatinga Vegetation.” *Brazilian Journal of Biology* 72 (1): 51–58. <https://doi.org/10.1590/S1519-69842012000100006>.

Lasprilla, Liliana Rosero, and others. 2003. “Interações Planta/Beija-Flor Em Três Comunidades Vegetais Da Parte Sul Do Parque Nacional Natural Chiribiquete, Amazonas (Colômbia).” PhD Thesis, Campinas, Brazil: Universidade Estadual de Campinas.

Lavandero, Blas, and Jason M. Tylianakis. 2013. “Genotype Matching in a Parasitoid-Host Genotypic Food Web: An Approach for Measuring Effects of Environmental Change.” *Molecular Ecology* 22 (1): 229–38. <https://doi.org/10.1111/mec.12100>.

Lundgren, Rebekka, and Jens M. Olesen. 2005. “The Dense and Highly Connected World of Greenland’s Plants and Their Pollinators.” *Arctic, Antarctic, and Alpine Research* 37 (4): 514–20. [https://doi.org/10.1657/1523-0430\(2005\)037\[0514:TDAHWC\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0514:TDAHWC]2.0.CO;2).

Martín González, Ana M., Bo Dalsgaard, and Jens M. Olesen. 2010. “Centrality Measures and the Importance of Generalist Species in Pollination Networks.” *Ecological Complexity* 7 (1): 36–43. <https://doi.org/10.1016/j.ecocom.2009.03.008>.

736 Maruyama, Pietro K., Jeferson Vizentin-Bugoni, Genilda M. Oliveira, Paulo  
737 E. Oliveira, and Bo Dalsgaard. 2014. "Morphological and Spatio-Temporal  
738 Mismatches Shape a Neotropical Savanna Plant-Hummingbird Network."  
739 *Biotropica* 46 (6): 740–47. <https://doi.org/10.1111/btp.12170>.

740 McKinnon, L., P. A. Smith, E. Nol, J. L. Martin, F. I. Doyle, K. F. Abraham,  
741 H. G. Gilchrist, R. I. G. Morrison, and J. Bêty. 2010. "Lower Predation Risk  
742 for Migratory Birds at High Latitudes." *Science, New Series* 327 (5963):  
743 326–27.

744 McMullen, C. K. 106AD. "Flower-Visiting Insects of the Galapagos Islands."  
745 *The Pan-Pacific Entomologist* 69 (1): 95.

746 Medan, Diego, Norberto H. Montaldo, Mariano Devoto, Anita Mantese,  
747 Viviana Vasellati, German G. Roitman, and Norberto H. Bartoloni. 2002.  
748 "Plant-Pollinator Relationships at Two Altitudes in the Andes of Mendoza,  
749 Argentina." *Arctic, Antarctic, and Alpine Research* 34 (3): 233. <https://doi.org/10.2307/1552480>.

750

751 Memmott, J. 1999. "The Structure of a Plant-Pollinator Food Web." *Ecology*  
752 *Letters* 2 (5): 276–80. <https://doi.org/10.1046/j.1461-0248.1999.00087.x>.

753 Montero, AC. 2005. "The Ecology of Three Pollination Networks." MSc  
754 Thesis, Aarhus, Denmark: Aarhus University.

755 Mosquin, Theodore. 1967. "Observations on the Pollination Biology of  
756 Plants on Melville Island, NWT." *Can. Fld Nat.* 81: 201–5.

757 Motten, Alexander F. 1986. "Pollination Ecology of the Spring Wildflower  
758 Community of a Temperate Deciduous Forest." *Ecological Monographs* 56  
759 (1): 21–42. <https://doi.org/10.2307/2937269>.

760 Olesen, Jens M., Louise I. Eskildsen, and Shadila Venkatasamy. 2002.  
761 "Invasion of Pollination Networks on Oceanic Islands: Importance of In-  
762 vader Complexes and Endemic Super Generalists." *Diversity & Distribution*

763 *Glyph="&" Ascii="É&"/> Distributions* 8 (3): 181–92. [https:](https://doi.org/10.1046/j.1472-4642.2002.00148.x)  
764 [//doi.org/10.1046/j.1472-4642.2002.00148.x](https://doi.org/10.1046/j.1472-4642.2002.00148.x).

765 Ollerton, J. 2003. “The Pollination Ecology of an Assemblage of Grassland  
766 Asclepiads in South Africa.” *Annals of Botany* 92 (6): 807–34. [https:](https://doi.org/10.1093/aob/mcg206)  
767 [//doi.org/10.1093/aob/mcg206](https://doi.org/10.1093/aob/mcg206).

768 Pearce, Jennie, and Simon Ferrier. 2001. “The Practical Value of Modelling  
769 Relative Abundance of Species for Regional Conservation Planning: A Case  
770 Study.” *Biological Conservation* 98 (1): 33–43. [https://doi.org/10.1016/](https://doi.org/10.1016/S0006-3207(00)00139-7)  
771 [S0006-3207\(00\)00139-7](https://doi.org/10.1016/S0006-3207(00)00139-7).

772 Pellissier, Loïc, Julien Pottier, Pascal Vittoz, Anne Dubuis, and Antoine  
773 Guisan. 2010. “Spatial Pattern of Floral Morphology: Possible Insight into  
774 the Effects of Pollinators on Plant Distributions.” *Oikos* 119 (11): 1805–13.  
775 <https://doi.org/10.1111/j.1600-0706.2010.18560.x>.

776 Peralta, Guadalupe, Carol M. Frost, Raphael K. Didham, Arvind Varsani,  
777 and Jason M. Tylianakis. 2015. “Phylogenetic Diversity and Co-Evolutionary  
778 Signals Among Trophic Levels Change Across a Habitat Edge.” Edited  
779 by Frank van Veen. *Journal of Animal Ecology* 84 (2): 364–72. [https:](https://doi.org/10.1111/1365-2656.12296)  
780 [//doi.org/10.1111/1365-2656.12296](https://doi.org/10.1111/1365-2656.12296).

781 Percival, Mary. 1974. “Floral Ecology of Coastal Scrub in Southeast Jamaica.”  
782 *Biotropica* 6 (2): 104. <https://doi.org/10.2307/2989824>.

783 Petanidou, Theodora, and Despina Vokou. 1993. “POLLINATION ECOLOGY OF LABIATAE IN A PHRYGANIC (EAST MEDITERRANEAN)  
784 ECOSYSTEM.” *American Journal of Botany* 80 (8): 892–99. [https://doi.](https://doi.org/10.1002/j.1537-2197.1993.tb15310.x)  
785 [org/10.1002/j.1537-2197.1993.tb15310.x](https://doi.org/10.1002/j.1537-2197.1993.tb15310.x).

787 Philipp, Marianne, Jens Böcher, Hans R. Siegismund, and Lene R. Nielsen.  
788 2006. “Structure of a Plant-Pollinator Network on a Pahoehoe Lava Desert  
789 of the Galápagos Islands.” *Ecography* 29 (4): 531–40. [https://doi.org/10.](https://doi.org/10.1111/j.0906-7590.2006.04546.x)  
790 [1111/j.0906-7590.2006.04546.x](https://doi.org/10.1111/j.0906-7590.2006.04546.x).

791 Pires, Aliny P. F., Nicholas A. C. Marino, Diane S. Srivastava, and Vinicius  
792 F. Farjalla. 2016. "Predicted Rainfall Changes Disrupt Trophic Interactions  
793 in a Tropical Aquatic Ecosystem." *Ecology* 97 (10): 2750–9. [https://doi.org/](https://doi.org/10.1002/ecy.1501)  
794 10.1002/ecy.1501.

795 Primack, Richard B. 1983. "Insect Pollination in the New Zealand Mountain  
796 Flora." *New Zealand Journal of Botany* 21 (3): 317–33. [https://doi.org/10.](https://doi.org/10.1080/0028825X.1983.10428561)  
797 1080/0028825X.1983.10428561.

798 Ramirez, Nelson. 1989. "Biología de Polinización En Una Comunidad  
799 Arbustiva Tropical de La Alta Guayana Venezolana." *Biotropica* 21 (4): 319.  
800 <https://doi.org/10.2307/2388282>.

801 Ramirez, Nelson, and Yseleny Brito. 1992. "Pollination Biology in a Palm  
802 Swamp Community in the Venezuelan Central Plains." *Botanical Journal of*  
803 *the Linnean Society* 110 (4): 277–302. [https://doi.org/10.1111/j.1095-8339.](https://doi.org/10.1111/j.1095-8339.1992.tb00294.x)  
804 1992.tb00294.x.

805 Robertson, Charles. 1929. *Flowers and Insects Lists of Visitors of Four*  
806 *Hundred and Fifty Three Flowers*. Cairnville: The Science Press Printing  
807 Company.

808 Sabatino, Malena. 2010. "Direct Effects of Habitat Area on Interaction  
809 Diversity in Pollination Webs." *Ecological Applications* 20 (6): 7.

810 Sagarin, Raphael D., Steven D. Gaines, and Brian Gaylord. 2006. "Moving  
811 Beyond Assumptions to Understand Abundance Distributions Across the  
812 Ranges of Species." *Trends in Ecology & Evolution* 21 (9): 524–30. <https://doi.org/10.1016/j.tree.2006.06.008>.  
813 <https://doi.org/10.1016/j.tree.2006.06.008>.

814 Schemske, Douglas W., Mary F. Willson, Michael N. Melampy, Linda J.  
815 Miller, Louis Verner, Kathleen M. Schemske, and Louis B. Best. 1978.  
816 "Flowering Ecology of Some Spring Woodland Herbs." *Ecology* 59 (2): 351–  
817 66. <https://doi.org/10.2307/1936379>.

- 818 Small, Ernest. 1976. "Insect Pollinators of the Mer Bleue Peat Bog of  
819 Ottawa." *Canadian Field-Naturalist*.
- 820 Smith-Ramírez, C., P. Martinez, M. Nuñez, C. González, and J. J. Armesto.  
821 2005. "Diversity, Flower Visitation Frequency and Generalism of Pollinators  
822 in Temperate Rain Forests of Chiloé Island, Chile." *Botanical Journal of the*  
823 *Linnean Society* 147 (4): 399–416. [https://doi.org/10.1111/j.1095-8339.2005.](https://doi.org/10.1111/j.1095-8339.2005.00388.x)  
824 00388.x.
- 825 Stald, L, A Valido, and J Olesen. 2003. "Struktur Og Dynamik I Rum Og  
826 Tid at et Bestøvningsnetværk På a Tenerife, De Kanariske Øer." MSc-Thesis,  
827 Aarhus, Denmark: University of Aarhus.
- 828 Title, Pascal O., and Jordan B. Bemmels. 2017. "ENVIREM: ENVironmen-  
829 tal Rasters for Ecological Modeling Version 1.0." University of Michigan.  
830 <https://doi.org/10.7302/Z2BR8Q40>.
- 831 ———. 2018. "ENVIREM: An Expanded Set of Bioclimatic and Topographic  
832 Variables Increases Flexibility and Improves Performance of Ecological Niche  
833 Modeling." *Ecography* 41 (2): 291–307. <https://doi.org/10.1111/ecog.02880>.
- 834 Tur, C., A. Sáez, A. Traveset, and M. A. Aizen. 2016. "Evaluating the  
835 Effects of Pollinator-Mediated Interactions Using Pollen Transfer Networks:  
836 Evidence of Widespread Facilitation in South Andean Plant Communities."  
837 Edited by José María Gómez. *Ecology Letters* 19 (5): 576–86. <https://doi.org/10.1111/ele.12594>.  
838
- 839 Tylianakis, Jason M., and Rebecca J. Morris. 2017. "Ecological Net-  
840 works Across Environmental Gradients." *Annual Review of Ecology,*  
841 *Evolution, and Systematics* 48 (1): 25–48. [https://doi.org/10.1146/](https://doi.org/10.1146/annurev-ecolsys-110316-022821)  
842 annurev-ecolsys-110316-022821.
- 843 Vázquez, Diego P. 2002. "Interactions Among Introduced Ungulates, Plants,  
844 and Pollinators: A Field Study in the Temperate Forest of the Southern  
845 Andes." PhD Thesis, Knoxville, United States of America: University of

846 Tennessee.

847 Vizentin-Bugoni, Jeferson, Pietro K. Maruyama, Vanderlei J. Debastiani, L.  
848 da S. Duarte, Bo Dalsgaard, and Marlies Sazima. 2016. “Influences of Sam-  
849 pling Effort on Detected Patterns and Structuring Processes of a Neotropical  
850 Plant-Hummingbird Network.” Edited by Daniel Stouffer. *Journal of Animal*  
851 *Ecology* 85 (1): 262–72. <https://doi.org/10.1111/1365-2656.12459>.

852 Vizentin-Bugoni, Jeferson, Pietro Kiyoshi Maruyama, and Marlies Sazima.  
853 2014. “Processes Entangling Interactions in Communities: Forbidden Links  
854 Are More Important Than Abundance in a HummingbirdPlant Network.”  
855 *Proceedings of the Royal Society B: Biological Sciences* 281 (1780): 20132397.  
856 <https://doi.org/10.1098/rspb.2013.2397>.

857 Vucic-Pestic, Olivera, Roswitha B. Ehnes, Björn C. Rall, and Ulrich Brose.  
858 2011. “Warming up the System: Higher Predator Feeding Rates but Lower  
859 Energetic Efficiencies: WARMING AND FUNCTIONAL RESPONSES.”  
860 *Global Change Biology* 17 (3): 1301–10. [https://doi.org/10.1111/j.1365-2486.](https://doi.org/10.1111/j.1365-2486.2010.02329.x)  
861 2010.02329.x.

862 Yamazaki, Kyoko, and Makoto Kato. 2003. “Flowering Phenology and  
863 Anthophilous Insect Community,” 67.

864 Zizka, Alexander, Daniele Silvestro, Tobias Andermann, Josué Azevedo,  
865 Camila Duarte Ritter, Daniel Edler, Harith Farooq, et al. 2019. “Coordi-  
866 nateCleaner: Standardized Cleaning of Occurrence Records from Biological  
867 Collection Databases.” Edited by Tiago Quental. *Methods in Ecology and*  
868 *Evolution* 10 (5): 744–51. <https://doi.org/10.1111/2041-210X.13152>.

# Environmental stress affects niche breadth in plant-pollinator communities

Supplementary information

*E. Fernando Cagua, Hugo J. Marrero, Jason M. Tylianakis, Daniel B. Stouffer*

Table S1: Manually corrected canonical names. More than one correct name have been included when an accepted/valid synonym the canonical name exists.

incorrect name	corrected name	guild
Acaena pinn	<i>Acaena pinnatifida</i>	plant
Adesmia brachy	<i>Adesmia brachysemeon</i>	plant
Aesculus camea	<i>Aesculus X carnea</i>	plant
Brachyome sinclairii	<i>Brachyscome sinclairii</i>	plant
Calceolaria arac	<i>Calceolaria arachnoidea</i>	plant
Equium sabulicola	<i>Echium sabulicola</i>	plant
Euonymus fo rtunei	<i>Euonymus fortunei</i>	plant
Galvezia leucantha pubescen	<i>Galvezia leucantha</i>	plant
Heliconia simulans	<i>Heliconia angusta</i>	plant
Pitcaimia flammea	<i>Pitcairnia flammea</i>	plant
Psittacanthus flavo viridis	<i>Psittacanthus flavo-viridis</i>	plant
Rhodophiala bifidum	<i>Rhodophiala bifida</i>	plant
Stachys albi	<i>Stachys albicaulis</i>	plant
Stenactis annuus	<i>Erigeron annuus</i>	plant
Thaspium aureum atropurpurem	<i>Thaspium trifoliatum</i>	plant
Tristhema mauritiana	<i>Tristemma mauritianum</i>	plant
Tropaeolum polyph	<i>Tropaeolum polyphyllum</i>	plant
Tyttnera scabra	<i>Turnera scabra</i>	plant
	<i>Turnera ulmifolia</i>	plant
VVedelia biflora	<i>Melanthera biflora</i>	plant
	<i>Wedelia biflora</i>	plant
Cateres pennatus	<i>Kateretes pennatus</i>	pollinator
Eclimus harrisi	<i>Condyllostylus crinicauda</i>	pollinator
Ptilandrena g. maculati	<i>Andrena distans</i>	pollinator
Tapinotaspis caerulea	<i>Chalepogenus caeruleus</i>	pollinator
Tapinotaspis herbsti	<i>Chalepogenus herbsti</i>	pollinator