

# Draft information

version: draft-0.5

2020-04-18

## Done

- Included feedback to paper (thanks Jason)
- Added many references including some to GBIF datasets
- Minor adjustments to figure captions
- Tidied up the supplementary materials

## To-do

- Add a few references to the methods justifying occurrence cleaning approach (thanks Audrey)

## Won't-do

- Finish the manual checks for species names that we were unable to correct automatically. It takes ~5-10 mins per species and we have ~600. Given the amount of data we already have it probably won't make much of a difference.

## Comments

- Jason, one of your comments was: *"I can't remember if I've raised this before, but if species are near to the boundaries of number of partners (close to zero or close to  $N$ , where  $N$  is the number of species in a community), a random change in the number of partners is more likely to bring them towards the mean because they can't go any higher than  $N$  or lower than 0. Is there any way to deal with this?"*. That certainly might be happening for species with few partners. I try to address that in the discussion (see page 15, line 308). I don't think it happens for species with many partners because very few of the species analysed are close to  $N$  (only ~2% of plants and ~3% of animals have a normalised degree  $> 0.5$ , ~0.5% of both have a normalised degree  $> 0.75$ ). To

avoid confusion I've removed the "regression to the mean" statements in the discussion.

- Appart from that, I think I followed most suggestions one way or another. **Thanks!**

# Environmental stress drives intermediate plant and pollinator specialisation

*E. Fernando Cagua*<sup>1</sup> (*efc29@uclive.ac.nz*)

*Audrey Lustig*<sup>2</sup> (*audrey.lustig@canterbury.ac.nz*)

*Jason M. Tylianakis*<sup>1</sup> (*jason.tylianakis@canterbury.ac.nz*)

*Daniel B. Stouffer*<sup>1</sup> (*daniel.stouffer@canterbury.ac.nz*)

<sup>1</sup> Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury, Private Bag 4800, Christchurch 8041, New Zealand

<sup>2</sup> Geospatial Research Institute, University of Canterbury, Private Bag 4800, Christchurch 8041, New Zealand

**Running title:** Environmental stress drives specialisation.

**Keywords:** eltonian niche, environmental effects, generalisation & specialisation, species degree, species interactions, and trophic niche

**Type of article:** Letter.

**Number of words:** 201 in abstract; 4,163 in main text.

**Number of displays:** 4 figures; 1 tables; 0 text boxes.

**Number of references:** 92

**Author for correspondence:** E. Fernando Cagua (+64 20 4026 8153).

**Data accessibility:** Data supporting the results will be accessible in an appropriate data repository after publication. We will include the DOI here.

**Author contributions:** EFC conceived the study. EFC and AL compiled cleaned the data. EFC analysed the data. All authors contributed to the writing of the manuscript and study design.

## Abstract

What determines whether or not a species is a generalist or a specialist? Evidence that the environment can influence species interactions is rapidly accumulating. However, a systematic link between environment and the number of partners a species interacts with has been elusive so far; presumably, because environmental gradients appear to have contrasting effects on species depending on the specific environmental variable. Here, we test whether the stresses imposed by the environment, instead of environmental gradients directly, influence species specialisation using a global dataset of plant-pollinator interactions. We found that environmental stress can significantly affect specialisation, even when accounting for changes in community composition, likely by interacting with species' traits and evolutionary history. Species that have a large number of interactions tended to focus on fewer, presumably higher-quality, interactions under stressful environmental conditions. Contrastingly, the specialists present in multiple locations tended to broaden their niche, presumably engaging in opportunistic interactions to cope with increased environmental stress. Indeed, many apparent specialists effectively behaved as facultative generalists. Overall, many of the species we analysed are not inherently generalist or specialist. Instead, the species' level of specialisation should be considered on a relative scale depending on the environmental conditions at a given location.

## Introduction

Species interactions are known to vary widely across space and time (Laliberté & Tylianakis 2012; Schleuning *et al.* 2012; Poisot *et al.* 2015; Trøjelsgaard *et al.* 2015). There are multiple examples of species that interact with a large number of partners in a particular community or season, but with fewer in another (Olesen *et al.* 2008, 2011; Dupont *et al.* 2009; Benadi *et al.* 2014; Rabeling *et al.* 2019). Some of this variation can be attributed to environmental drivers (Tylianakis & Morris 2017). However, it remains unknown how generalisable characteristics of the environment, specifically the stress it imposes on species, affects whether two species interact, and, ultimately, the species' specialisation. Understanding how the environment drives the number of partners is crucial because it underpins the species' role in its community and shapes the structure of the network of interactions (Cirtwill *et al.* 2018). This structure, in turn, determines ecosystem function and stability (Thebault & Fontaine 2010).

Species interactions are determined in part by niche processes (the matching of traits) and partly by neutral processes (more abundant species are more likely to encounter each other and, thus, interact) (Vázquez *et al.* 2009a, b). The environment can influence both of these processes (Godsoe *et al.* 2017). It is, therefore, not surprising that, despite limitations on the spatial extent or the number of environmental gradients considered, multiple studies have shown how changes to interactions can be related to environmental change (Tylianakis & Morris 2017). Despite widespread evidence that species interactions and their networks can change along environmental gradients, the direction of change can be variable (Devoto *et al.* 2005; Pires *et al.* 2016; Baskett & Schemske 2018), such that until now it has been impossible to extrapolate previous findings to new kinds of gradients. Overall, while it looks clear that pairwise interactions respond to environmental drivers, there is high variability in the response (Tylianakis *et al.* 2008).

One possible explanation for the seemingly contradictory evidence is that different bioclimatic factors (like temperature or precipitation) can have contrasting effects on species and their partners, such that we lack a common currency with which to derive general responses to the environment. Here we attempt to overcome these barriers to deriving general rules by reducing multiple factors into a single measure of environmental stress. Previous research suggests that environmental stress may

affect the number of interaction partners a species has in different ways depending on the species' role in the community (for example its trophic guild) (Tylianakis & Morris 2017). Specifically, we propose two alternative hypotheses of how environmental stress may affect specialisation. First, species under environmental stress might be “pressured” to focus on partners with which they are best adapted to interact. For instance, Hoiss *et al.* (2012) found increased phylogenetic clustering between plants and pollinators at higher altitudes; while Peralta *et al.* (2015) found that parasitoids in plantation forest, where environmental stress was higher than in native forests, constrained their host use to those with which phylogenetic matching was strongest. Similarly, Lavandero & Tylianakis (2013) showed that environmental stress due to higher temperature reduced the trophic niche breadth (range of host genotypes used) of parasitoids, suggesting higher specialisation.

Alternatively, it is also possible that when species are under environmental stress, they are forced to be more flexible in their interactions. Higher environmental stress is likely to be reflected in greater energetic or reproductive costs. Therefore, they might not be able to sustain encounter rates with their preferred partners at sufficient levels. In line with this hypothesis, Hoiss *et al.* (2015) found that the specialisation of plant-pollinator networks decreased both with elevation and after extreme drought events. Likewise, Pellissier *et al.* (2010) found a positive relationship between niche breadth and environmental stress: disk- or bowl-shaped blossoms (which allow a large number of potential pollinator species to access pollen and nectar rewards) dominated at high altitude flower communities.

Here, we investigate whether and how environmental stress can systematically affect specialisation. Our main aim is to test the two hypotheses mentioned above that relate environmental stress with a species' number of partners and investigate whether this changes systematically across species or between trophic guilds. We propose that specialist species can become “facultative” generalists to reduce their vulnerability to the absence of preferred partners (for example, when variations in climate decouple phenologies; Benadi *et al.* 2014). In other words, we expect that, as environmental stress increases, specialists should be more likely to engage with more partners. Species with many partners, conversely, should have a larger pool of available partners and might, therefore, be more likely to specialise on the most beneficial partners under environmental stress. Importantly, when testing these hypotheses, we control for the potential effects of the environment on community

composition (which has been previously shown to be a determinant factor; Gravel *et al.* 2018). We test these hypotheses using data on plant-pollinator interactions. These interactions provide a particularly interesting system to test these hypotheses because, due to the multiple trade-offs involved in the pollination service, there are multiple intuitive ways in which we could imagine species responding to environmental stress given the available partners. We estimate the stress species might experience a given community by calculating its bioclimatic suitability based on the species' patterns of global occurrence.

## Methods

We retrieved plant-pollinator networks from the Web of Life database (Fortuna *et al.* 2014). This database contains datasets originating from 58 studies published in the primary literature between 1923 and 2016. Calculating the environmental stress of species in their community and their potential partners required us to reduce both the taxonomic and distributional/location uncertainty. A critical step towards reducing this uncertainty is to ensure that the names used to identify species are valid and unambiguous, which in turn allows us to obtain further information from biological databases and accurately match species across studies. Therefore, our first step was to ensure consistent spelling and standardisation of species names and synonyms (see Supplementary Methods). The cleaning process resulted on a total of 2,555 plants and 8,406 pollinator species distributed across 73 locations around the globe (Figure 1 and S1).

After matching species across studies as accurately as possible, we performed two more steps. First, we calculated the environmental stress of species in their communities. Second, we relate each species' stress in its community with the number of partner species it has as a metric of its level of specialisation.

## Environmental stress display

We calculated the environmental stress of species in their communities. We assume that the stress a species experiences in a particular location is inversely related to the suitability of the average




Figure 1: Frequency distribution of the number of locations in which a species is present. The most common pollinator species was *Apis mellifera*, which was sampled at 42 locations, while the most common plant species was *Trifolium repens*, which was sampled at 11 locations.

environmental conditions in that place. As we aim to compare specialisation levels for different levels of environmental stress, we only calculate bioclimatic suitability for species that were present in at least two communities. To calculate the bioclimatic suitability of a species in a particular location, we used a niche-factor analysis (Hirzel *et al.* 2002; Broennimann *et al.* 2012). This approach is based on the probability density function of species distribution in an environmental variable space. Habitats are characterised by a collection of environmental variables. In a nutshell, those habitats in which the species occurs more often are deemed to be more suitable for the species than habitats in which the species has never been observed. As bioclimatic suitability is calculated in a scale from zero to one following the niche-factor analysis, for simplicity, we define environmental stress as one minus suitability.

The niche factor analysis requires two critical pieces of information. First, it requires information about the occurrences of the species of interest. Second, the method requires information about the environmental conditions for all the locations in which the species occurs. We retrieved 38.1 million occurrences of our focal species from the Global Biodiversity Information Facility (GBIF.org 2019a, b, c, d, e, f, g, h, i, j). Issues with data quality are a central issue hampering the use of publicly available species occurrence GBIF data in ecology and biogeography (Jetz *et al.* 2019). We, therefore, followed a series of filters and geographic heuristics to correct or remove erroneous and imprecise referencing records (see supplementary methods; Zizka *et al.* 2019) which allowed us to identify and remove 7.5 million potentially problematic occurrences from further analysis. We



integrated the occurrence records from our plant-pollinator communities to the cleaned occurrences retrieved from GBIF.

We retrieved environmental data from WorldClim V2.0, which includes 19 bioclimatic variables commonly used in species distribution modelling (Fick & Hijmans 2017). We then complemented data obtained from WorldClim with data from Envirem (Title & Bemmels 2017), which includes 16 extra bioclimatic and two topographic variables. The additional set of variables from Envirem are relevant to ecological or physiological processes and thus have the potential to improve our suitability estimation (Title & Bemmels 2018). We obtained all environmental data as rasters composed by cells of 2.5 arc-minutes. We chose this resolution because it provides a reasonable match to the locational accuracy of the species occurrences found in GBIF, particularly those that originate from preserved specimens in museum collections.

After obtaining information about species occurrence and the environment, we then merged these two datasets such that a vector with details of our 37 bioclimatic and topographic variables characterised the location of each occurrence. Sets of occurrence data tend to be spatially aggregated due to sample bias (tendency to collect close to cities, certain countries). Moreover, spatial autocorrelation arises in ecological data because geographically clumped records tend to be more similar in physical characteristics and/or species abundances than do pairs of locations that are farther apart. To account for such spatial dependency in occurrence data, we only included one occurrence record if a species had more than one within a cell of the bioclimatic raster. We did this to avoid giving more weight to areas with a high number of occurrences, a common scenario in occurrence records collected opportunistically as the ones we use here. In this step we removed 85.4% of the occurrences which resulted in a total of 4.5 million occurrences used in our niche analysis.

A common issue of terrestrial bioclimatic datasets is that the boundaries of the cells with information do not precisely match the landmass boundaries. The result of this mismatch is that not all environmental variables were available for 3,273 of the raster cells with occurrences (0.8% of the total). As expected, the vast majority of these problematic cells were close to the shore. To address this issue, we calculated the average value of environmental variables within a 5km buffer of the centre of the cell where the variable was missing and used it to approximate the value of the variable in that cell. Using this procedure, we were able to fill environmental variables for 89.3% of the cells

where they were missing. To fill the remaining 350 cells, we repeated the aforementioned procedure but instead using a 10km buffer. We removed from further analysis occurrences located within the 135 cells for which we were unable to fill environmental variables (0.03% of the total).

Next, we calculated the probability density function of the species distribution in environmental space. To determine the environmental space, we used the first two components from a principal component analysis of the 37 bioclimatic variables associated with the species occurrences. Specifically we used the `dudi.pca` function from the R package `ade4` 1.7.13 (Dray & Dufour 2007) and centered and scaled all bioclimatic variables to have a mean of zero and a unit variance. We then determined the position of species occurrences in the environmental space and estimated their bivariate probability density function. We used a kernel method to estimate this density and normalised it such that it ranges between zero and one. We used the kernel density method in the niche-factor analysis (Broennimann *et al.* 2012) rather than the distance from the mode (Hirzel *et al.* 2002) (as has been proposed earlier) because it has been shown to reduce the procedure’s sensitivity to sampling effort and the resolution of the environmental space. Specifically, to calculate the probability density function we used `ecospat.grid.clim.dyn` from the R package `ecospat` 3.0 (Broennimann *et al.* 2018) with a grid resolution of 200. We then determined the location in the environmental space of the plant-pollinator communities using the function `suprow` from `ade4`. The normalised density at that particular location (which we calculated using the R package `raster` 2.8.19; Hijmans 2019) corresponds with the bioclimatic suitability. The result of all these steps is the generation of a value of environmental stress, which corresponds to one minus the bioclimatic suitability for a species of a particular location.

We used a sensitivity analysis to determine the minimum number of occurrences that are necessary to have robust environmental stress estimations. For that we used the species with most occurrences available, *Archilochus colubris*, and calculated the mean absolute error of the bioclimatic suitability values obtained with one thousand subsamples from the 74,791 occurrences available from GBIF. This analysis indicated that, for a species, we need roughly 18 independent occurrences for each community for which we aim to estimate the environmental stress. This is the number of occurrences necessary to maintain the mean absolute error of bioclimatic suitability below 0.1 (Figure 2). We therefore removed from further analyses 286 species for which we did not have enough occurrences




Figure 2: Sensitivity analysis of environmental stress error. The number of independent occurrences retrieved from GBIF is inversely related to the error of bioclimatic suitability for our plant-pollinator networks. The sensitivity analysis was performed by subsampling occurrences of *Archilochus colubris*, the species in our dataset with the largest number of occurrences in GBIF, which was recorded in two of our communities.

to obtain robust estimates.

## Data analysis

We then used a set of Bayesian multilevel models to evaluate the impact of environmental stress on species specialisation. Specifically, we use the normalised degree of species as our response variable; that is, the number of species it interacts with given the number of species in the opposite guild (Martín González *et al.* 2010). The normalised degree was modelled using a logit link function and a binomial distribution in which the number of partner species a focal species interacts with is the number of successes, and the number of species in the opposite guild is the number of trials. We are aware that whether species interact or not is not a Bernoulli process as species interactions are not strictly independent from each other. However, the use of a binomial distribution allows us to account for the differences in species richness across communities indirectly. Importantly, results are qualitatively similar when we model species degree directly using a Poisson distribution and a logarithmic link function.

We evaluated four models to assess the relative importance of stress for each focal species.

A first model, our baseline model, included five variables. The predictors in the baseline model were the environmental stress of the community for each focal species, the species' number of

known possible partners in the community, and both the species guild (plant or a pollinator) and its interaction with environmental stress. We included the number of known possible partners as a predictor in our models as it allows us to control for the effects of the environment on community composition, effectively accounting for species co-occurrence. We calculated this metric by determining the number of partners with which the species is known to interact in any other community and assessing how many of these potential partners are present in the local community. Controlling for the number of potential partners makes our model a particularly stringent test of our environmental-stress hypotheses because this variable could explain a large proportion of variance. Often, the potential and the actual number of partners is the same or very close to each other, especially for rare species present only in a few communities.

We allowed the intercept and slope of the stress-specialisation relationship to vary among species. This approach allowed us to investigate two questions. First, it allows us to inspect the extent to which environmental stress affects species similarly. Second, by investigating the correlation between the intercept and the slope as a model parameter, it allowed us to inspect the extent by which species with a small or large number of partner species respond to increasing levels of environmental stress. To account for unmeasured differences between communities, like sampling effort, sampling method, or diversity, we also allowed the model intercept to be different for each community in our study. To facilitate model interpretation and convergence, we scaled all continuous variables to have a mean of zero and a unit variance.

We compared this baseline model with three alternative models in which we removed one predictor at a time. To quantify the difference between models, in terms of their expected out-of-sample performance, we use the Watanabe-Akaike information criterion (WAIC). All models were fitted under a Bayesian framework using the R package `brms` 2.8.0 (Bürkner 2017, 2018) as an interface for Stan (Carpenter *et al.* 2017). For each model, we used four Markov chains of 4,000 iterations each; we used half of the iterations for warmup. We used weakly informative priors for all model parameters. Specifically we used normal priors of mean zero and standard deviation ten for the population-level effects and the intercepts, a half-Cauchy prior with a location of zero and a scale of two for the standard deviations, and, when applicable, an LKJ-correlation prior with parameter  $\zeta = 1$  for the correlation matrix between group-level parameters.




Figure 3: Conditional effects of predictors in our baseline model. The shown values are based on predictions for a hypothetical community with 76 plants and 33 pollinators. These values correspond to the median number of species in each guild across communities. In each panel, we condition on the mean value of the other predictor in the model. We indicate mean values for each predictor with a vertical dashed line. For model fitting, we scaled all predictors to have a mean of zero and unit variance; however, here we show the unscaled predictors to facilitate interpretation. To illustrate the uncertainty around the fitted estimates, we plot the fits of 100 independent draws from the posterior distribution. The thick lines indicate the mean values of the response distribution. As there was no interaction between the guild and the number of possible interactions, we only show the conditional effect of pollinators.

## Results

Our models performed relatively well. The Bayesian  $R^2$  for our baseline model was 0.89, which indicates that our models were able to capture a large proportion of the variability on the data. Overall, we found that environmental stress does not have a consistent effect across species. Indeed, when looking at the fixed effects, stress has virtually no relationship with the normalised degree—our metric of specialisation (Figure 3a). However, environmental stress was still an important predictor in our model. The difference in WAIC between our baseline model and the model that did not include environmental stress was  $489 \pm 94$  (Table 1). This apparent discrepancy can be explained by the variability of the specialisation-stress relationship across species.

For some species, there is a strong negative relationship between stress and specialisation, while for others, there is a strong positive relationship (Figure 4a). Interestingly, the slope of this

Table 1: Comparison in out of sample predictive power of the baseline model (bold) and their alternatives. We rank models by their expected log predictive density based on their Wanatabe-Akaike information criterion (WAIC).

| predictors                                  | WAIC  | SE  |
|---|-------|-----|
| <b>stress x guild + # possible partners</b> | 6,592 | 170 |
| stress + # possible partners                | 6,595 | 166 |
| guild + # possible partners                 | 7,081 | 202 |
| stress x guild                              | 8,041 | 290 |

relationship was correlated with the species’ specialisation intercept in the model (Figure 4b and c; the mean correlation coefficient was 0.52 [0.33, 0.67]). This correlation indicates that the slope of the stress-specialisation relationship was more likely to be positive for species with many partners under average stress conditions (and negative for species fewer partners). Recall that the model estimates the intercept of the number of partners for each species at the mean value for stress across communities (0.68). Extrapolating to no-stress conditions this means that species that would interact with few partners under no stress are more likely to interact with more partners as stress increases, whereas those that would interact with many partners are more likely to interact fewer. As expected, we found a strong and positive relationship between the number of possible interactions and the number of realised interactions in the community (i.e. species that were capable of being generalists tended to realise this generality at all locations). There was also a large difference of WAIC between the model that included this predictor and that that excluded it. This result indicates that the availability of potential partners—this is, community composition—accounts for a large proportion of the variability in species degree. Importantly, our findings relating to the variability of the stress-specialisation relationship were qualitatively unchanged, whether we accounted for community composition or not.

The standard deviation (in the parameters scale) of the community intercepts was 1.02 [0.85, 1.23] which indicates the importance of the local context when determining specialisation. The standard deviation of the species intercept was 0.54 [0.48, 0.61], and that of the species’ stress slope was 0.38 [0.32, 0.44] (95% credible intervals shown within square brackets).




Figure 4: Species-level effects of environmental stress (a) Conditional effect of stress for individual species. Each line corresponds to the median relationship for each species. Although we included in the analysis all species that are present in two or more communities, to facilitate visualisation here, we show only species present in at least six communities (10 plants and 33 pollinators). As in the previous figure, fitted values assume a hypothetical community of median size. In each panel, we highlight two species for which the relationship between environmental suitability and the normalised degree was particularly strong. (b and c) There was a positive correlation between the species' intercept and the species' stress-specialisation slope. The species' intercept can be interpreted as the number of partners a species would have under mean levels of environmental stress.

## Discussion

We set out to explore whether and how environmental stress can systematically affect specialisation. After accounting for the pool of potential partners, we found that environmental conditions contribute to determining whether a species is a generalist or a specialist *in their community*. We also found that the particular effect of the environment depends strongly on the species' inherent tendency to be a specialist or generalist. Based on existing literature, we proposed two alternative hypotheses of how environmental stress may affect species' specialisation, and we found evidence for both. Species with many partners in low-stress communities were more likely to have a negative relationship and hence reduce the number of partners as stress increases. Contrastingly, species in our datasets with few partners in low-stress communities were more likely to interact with more partners in more stressful communities.

Our results suggest that changes in community composition are indeed the primary channel through which the environment determines changes interaction probability, and this is a common mechanism through which ecological networks respond to environmental change (Tylianakis & Morris 2017). However, they also show that, for a large number of species, the environment may also play a substantial role in determining their level of specialisation. Previous research has recognised that environmental factors may help explain the changes in network structure along environmental gradients that cannot be explained by community composition (Tylianakis *et al.* 2007). However, how these two factors were linked had been elusive so far (Gravel *et al.* 2018). We believe that part of this difficulty could have arisen because species, and ultimately network structure, can respond in multiple, and contrasting, ways depending on the particular bioclimatic variable examined (e.g. temperature or precipitation). Using stress to summarise the effect on species of multiple environmental gradients allowed us to detect a clear signal of the environment in species' interaction patterns.

Recent research suggests that species are continuously changing their interaction partners wherever environmental conditions change in space or time (Raimundo *et al.* 2018). So far it appears that this rewiring is primarily driven by generalist species (Burkle *et al.* 2013; Ponisio *et al.* 2017), presumably because generalist species are less sensitive to trait matching of their interaction partners



(CaraDonna *et al.* 2017). Our results add two important nuances to these findings. First, because “generalists” seem to focus on fewer partners as environmental conditions deteriorate, we show that trait matching might still play a role in determining the less-flexible core interactions of generalist species. Second, and most importantly, our results suggest that only a small proportion of species are “true generalists” or “true specialists” this is, species that interact with a large or small number of partners regardless of the environmental stress, respectively. This pattern implies that rewiring is not exclusive to species with many partners. Instead, at least a fraction of the species that appear to be specialist *in their communities* might be as flexible, if not more, than those with a large number of partners, effectively behaving as facultative generalists in the face of environmental change. These flexible “specialists” might therefore have a more significant role in network persistence than previously expected.

In our model, we can roughly divide species between true specialists, true generalists, and flexible species. However, there is a fourth group that remained invisible to our model but has important implications for network persistence and stability. Species that can vary their interaction partners flexibly and their role in the network are more likely to persist in their community as environmental conditions vary (Gaiarsa *et al.* 2019). We propose this fourth group is composed of true specialists that are constrained to interact with partners of high trait-matching and therefore were not likely to be found in more than one community. If species that are not flexible are unlikely to persist over temporal or spatial environmental gradients, we can expect specialised communities that are highly constrained by trait-matching (like some plant-hummingbird networks; Vizentin-Bugoni *et al.* 2014; Maruyama *et al.* 2014) to be far more vulnerable to increased climate change-induced environmental stress and habitat degradation than communities where role and interaction flexibility are more prevalent.

Similarly, if the patterns we see in our models have also played a role during the evolutionary history of pollination communities, our results also help to explain why only a small fraction of plant-pollinator interactions shows a strong signature of deep co-evolutionary history (Hutchinson *et al.* 2017). The increases in the stress that species are predicted to experience due to rapid environmental change might further erode the co-evolutionary history of specialist species. Communities as a whole might be in a trajectory of even more diffuse co-evolution. For specialists, at least, the

longer-term benefits of being able to interact with multiple partners might be more important than the shorter-term benefits of interacting with partners of high trait matching.

The structural implications of the “regression towards the mean” that environmental stress promotes are less clear. However, it is plausible to expect that nestedness, and therefore network stability, might be reduced in the face of rapid environmental change. Determining exactly how the changes in degree caused by environmental stress are reflected by systematic changes in network structure would be an interesting avenue of research. Answering this question would require expanding our suitability analysis to all species in the community and comparing the degree distribution of networks along a gradient of stress for the community as a whole.

In conclusion, we show that the environment can affect the specialisation level of plants and pollinators in systematic ways beyond community composition. Species that are inflexible with their interaction partners are unlikely to persist under more stressful environmental conditions. However, we show that many species are flexible in regards to their specialisation levels and therefore are not inherently generalists or specialists. Instead, the species’ level of specialisation/generalisation should be considered on a relative scale depending on environmental stress at a given location.

## Acknowledgements

We thank Warwick Allen, Marilia Gaiarsa, and Guadalupe Peralta for feedback and valuable discussions. EFC acknowledges the support from the University of Canterbury Doctoral Scholarship and a New Zealand International Doctoral Research Scholarship administered by New Zealand Education. DBS and JMT acknowledge the support of Rutherford Discovery Fellowships (RDF-13-UOC-003 and RDF-UOC-1002) and the Marsden Fund Council (UOC-1705), administered by the Royal Society of New Zealand Te Apārangi.

## References

Baskett, C.A. & Schemske, D.W. (2018). Latitudinal patterns of herbivore pressure in a temperate herb support the biotic interactions hypothesis. *Ecology Letters*, 21, 578–587.

351 Benadi, G., Hovestadt, T., Poethke, H.-J. & Blüthgen, N. (2014). Specialization and phenological  
352 synchrony of plant-pollinator interactions along an altitudinal gradient. *Journal of Animal Ecology*,  
353 83, 639–650.

354 Broennimann, O., Di Cola, V. & Guisan, A. (2018). *Ecospat: Spatial ecology miscellaneous methods*.  
355 Broennimann, O., Fitzpatrick, M.C., Pearman, P.B., Petitpierre, B., Pellissier, L. & Yoccoz, N.G.  
356 *et al.* (2012). Measuring ecological niche overlap from occurrence and spatial environmental data:  
357 Measuring niche overlap. *Global Ecology and Biogeography*, 21, 481–497.

358 Burkle, L.A., Marlin, J.C. & Knight, T.M. (2013). Plant-pollinator interactions over 120 years:  
359 Loss of species, co-occurrence, and function. *Science*, 339, 1611–1615.

360 Bürkner, P.-C. (2017). Brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of*  
361 *Statistical Software*, 80.

362 Bürkner, P.-C. (2018). Advanced Bayesian Multilevel Modeling with the R Package brms. *The R*  
363 *Journal*, 10, 395.

364 CaraDonna, P.J., Petry, W.K., Brennan, R.M., Cunningham, J.L., Bronstein, J.L. & Waser, N.M.  
365 *et al.* (2017). Interaction rewiring and the rapid turnover of plant-pollinator networks. *Ecology*  
366 *Letters*, 20, 385–394.

367 Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B. & Betancourt, M. *et al.* (2017).  
368 Stan: A Probabilistic Programming Language. *Journal of Statistical Software*, 76.

369 Cirtwill, A.R., Dalla Riva, G.V., Gaiarsa, M.P., Bimler, M.D., Cagua, E.F. & Coux, C. *et al.* (2018).  
370 A review of species role concepts in food webs. *Food Webs*, 16, e00093.

371 Devoto, M., Medan, D. & Montaldo, N.H. (2005). Patterns of interaction between plants and  
372 pollinators along an environmental gradient. *Oikos*, 109, 461–472.

373 Dray, S. & Dufour, A.-B. (2007). The ade4 package: Implementing the duality diagram for ecologists.  
374 *Journal of Statistical Software*, 22.

375 Dupont, Y.L., Padrón, B., Olesen, J.M. & Petanidou, T. (2009). Spatio-temporal variation in the  
376 structure of pollination networks. *Oikos*, 118, 1261–1269.

377 Fick, S.E. & Hijmans, R.J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for  
 378 global land areas: New climate surfaces for global land areas. *International Journal of Climatology*,  
 379 37, 4302–4315.

380 Fortuna, M.A., Ortega, R. & Bascompte, J. (2014). The Web of Life. *arXiv:1403.2575 [q-bio]*.

381 Gaiarsa, M.P., Kremen, C. & Ponisio, L.C. (2019). Interaction flexibility predicts pollinator  
 382 population dynamics. *Manuscript submitted for publication*, 15.

383 GBIF.org. (2019a). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 384 download/0004943-190621201848488](https://www.gbif.org/occurrence/download/0004943-190621201848488). Last accessed 2 July 2019.

385 GBIF.org. (2019b). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 386 download/0004939-190621201848488](https://www.gbif.org/occurrence/download/0004939-190621201848488). Last accessed 2 July 2019.

387 GBIF.org. (2019c). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 388 download/0004937-190621201848488](https://www.gbif.org/occurrence/download/0004937-190621201848488). Last accessed 2 July 2019.

389 GBIF.org. (2019d). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 390 download/0004935-190621201848488](https://www.gbif.org/occurrence/download/0004935-190621201848488). Last accessed 2 July 2019.

391 GBIF.org. (2019e). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 392 download/0004929-190621201848488](https://www.gbif.org/occurrence/download/0004929-190621201848488). Last accessed 2 July 2019.

393 GBIF.org. (2019f). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 394 download/0004928-190621201848488](https://www.gbif.org/occurrence/download/0004928-190621201848488). Last accessed 2 July 2019.

395 GBIF.org. (2019g). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 396 download/0004927-190621201848488](https://www.gbif.org/occurrence/download/0004927-190621201848488). Last accessed 2 July 2019.


397 GBIF.org. (2019h). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 398 download/0004918-190621201848488](https://www.gbif.org/occurrence/download/0004918-190621201848488). Last accessed 2 July 2019.

399 GBIF.org. (2019i). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 400 download/0004917-190621201848488](https://www.gbif.org/occurrence/download/0004917-190621201848488). Last accessed 2 July 2019.

401 GBIF.org. (2019j). *GBIF occurrence download*. Available at: [https://www.gbif.org/occurrence/  
 402 download/0004916-190621201848488](https://www.gbif.org/occurrence/download/0004916-190621201848488). Last accessed 2 July 2019.

403 Godsoe, W., Jankowski, J., Holt, R.D. & Gravel, D. (2017). Integrating Biogeography with  
 404 Contemporary Niche Theory. *Trends in Ecology & Evolution*, 32, 488–499.

405 Gravel, D., Baiser, B., Dunne, J.A., Kopelke, J.-P., Martinez, N.D. & Nyman, T. *et al.* (2018).  
 406 Bringing Elton and Grinnell together: A quantitative framework to represent the biogeography of  
 407 ecological interaction networks. *Ecography*, 41, 1–15.

408 Hijmans, R.J. (2019). *Raster: Geographic data analysis and modeling*. 

409 Hirzel, A.H., Hausser, J., Chessel, D. & Perrin, N. (2002). Ecological-niche factor analysis: How to  
 410 compute habitat-suitability maps without absence data?, 83, 10.

411 Hoiss, B., Krauss, J., Potts, S.G., Roberts, S. & Steffan-Dewenter, I. (2012). Altitude acts as  
 412 an environmental filter on phylogenetic composition, traits and diversity in bee communities.  
 413 *Proceedings of the Royal Society B: Biological Sciences*, 279, 4447–4456.

414 Hoiss, B., Krauss, J. & Steffan-Dewenter, I. (2015). Interactive effects of elevation, species richness  
 415 and extreme climatic events on plant-pollinator networks. *Global Change Biology*, 21, 4086–4097.

416 Hutchinson, M.C., Cagua, E.F. & Stouffer, D.B. (2017). Cophylogenetic signal is detectable in  
 417 pollination interactions across ecological scales. *Ecology*, 98, 2640–2652.

418 Jetz, W., McGeoch, M.A., Guralnick, R., Ferrier, S., Beck, J. & Costello, M.J. *et al.* (2019).  
 419 Essential biodiversity variables for mapping and monitoring species populations. *Nature Ecology &*  
 420 *Evolution*, 3, 539–551.

421 Laliberté, E. & Tylianakis, J.M. (2012). Cascading effects of long-term land-use changes on plant  
 422 traits and ecosystem functioning. *Ecology*, 93, 145–155.

423 Lavandero, B. & Tylianakis, J.M. (2013). Genotype matching in a parasitoid-host genotypic food  
 424 web: An approach for measuring effects of environmental change. *Molecular Ecology*, 22, 229–238.

425 Martín González, A.M., Dalsgaard, B. & Olesen, J.M. (2010). Centrality measures and the  
 426 importance of generalist species in pollination networks. *Ecological Complexity*, 7, 36–43.

427 Maruyama, P.K., Vizentin-Bugoni, J., Oliveira, G.M., Oliveira, P.E. & Dalsgaard, B. (2014).  
 428 Morphological and Spatio-Temporal Mismatches Shape a Neotropical Savanna Plant-Hummingbird

429 Network. *Biotropica*, 46, 740–747.

430 Olesen, J.M., Bascompte, J., Elberling, H. & Jordano, P. (2008). Temporal dynamics in a pollination  
431 network. *Ecology*, 89, 1573–1582.

432 Olesen, J.M., Stefanescu, C. & Traveset, A. (2011). Strong, Long-Term Temporal Dynamics of an  
433 Ecological Network. *PLoS ONE*, 6, e26455.

434 Pellissier, L., Pottier, J., Vittoz, P., Dubuis, A. & Guisan, A. (2010). Spatial pattern of floral  
435 morphology: Possible insight into the effects of pollinators on plant distributions. *Oikos*, 119,  
436 1805–1813.

437 Peralta, G., Frost, C.M., Didham, R.K., Varsani, A. & Tylianakis, J.M. (2015). Phylogenetic  
438 diversity and co-evolutionary signals among trophic levels change across a habitat edge. *Journal of*  
439 *Animal Ecology*, 84, 364–372.

440 Pires, A.P.F., Marino, N.A.C., Srivastava, D.S. & Farjalla, V.F. (2016). Predicted rainfall changes  
441 disrupt trophic interactions in a tropical aquatic ecosystem. *Ecology*, 97, 2750–2759.

442 Poisot, T., Stouffer, D.B. & Gravel, D. (2015). Beyond species: Why ecological interaction networks  
443 vary through space and time. *Oikos*, 124, 243–251.

444 Ponisio, L.C., Gaiarsa, M.P. & Kremen, C. (2017). Opportunistic attachment assembles plant-  
445 pollinator networks. *Ecology Letters*, 20, 1261–1272.

446 Rabeling, S.C., Lim, J.L., Tidon, R., Neff, J.L., Simpson, B.B. & Pawar, S. (2019). Seasonal variation  
447 of a plant-pollinator network in the Brazilian Cerrado: Implications for community structure and  
448 robustness. *PLOS ONE*, 14, e0224997.

449 Raimundo, R.L., Guimarães, P.R. & Evans, D.M. (2018). Adaptive Networks for Restoration  
450 Ecology. *Trends in Ecology & Evolution*, 33, 664–675.

451 Schleuning, M., Fründ, J., Klein, A.-M., Abrahamczyk, S., Alarcón, R. & Albrecht, M. *et al.* (2012).  
452 Specialization of Mutualistic Interaction Networks Decreases toward Tropical Latitudes. *Current*  
453 *Biology*, 22, 1925–1931.

454 Thebault, E. & Fontaine, C. (2010). Stability of Ecological Communities and the Architecture of

455 Mutualistic and Trophic Networks. *Science*, 329, 853–856.

456 Title, P.O. & Bemmels, J.B. (2017). ENVIREM: ENVironmental Rasters for Ecological Modeling  
457 version 1.0.

458 Title, P.O. & Bemmels, J.B. (2018). ENVIREM: An expanded set of bioclimatic and topographic  
459 variables increases flexibility and improves performance of ecological niche modeling. *Ecography*, 41,  
460 291–307.

461 Trøjelsgaard, K., Jordano, P., Carstensen, D.W. & Olesen, J.M. (2015). Geographical variation in  
462 mutualistic networks: Similarity, turnover and partner fidelity. *Proceedings of the Royal Society B:  
463 Biological Sciences*, 282, 20142925.

464 Tylianakis, J.M., Didham, R.K., Bascompte, J. & Wardle, D.A. (2008). Global change and species  
465 interactions in terrestrial ecosystems. *Ecology Letters*, 11, 1351–1363.

466 Tylianakis, J.M. & Morris, R.J. (2017). Ecological Networks Across Environmental Gradients.  
467 *Annual Review of Ecology, Evolution, and Systematics*, 48, 25–48.

468 Tylianakis, J.M., Tcharntke, T. & Lewis, O.T. (2007). Habitat modification alters the structure of  
469 tropical hostParasitoid food webs. *Nature*, 445, 202–205.

470 Vázquez, D.P., Blüthgen, N., Cagnolo, L. & Chacoff, N.P. (2009a). Uniting pattern and process in  
471 plantAnimal mutualistic networks: A review. *Annals of Botany*, 103, 1445–1457.

472 Vázquez, D.P., Chacoff, N.P. & Cagnolo, L. (2009b). Evaluating multiple determinants of the  
473 structure of plantAnimal mutualistic networks. *Ecology*, 90, 2039–2046.

474 Vizentin-Bugoni, J., Maruyama, P.K. & Sazima, M. (2014). Processes entangling interactions in  
475 communities: Forbidden links are more important than abundance in a hummingbirdPlant network.  
476 *Proceedings of the Royal Society B: Biological Sciences*, 281, 20132397.

477 Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C. & Edler, D. *et al.* (2019).  
478 CoordinateCleaner: Standardized cleaning of occurrence records from biological collection databases.  
479 *Methods in Ecology and Evolution*, 10, 744–751.

# Environment affects specialisation of plants and pollinators

Supplementary information

*E. Fernando Cagua, Audrey Lustig, Jason M. Tylianakis, Daniel B. Stouffer*

## Supplementary methods

### Reducing taxonomic uncertainty

Data were obtained from the Web of Life database (Fortuna, Ortega, and Bascompte 2014) which includes data from 58 published studies (Abreu and Vieira 2004; Arroyo, Primack, and Armesto 1982; Barrett and Helenurm 1987; Bartomeus, Vilà, and Santamaría 2008; Bek 2006; Bezerra, Machado, and Mello 2009; Bundgaard 2003; Canela 2006; Clements and Long 1923; del Coro Arizmendi and Ornelas 1990; Dicks, Corbet, and Pywell 2002; Dupont and Olesen 2009; Dupont, Hansen, and Olesen 2003; Elberling and Olesen 1999; Gutierrez, Rojas-Nossa, and Stiles 2004; Hattersley-Smith 1985; Herrera 1988; Hocking 1968; Ingversen 2006; INouE et al. 1990; Inouye and Pyke 1988; Kaiser-Bunbury et al. 2014, 2010; Kakutani et al. 1990; Kato 2000; Kato, Matsumoto, and Kato 1993; Kato and Miura 1996; Kato et al. 1990; Kevan 1970; Kohler 2011; Lara 2006; Las-Casas, Azevedo Júnior, and Dias Filho 2012; Lundgren and Olesen 2005; McMullen 1993; Medan et al. 2002; Memmott 1999; Montero 2005; Mosquin 1967; Motten 1986; Olesen, Eskildsen, and Venkatasamy 2002; Ollerton 2003; Percival 1974; Petanidou and Vokou 1993; Philipp et al. 2006; Primack 1983; Ramirez 1989; Ramirez and Brito 1992; Robertson 1929; Rosero and others 2003; Sabatino 2010; Schemske et al. 1978; Small 1976; Smith-Ramírez et al. 2005; Stald, Valido, and Olesen 2003; Vázquez 2002; Vizentin-Bugoni et al. 2016; Witt 1998; Yamazaki and Kato 2003).

Interaction data from the included studies included 11,231 unique organism names. From these, 1,166 were present in more than one study. From the total number of organisms, 159 were identified to the subspecies or variety level, 6,759 to the species level, 1,755 to the genus level, whereas the remaining 2,558 were unidentified. As the species level was the most common taxonomic rank available in our interaction datasets, in all further analysis, we grouped together subspecies or varieties within the same species.



We were able to confirm the validity of 5,263 of the scientific names used to identify organisms (roughly 76%). We assessed the validity of a name by querying the Global Names Resolver database (<https://resolver.globalnames.org>) which includes data from 98 taxonomic sources. We accessed this database using the function `gnr_resolve` from the R package `taxize` 0.9.6 (Chamberlain and Szocs 2013; S. Chamberlain, Szocs, et al. 2019).

From the remaining 1,655 names we were unable to validate, we were able to identify and correct 726 that contained spelling mistakes. These spelling mistakes were corrected automatically by fuzzy matching the canonical names in our data sources with those in the Global Names Resolver database. However, on rare occasions, the fuzzy matching algorithm can suggest a scientific name that has a similar spelling, but that corresponds to an organism in a different taxonomic group, often a separate kingdom. To address this potential problem, we checked the taxonomic hierarchy of suggested names and confirmed that it matched our expected taxon. We retrieved all taxonomic hierarchies from the National Center for Biotechnology Information taxonomic database (<https://www.ncbi.nlm.nih.gov/taxonomy>).

As species names are constantly changing, we subsequently checked for possible synonyms of the canonical names in our data sources. Using data from the Integrated Taxonomic Information System database (<http://www.itis.gov>), we found synonyms and alternative names for 611 species.

Finding these alternative names was required for two main reasons. First, because we wanted to be able to identify the cases in which the same species might have been recorded with different names in various data sources. This can occur not only when the canonical name has been changed but also when there are widely used orthographic variants. Second, because retrieving occurrence data is often only possible using the latest accepted/valid name for a particular species.

All together, from the 1,655 names we were unable to validate, it was not possible to automatically correct or find synonyms for 332 of them. We then manually consulted multiple online databases, chiefly Wikispecies (<https://species.wikimedia.org/>), and looked for canonical names that both, resembled the unvalidated names and matched the geographic and taxonomic expectations. In this fashion, we were able to further correct 25 names. Most manual corrections were made on names that have been abbreviated or had more than two spelling mistakes. A complete list of manual name corrections can be seen in Table S1.

This cleaning process allowed us to match a further 270 names across data

Table S1: Manually corrected canonical names. More than one correct name have been included when an accepted/valid synonym the canonical name exists.

| incorrect name               | corrected name                     | guild      |
|------------------------------|------------------------------------|------------|
| Acaena pinn                  | <i>Acaena pinnatifida</i>          | plant      |
| Adesmia brachy               | <i>Adesmia brachysemeon</i>        | plant      |
| Aesculus camea               | <i>Aesculus X carnea</i>           | plant      |
| Brachyome sinclairii         | <i>Brachyscome sinclairii</i>      | plant      |
| Calceolaria arac             | <i>Calceolaria arachnoidea</i>     | plant      |
| Equium sabulicola            | <i>Echium sabulicola</i>           | plant      |
| Euonymus fo rtunei           | <i>Euonymus fortunei</i>           | plant      |
| Galvezia leucantha pubescen  | <i>Galvezia leucantha</i>          | plant      |
| Heliconia simulans           | <i>Heliconia angusta</i>           | plant      |
| Pitcaimia flammea            | <i>Pitcairnia flammea</i>          | plant      |
| Psittacanthus flavo viridis  | <i>Psittacanthus flavo-viridis</i> | plant      |
| Rodophiala bifidum           | <i>Rhodophiala bifida</i>          | plant      |
| Stachys albi                 | <i>Stachys albicaulis</i>          | plant      |
| Stenactis annuus             | <i>Erigeron annuus</i>             | plant      |
| Thaspium aureum atropurpurem | <i>Thaspium trifoliatum</i>        | plant      |
| Tristhema mauritiana         | <i>Tristemma mauritianum</i>       | plant      |
| Tropaeolum polyph            | <i>Tropaeolum polyphyllum</i>      | plant      |
| Tyttnera scabra              | <i>Turnera scabra</i>              | plant      |
|                              | <i>Turnera ulmifolia</i>           | plant      |
| VVedelia biflora             | <i>Melanthera biflora</i>          | plant      |
|                              | <i>Wedelia biflora</i>             | plant      |
| Cateres pennatus             | <i>Kateretes pennatus</i>          | pollinator |
| Eclimus harrisi              | <i>Condyllostylus crinicauda</i>   | pollinator |
| Ptilandrena g. maculati      | <i>Andrena distans</i>             | pollinator |
| Tapinotaspis caerulea        | <i>Chalepogenus caeruleus</i>      | pollinator |
| Tapinotaspis herbsti         | <i>Chalepogenus herbsti</i>        | pollinator |

sources and, by doing so, identify another 72 species that were present in more than one study. The process also allowed us to identify problematic data sources in which some names were included as both plants and pollinators. These data sources were removed from further analysis. In seven of our data sources, interaction data were recorded at multiple points in time. When this was the case, we combined interaction data into one single interaction network.

## Reducing location uncertainty

We retrieved occurrences from the Global Biodiversity Information Facility (GBIF; <https://www.gbif.org>) using the R package `rgbif` 0.9.6 (Chamberlain and Boettiger 2017; S. Chamberlain, Barve, et al. 2019). Specifically, for each species, we only requested occurrences for which the coordinates of the observation were available and that had no known geospatial issue in the GBIF database. We downloaded roughly 38.1 million occurrences for the 986 species in which we were interested. These occurrences, however, contain observations of mixed quality. Therefore, we followed Zizka et al. (2019) and




Figure S1: Worldwide distribution of pollination communities included in this study

applied a series of simple filters and geographic heuristics to remove those of lower quality. Specifically, we removed all occurrences with (i) a coordinate uncertainty larger than 100km; (ii) those recorded prior to 1945 (as records prior to this date have been shown to be often imprecise); (iii) those in which the number of counts in the occurrence was registered as zero (as that indicates that the species has not been recorded); and (iv) those occurrences in which the “basis of record” was not a human observation or a preserved specimen (as occurrences from unknown and fossil records are known to be highly unreliable). We then used the R package **CoordinateCleaner** 0.9.6 (Zizka et al. 2019) and land mass and country data from Natural Earth (<https://www.naturalearthdata.com>) with a 1:10,000,000 scale to further identify and remove problematic occurrences. We removed occurrences for which their coordinates (v) fell outside the borders of the country where they were recorded; (vi) those around a country capital or the centroid of the country and province centroids; (vii) those around a biodiversity institution; and (viii) those located within oceans. Through this cleaning process, we removed with 7.5 million occurrences distributed across 916 species.

## Supplementary references

Abreu, Carlos R M, and Milene F Vieira. 2004. “Os beija-flores e seus recursos florais em um fragmento florestal de Viçosa, sudeste brasileiro.” *Lundiana* 5 (2): 129–34.

Arroyo, Mary T. Kalin, Richard Primack, and Juan Armesto. 1982. “Community Studies in Pollination Ecology in the High Temperate Andes of Central Chile. I. Pollination Mechanisms and Altitudinal Variation.” *American Journal of Botany* 69 (1): 82–97. <https://doi.org/10.1002/j.1537-2197.1982.tb13237.x>.




Figure S2: Median habitat suitability of communities in our dataset. Each row represents a different community and horizontal lines represent span the 2.5 and 97.5 quantiles.

Barrett, Spencer C. H., and Kaius Helenurm. 1987. “The Reproductive Biology of Boreal Forest Herbs. I. Breeding Systems and Pollination.” *Canadian Journal of Botany* 65 (10): 2036–46. <https://doi.org/10.1139/b87-278>.

Bartomeus, Ignasi, Montserrat Vilà, and Luís Santamaría. 2008. “Contrasting Effects of Invasive Plants in Plant-Pollinator Networks.” *Oecologia* 155 (4): 761–70. <https://doi.org/10.1007/s00442-007-0946-1>.

Bek, S. 2006. “A Pollination Network from a Danish Forest Meadow.” MSc Thesis, Aarhus, Denmark: Aarhus University.

Bezerra, Elisângela L.S., Isabel C. Machado, and Marco A. R. Mello. 2009. “Pollination Networks of Oil-Flowers: A Tiny World Within the Smallest of All Worlds: Pollination Networks of Oil-Flowers.” *Journal of Animal Ecology* 78 (5): 1096–1101. <https://doi.org/10.1111/j.1365-2656.2009.01567.x>.

Bundgaard, M. 2003. “Tidslig Og Rumlig Variation I et Plante-Bestøvernetværk.” MSc Thesis, Aarhus, Denmark: Aarhus University.

Canela, Maria Bernadete. 2006. “Interações Entre Plantas E Beija-Flores Numa Comunidade de Floresta Atlântica Montana Em Itatiaia, Rio de Janeiro.” PhD Thesis, Campinas, Brazil: Universidade Estadual de Campinas.

Chamberlain, Scott A, and Carl Boettiger. 2017. “R Python, and Ruby

Clients for GBIF Species Occurrence Data.” Preprint. PeerJ Preprints. <https://doi.org/10.7287/peerj.preprints.3304v1>.

Chamberlain, Scott, Vijay Barve, Dan Mcglinn, Damiano Oldoni, Peter Desmet, Laurens Geffert, and Karthik Ram. 2019. *Rgbif: Interface to the Global Biodiversity Information Facility API*.

Chamberlain, Scott, and Eduard Szocs. 2013. “Taxize - Taxonomic Search and Retrieval in R.” *F1000Research*.

Chamberlain, Scott, Eduard Szocs, Zachary Foster, Zebulun Arendsee, Carl Boettiger, Karthik Ram, Ignasi Bartomeus, et al. 2019. *Taxize: Taxonomic Information from Around the Web*.

Clements, Frederic Edward, and Frances Louise Long. 1923. *Experimental Pollination: An Outline of the Ecology of Flowers and Insects*. 336. Carnegie Institution of Washington.

del Coro Arizmendi, Ma., and Juan Francisco Ornelas. 1990. “Hummingbirds and Their Floral Resources in a Tropical Dry Forest in Mexico.” *Biotropica* 22 (2): 172. <https://doi.org/10.2307/2388410>.

Dicks, L. V., S. A. Corbet, and R. F. Pywell. 2002. “Compartmentalization in Plant-Insect Flower Visitor Webs.” *Journal of Animal Ecology* 71 (1): 32–43. <https://doi.org/10.1046/j.0021-8790.2001.00572.x>.

Dupont, Yoko L., Dennis M. Hansen, and Jens M. Olesen. 2003. “Structure of a Plant-Flower-Visitor Network in the High-Altitude Sub-Alpine Desert of Tenerife, Canary Islands.” *Ecography* 26 (3): 301–10. <https://doi.org/10.1034/j.1600-0587.2003.03443.x>.

Dupont, Yoko L., and Jens M. Olesen. 2009. “Ecological Modules and Roles of Species in Heathland Plant-Insect Flower Visitor Networks.” *Journal of Animal Ecology* 78 (2): 346–53. <https://doi.org/10.1111/j.1365-2656.2008.01501.x>.

Elberling, Heidi, and Jens M. Olesen. 1999. “The Structure of a High Latitude Plant-Flower Visitor System: The Dominance of Flies.” *Ecography* 22 (3): 314–23. <https://doi.org/10.1111/j.1600-0587.1999.tb00507.x>.

Fortuna, Miguel A., Raul Ortega, and Jordi Bascompte. 2014. “The Web of Life.” *arXiv:1403.2575 [Q-Bio]*, March. <http://arxiv.org/abs/1403.2575>.

Gutierrez, Aquiles, Sandra Victoria Rojas-Nossa, and Gary F Stiles. 2004. “Dinámica Anual Del a Interacción Colibrí-Flor En Ecosistemas Altoandinos.” *Ornitologia Neotropical* 15: 205–13.

- Hattersley-Smith, G. 1985. "Botanical Studies in the Lake Hazen Region, Northern Ellesmere Island, Northwest Territories, Canada, by James H. Soper and John M. Powell." *ARCTIC* 38 (4): 346. <https://doi.org/10.14430/arctic2421>.
- Herrera, Javier. 1988. "Pollination Relationships in Southern Spanish Mediterranean Shrublands." *The Journal of Ecology* 76 (1): 274. <https://doi.org/10.2307/2260469>.
- Hocking, Brian. 1968. "Insect-Flower Associations in the High Arctic with Special Reference to Nectar." *Oikos* 19 (2): 359. <https://doi.org/10.2307/3565022>.
- Ingversen, TT. 2006. "Plant-Pollinator Interactions on Jamaica and Dominica: The Centrality, Asymmetry and Modularity of Networks." MSc Thesis, Aarhus, Denmark: University of Aarhus.
- INouE, Tamiji, Makoto KATo, Takehiko KAKuTANi, Takeshi SuKA, and Takao It. 1990. "Of Kibune, Kyoto: An Overview of the Flowering Phenology and the Seasonal Pattern of Insect Visits'," 88.
- Inouye, David W., and Graham H. Pyke. 1988. "Pollination Biology in the Snowy Mountains of Australia: Comparisons with Montane Colorado, USA." *Austral Ecology* 13 (2): 191–205. <https://doi.org/10.1111/j.1442-9993.1988.tb00968.x>.
- Kaiser-Bunbury, Christopher N., Stefanie Muff, Jane Memmott, Christine B. Müller, and Amedeo Calfisch. 2010. "The Robustness of Pollination Networks to the Loss of Species and Interactions: A Quantitative Approach Incorporating Pollinator Behaviour." *Ecology Letters* 13 (4): 442–52. <https://doi.org/10.1111/j.1461-0248.2009.01437.x>.
- Kaiser-Bunbury, Christopher N., Diego P. Vázquez, Martina Stang, and Jaboury Ghazoul. 2014. "Determinants of the Microstructure of PlantPollinator Networks." *Ecology* 95 (12): 3314–24. <https://doi.org/10.1890/14-0024.1>.
- Kakutani, Takehiko, Tamiji Inoue, Makoto Kato, and Hidekuji Ichihashi. 1990. "Insect-Flower Relationship in the Campus of Kyoto University, Kyoto : An Overview of the Flowering Phenology and the Seasonal Pattern of Insect Visits." *Contributions from the Biological Laboratory, Kyoto University* 27 (4): 465–522.
- Kato, Makoto. 2000. "Anthophilous Insect Community and Plant-Pollinator Interactions on Amami Islands in the Ryukyu Archipelago, Japan," 101.

Kato, Makoto, Takehiko Kakutani, Tamiji Inoue, and Takao Itino. 1990. "Insect-Flower Relationship in the Primary Beech Forest of Ashu, Kyoto: An Overview of the Flowering Phenology and the Seasonal Pattern of Insect Visits," 68.

Kato, Makoto, Masamichi Matsumoto, and Toru Kato. 1993. "Flowering Phenology and Anthophilous Insect Community in the Cool-Temperate Subalpine Forests and Meadows at Mt. Kushigata in the Central Part of Japan," 58.

Kato, Makoto, and Reiichi Miura. 1996. "Flowering Phenology and Anthophilous Insect Community at a Threatened Natural Lowland Marsh at Nakaikemi in Tsuruga, Japan." *Contributions from the Biological Laboratory, Kyoto University* 29 (1): 1.

Kevan, PG. 1970. "High Arctic Insect-Flower Relations: The Interrelationships of Arthropods and Flowers at Lake Hazen, Ellesmere Island, Northwest Territories, Canada." PhD Thesis, Edmonton, Canada: University of Alberta.

Kohler, Glauco Ubiratan. 2011. "Redes de Interação Planta Beija-Flor Em Um Gradiente Altitudinal de Floresta Atlântica No Sul Do Brasil." MSc Thesis, Curitiba, Brasil: Universidade Federal do Paraná.

Lara, Carlos. 2006. "Temporal Dynamics of Flower Use by Hummingbirds in a Highland Temperate Forest in Mexico." *Ecoscience* 13 (1): 23–29. [https://doi.org/10.2980/1195-6860\(2006\)13\[23:TDOFUB\]2.0.CO;2](https://doi.org/10.2980/1195-6860(2006)13[23:TDOFUB]2.0.CO;2).

Las-Casas, Fmg, Sm Azevedo Júnior, and Mm Dias Filho. 2012. "The Community of Hummingbirds (Aves: Trochilidae) and the Assemblage of Flowers in a Caatinga Vegetation." *Brazilian Journal of Biology* 72 (1): 51–58. <https://doi.org/10.1590/S1519-69842012000100006>.

Lundgren, Rebekka, and Jens M. Olesen. 2005. "The Dense and Highly Connected World of Greenland's Plants and Their Pollinators." *Arctic, Antarctic, and Alpine Research* 37 (4): 514–20. [https://doi.org/10.1657/1523-0430\(2005\)037\[0514:TDAHWC\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0514:TDAHWC]2.0.CO;2).

McMullen, C. K. 1993. "Flower-Visiting Insects of the Galapagos Islands." *The Pan-Pacific Entomologist* 69 (1): 95.

Medan, Diego, Norberto H. Montaldo, Mariano Devoto, Anita Mantese, Viviana Vasellati, German G. Roitman, and Norberto H. Bartoloni. 2002. "Plant-Pollinator Relationships at Two Altitudes in the Andes of Mendoza, Argentina." *Arctic, Antarctic, and Alpine Research* 34 (3): 233. <https://doi.org/10.2307/1552480>.

- Memmott, Jane. 1999. "The Structure of a Plant-Pollinator Food Web." *Ecology Letters* 2 (5): 276–80. <https://doi.org/10.1046/j.1461-0248.1999.00087.x>.
- Montero, AC. 2005. "The Ecology of Three Pollination Networks." MSc Thesis, Aarhus, Denmark: Aarhus University.
- Mosquin, Theodore. 1967. "Observations on the Pollination Biology of Plants on Melville Island, NWT." *Can. Fld Nat.* 81: 201–5.
- Motten, Alexander F. 1986. "Pollination Ecology of the Spring Wildflower Community of a Temperate Deciduous Forest." *Ecological Monographs* 56 (1): 21–42. <https://doi.org/10.2307/2937269>.
- Olesen, Jens M., Louise I. Eskildsen, and Shadila Venkatasamy. 2002. "Invasion of Pollination Networks on Oceanic Islands: Importance of Invader Complexes and Endemic Super Generalists." *Diversity and Distributions* 8 (3): 181–92. <https://doi.org/10.1046/j.1472-4642.2002.00148.x>.
- Ollerton, J. 2003. "The Pollination Ecology of an Assemblage of Grassland Asclepiads in South Africa." *Annals of Botany* 92 (6): 807–34. <https://doi.org/10.1093/aob/mcg206>.
- Percival, Mary. 1974. "Floral Ecology of Coastal Scrub in Southeast Jamaica." *Biotropica* 6 (2): 104. <https://doi.org/10.2307/2989824>.
- Petanidou, Theodora, and Despina Vokou. 1993. "Pollination Ecology of Labiatae in a Phryganeic (East Mediterranean) Ecosystem." *American Journal of Botany* 80 (8): 892–99. <https://doi.org/10.1002/j.1537-2197.1993.tb15310.x>.
- Philipp, Marianne, Jens Böcher, Hans R. Siegmund, and Lene R. Nielsen. 2006. "Structure of a Plant-Pollinator Network on a Pahoehoe Lava Desert of the Galápagos Islands." *Ecography* 29 (4): 531–40. <https://doi.org/10.1111/j.0906-7590.2006.04546.x>.
- Primack, Richard B. 1983. "Insect Pollination in the New Zealand Mountain Flora." *New Zealand Journal of Botany* 21 (3): 317–33. <https://doi.org/10.1080/0028825X.1983.10428561>.
- Ramirez, Nelson. 1989. "Biología de Polinización En Una Comunidad Arbustiva Tropical de La Alta Guayana Venezolana." *Biotropica* 21 (4): 319. <https://doi.org/10.2307/2388282>.
- Ramirez, Nelson, and Ysaly Brito. 1992. "Pollination Biology in a Palm Swamp Community in the Venezuelan Central Plains." *Botanical Journal of*



*the Linnean Society* 110 (4): 277–302. <https://doi.org/10.1111/j.1095-8339.1992.tb00294.x>.

Robertson, Charles. 1929. *Flowers and Insects Lists of Visitors of Four Hundred and Fifty Three Flowers*. Cairnville: The Science Press Printing Company.

Rosero, Liliana, and others. 2003. “Interações Planta/Beija-Flor Em Três Comunidades Vegetais Da Parte Sul Do Parque Nacional Natural Chiribiquete, Amazonas (Colômbia).” PhD Thesis, Campinas, Brazil: Universidade Estadual de Campinas.

Sabatino, Malena. 2010. “Direct Effects of Habitat Area on Interaction Diversity in Pollination Webs.” *Ecological Applications* 20 (6): 7.

Schemske, Douglas W., Mary F. Willson, Michael N. Melampy, Linda J. Miller, Louis Verner, Kathleen M. Schemske, and Louis B. Best. 1978. “Flowering Ecology of Some Spring Woodland Herbs.” *Ecology* 59 (2): 351–66. <https://doi.org/10.2307/1936379>.

Small, Ernest. 1976. “Insect Pollinators of the Mer Bleue Peat Bog of Ottawa.” *Canadian Field-Naturalist*.

Smith-Ramírez, C., P. Martinez, M. Nuñez, C. González, and J. J. Armesto. 2005. “Diversity, Flower Visitation Frequency and Generalism of Pollinators in Temperate Rain Forests of Chiloé Island, Chile.” *Botanical Journal of the Linnean Society* 147 (4): 399–416. <https://doi.org/10.1111/j.1095-8339.2005.00388.x>.

Stald, L, A Valido, and Jens M. Olesen. 2003. “Struktur Og Dynamik I Rum Og Tid at et Bestøvningsnetværk På a Tenerife, De Kanariske Øer.” MSc-Thesis, Aarhus, Denmark: University of Aarhus.

Vázquez, Diego P. 2002. “Interactions Among Introduced Ungulates, Plants, and Pollinators: A Field Study in the Temperate Forest of the Southern Andes.” PhD Thesis, Knoxville, United States of America: University of Tennessee.

Vizentin-Bugoni, Jeferson, Pietro Kiyoshi Maruyama, Vanderlei J. Debastiani, L. da S. Duarte, Bo Dalsgaard, and Marlies Sazima. 2016. “Influences of Sampling Effort on Detected Patterns and Structuring Processes of a Neotropical Plant-Hummingbird Network.” Edited by Daniel B. Stouffer. *Journal of Animal Ecology* 85 (1): 262–72. <https://doi.org/10.1111/1365-2656.12459>.

Witt, P. 1998. “BSc Thesis.” BSc Thesis, Aarhus, Denmark: University of

Aarhus.

Yamazaki, Kyoko, and Makoto Kato. 2003. “Flowering Phenology and Anthophilous Insect Community,” 67.

Zizka, Alexander, Daniele Silvestro, Tobias Andermann, Josué Azevedo, Camila Duarte Ritter, Daniel Edler, Harith Farooq, et al. 2019. “CoordinateCleaner: Standardized Cleaning of Occurrence Records from Biological Collection Databases.” Edited by Tiago Quental. *Methods in Ecology and Evolution* 10 (5): 744–51. <https://doi.org/10.1111/2041-210X.13152>.