

Doctoral Thesis Proposal

# BUILDING RESILIENCE: HOW DO SPECIES INTERACTIONS SHAPE ECOSYSTEM COLLAPSE?

Fernando Cagua

## Summary

Natural ecosystems provide important services—like food and water—we humans depend on to a large extent. Much like the failure of a single key financial institution can trigger unexpected crashes on the stock market, human pressures—such as biological invasions and species extinctions—can cause sudden collapses that severely transform the way ecosystems function. However, despite its importance, we do not completely understand the dynamics that make ecosystems resilient to collapse. Because the functioning of ecosystems is largely determined by the network of interactions between the species that inhabit them, my proposed research aims to quantify the role played by species interactions in determining the resilience of ecosystems. To achieve this, I will focus on networks of mutually beneficial interactions, like those between plants and their pollinators, and use a combination of empirical data, computer simulations and ecological theory. Ultimately I want to understand why, when and how ecosystem collapses occur, and how to recover from them.

# Introduction

From food and freshwater production, to recreation and carbon sequestration, ecosystems provide a wide range of services of considerable value to humans. Unfortunately, the frequency of undesired ecosystem collapses—like the (often sudden) shift from a transparent to a turbid lake or from a self-sustaining fishery to a collapsed one—is dramatically increasing worldwide<sup>1</sup>. When ecosystem resilience is limited, breakdowns are more likely, and their ability to provide those services we depend on is endangered. Therefore, a necessary step to anticipate, prevent, and reverse ecosystem collapse is to understand the processes that support or undermine ecosystem resilience<sup>2,3</sup>.

Resilience is related to the amount of disturbance that an ecosystem could withstand without collapsing, or, in ecological jargon, undergoing a regime shift—a large, persistent transformation in ecosystem functioning and structure<sup>4,5</sup>. Moreover, substantial research indicates that ecosystem functioning and structure are largely determined by the network of interactions formed by species in an ecological community<sup>3,6–8</sup>. Since these factors ultimately determine the ecosystem response to disturbances, **the overall objective of my proposed research is to quantify the role played by species interactions in modulating ecosystem resilience.**

To do so, I will focus on the network of mutualistic interactions between plants and pollinators<sup>6,9,10</sup>. These networks, which form the base of pollination systems, play a globally important role in the maintenance of biodiversity and crop production<sup>9,10</sup>. Pollination systems are locally important too; for instance two thirds of New Zealand plants are pollinated by birds or insects<sup>11</sup>, and this includes iconic native plants (like kōwhai and pōhutukawa), and economically important crops (like kiwifruit, apples and grapes). However, pollination systems worldwide are currently being disrupted by multiple drivers of human-driven global change<sup>11</sup>.

Pollination is being affected particularly affected by the simultaneous loss of previously important species, and the introduction of invasive species. My thesis will concentrate on the resistance and resilience of pollination systems to biotic invasions and defaunation—top components of human-caused global change. These two drivers have affected New Zealand with particular intensity, at the same time it remains notably vulnerable<sup>12</sup>. For instance, 50% of plant species in New Zealand are introduced<sup>13</sup>, and imported social bees are now an important component of pollinator fauna<sup>14,15</sup>. Moreover, the depletion of native birds<sup>16,17</sup> is a prime example of how pollination systems in New Zealand are losing key pollinators, plants and habitats<sup>11</sup>.

In the first chapter of my thesis I will concentrate on biotic invasions. Specifically I have a twofold objective **first, I aim to determine which network characteristics shape the its resistance and resilience to invasions**, and second **to determine how biotic invasions, by affecting existing interactions in the community, reshape network resistance and resilience**. Empirical, time continuous observations of ecosystem dynamics in networks that have been subjected to invasions are limited. Therefore, I will use computer simulated communities to estimate the population dynamics of the species in the community, and then I will directly quantify stability properties from fluctuations in the species populations<sup>18,19</sup>. Specifically I will simulate communities with different network structures, and analyze which structures are more favorable for the coexistence of the invasive species to co-exist in the community. I will then in turn, see how the structure itself is changed by the invasive species, and whether the change in

structure has stability implications.

However, it has been shown that biotic invasions often occur in ecosystems that have already been degraded by species removal<sup>20</sup>. Understanding how invasions interact with defaunation in ecological networks is a global research priority, and essential for conserving, restoring, and managing New Zealand ecosystems<sup>15</sup>. In the second chapter, I will follow **by determining the compound effect of biodiversity loss and invasive species on the resilience of an ecosystem**. I will be extending the models developed in the first chapter, but focusing on how the loss of species—particularly those with a unique role in the ecosystem—make the ecosystem more or less susceptible to invasions. Simultaneously, the population models will also serve to determine the the patterns of species extinctions that follows after a successful species invasion.

The first two chapters of my thesis are designed to answer underlying questions of resilience theory, however the underlying aim of my third chapter is to translate the gained insight into useful lessons for ecosystem management. Ecosystems are complex, non-linear systems that are very difficult to control. However, recent work in theoretical physics has highlighted that is indeed possible to regulate them using targeted interventions<sup>21</sup>. I propose to build upon these findings to **determine the optimal set of management actions—from both a theoretical and a feasibility perspective—that are required to modify an ecosystem state**. For that, using a previously collected empirical dataset that characterized the network of interactions before and after an ecological invasion<sup>22</sup>, I will first find the set of species that can act as drivers of ecosystem state. Then, using an extension of the population model developed in the first chapter, I aim to determine the characteristics—like the degree of generalization of trophic position—that make an species more likely to serve as a driver of ecosystem state. Rescuing ecosystems from the brink of collapse and recovering them from undesired shifts is a major goal in conservation science. Finding a way in which actions targeted to specific species can maximize the ecosystem resilience, will bring us much closer to that goal.

In a world of constant change, building resilience is our best insurance against losing the ecosystem services we value and depend on. Although we have identified some of the pervasive effects of invasions and defaunation at the species level, we currently do not understand how the changes on ecosystem dynamics is affecting the resilience of the ecosystems as a whole. The research I propose intends to establish the general theory necessary to answer this question. Doing so is especially important when the ecosystem response might be inconspicuous until transformation is imminent. Only by better understanding the dynamics behind species interactions, can we hopefully be better prepared to anticipate, prevent and reverse undesired ecosystem collapses.

# 1 Invasions & mutualism

The invasion of naturalized species is changing ecosystems worldwide. Fortunately, during the last four decades, there has been steady progress in understanding the causes and outcomes of biotic invasions. For instance, we now understand that the success of an ecological invasion depends on geographic, bio-climatic, and taxonomic factors, as well as aspects of reproductive biology and general ecology of the introduced organisms. However, we are still unable to successfully predict the outcome of species introductions in a large proportion of cases.

One of the potential explanations for this limited predictive success is that research on both the causes and consequences of invasive species, have focused on the negative interactions between species in the community (competition, and predator-prey relationships). However the establishment of an introduced species depends on, or at least is greatly enhanced, by the establishment of mutualistic relationships<sup>23</sup>—in which both interacting species have a positive outcome. In fact, there are multiple empirical examples of plants that only become successful invaders, when mutualistic partners that pollinate their flowers, or disperse their seeds, are available<sup>24–26</sup>. Understanding the reciprocal relationship between biotic invasions and mutualism is a key step necessary to both improving current predictions of invasion outcomes, as well as evaluating how invasive species are modifying mutualistic systems<sup>23</sup>.

## The invasibility of mutualistic systems

It has been recently shown that there exists a direct link between the structure of ecological networks and ecosystem stability<sup>6,18,27–32</sup>. In particular stable species coexistence in mutualistic networks seems to be favoured by highly diverse, connected, and nested structures<sup>18,28,30,32</sup>. For instance, the nested structure observed in many mutualistic networks—in which specialist species tend to interact with a subset of the species with which a generalist interact—support higher amounts of biodiversity, minimise competition among species in the community, and maximise the range of conditions necessary to have a stable community<sup>18,31</sup>. The unambiguous link between ecosystem stability and the ecosystem response to disturbances is an important argument for investigating the implications of network structure and its vulnerability to drivers of global change.

Perturbations caused by global change are severely modifying the structure of mutualistic networks<sup>33</sup>. Indeed, current evidence suggests that most of its effects are negative<sup>3,29</sup>. Although our understanding of the effects of those perturbations is limited, we know that, for instance, climate change, and habitat modification can lead to shifts on species abundances, and mismatches of phenology, behaviour, or geographic ranges of the interacting species<sup>3,33–35</sup>. Those changes can in turn disrupt the patterns of interactions that determine the structure of mutualistic networks. Some evidence suggest other factors of global change like biotic invasions can also modify the structure of mutualistic networks, for example through changes in the strength of species interactions, and the degree of network nestedness and connectivity<sup>22,36–39</sup>.

I hypothesise that network structure is not only modified by the successive invasion of alien species, but it also plays an important role in determining the invasibility of the community. Two facts provide support to this hypothesis. First, mutualistic interactions are key for the invasion




Figure 1: Each panel represents a plant-animal network with different network structures. Plants compete for resources such as nutrients, but also have indirect interactions mediated by their common pollinators. As the number of shared pollinators is higher, positive effects outweigh negative one. Current theory predicts a higher number of coexisting species as indicated by the size of the matrices.

success of introduced species because introduced species need to have suitable pollinators in the new habitat in order to establish themselves in the community<sup>23,40</sup>. Second, the structure of mutualistic interactions is known to have direct implications for the number of coexisting species in the community<sup>6,9,18,41</sup>. Specifically, structures that minimise plant competition seem to facilitate species coexistence and promote biodiversity<sup>18</sup>. Therefore structural attributes—like the degree of specialisation/generalisation, and the contribution to nestedness, of available pollinators—should influence the likelihood an introduced species becomes invasive<sup>42</sup>.

Previous studies have found that simpler networks, with less species and less interactions between species, are easier to invade<sup>43,44</sup>. However studies are strictly limited to trophic networks in which interactions are chiefly antagonistic (prey-predator)<sup>43–45</sup>. Lamentably, these results cannot be easily extrapolated because mutualistic interactions introduce facilitation and competition feedbacks that are not present in antagonistic interactions. Here I will fill that gap, and study the link between different structural attributes of mutualistic networks—degree of nestedness, contribution to nestedness, and compartmentalisation.

The first step to achieve that is to construct simulated communities of mutualistic interactions. The structure of those communities will be determined by implementing the varying structure of empirically observed networks, as well as by constructing simulated networks with a wide range of structural parameters (Figure 1). This will allow me to evaluate if the network structures that have been found to minimise competition also make the community more vulnerable to invasions.

The second step is to develop models of the population dynamics in mutualistic communities in which the abundances of each species present on the community depends on their growth rate (*births* – *deaths*). These growth rates can in turn be affected by the feedbacks imposed by the interactions with other species. Previous theoretical work has largely assumed that the feedbacks of mutualistic interactions increase the growth rates of the focal species, while antagonistic interactions decrease it, which in turn allows the coexistence of a larger number of species (Figure 1)<sup>18,41</sup>.

This assumption, however, ignores that in mutualistic systems, organisms might also compete




Figure 2: Three possible responses of ecosystems in stable coexistence to the introduction of an alien species (dark blue): a) the alien species did not succeed to coexist, and caused only minor changes in the abundances of the species already present in the community (light blue); b) the alien species persisted with the original species; or c) the alien species became invasive, and five of the original species went extinct.

to optimise the obtained benefits. Indeed evidence suggest that in plant-pollinator systems, the increase of mutualistic interactions caused by invasive species, does not translate in increased facilitation<sup>46</sup>. Because plant reproduction depends strongly in the quality of the mutualistic service, mutualisms can be strongly altered when co-flowering species compete for the service of shared pollinators<sup>40,47</sup>. This happens because the visit of a pollinator does not lead to reproduction if pollen is transferred between different species of plants<sup>48</sup>. Even though there are some mechanisms to mitigate the impacts of competition for mutualism and inter-specific pollen transfer<sup>49,50</sup>, it has been found that invasive species, at high densities, are able to co-opt pollinators from native species<sup>51</sup>, dominate the networks of pollen transport<sup>46,52</sup>, and ultimately decrease the seed output of native species<sup>53</sup>.

The third step is to analyse the ecosystem response to an alien species. Depending on the network structure, and the traits of the native and alien species, the outcome of an species introduction can vary. One possible outcome option is that alien species is not able to invade the community (Figure 2a). Another possibility is that the alien species becomes an invader, which can, in some instances, have catastrophic consequences for other species in the community (Figure 2b, c). respond. Therefore, I will quantify the invasibility of the community as the likelihood that an invader is successful<sup>43,54</sup>.

Because, as mentioned, a nested structure maximises the number of species that can stably coexist<sup>18</sup>, I predict that a similar relationship exists between invasibility and nestedness. However, I also argue that the competition for pollinators confers the community a resistance to invasions that has not been accounted for in previous theoretical or empirical research (Figure 3). These hypotheses link concepts of network structure, apparent competition/facilitation and ecosystem stability, and will allow us to put limited and somewhat disparate empirical observations in context.

Despite the general belief that successful invasive species tend to be super-generalists<sup>23,37,38,55</sup>, a recent meta-analysis has shown that invasive species become very integrated in pollination networks<sup>42</sup>. Specifically, invasive plants seem to interact preferentially with pollinators that are weak contributors to community nestedness, which have been shown to be the less vulnerable to extinction<sup>42,56</sup>. However this empirical observation has not yet been explained mechanistically.




Figure 3: It has been found that a) an increasingly nested structure enhances the number of species that can stable coexist, while b) a compartmentalized structure decreases it. I hypothesise that a similar relationship exist between these attributes of network structure and the invasibility of the community. In addition, I argue that the apparent competition for pollinators increases the resistance of mutualistic communities to invasions by counteracting the positive feedbacks of apparent facilitation.

By testing the hypotheses I propose, not only light will be shed on what makes ecosystem vulnerable to invasions, but also, because they deal with fundamental tenets in theoretical ecology, we will also gain a better understanding of the implications of network structure on the persistence of biodiversity, the link between stability and biodiversity, and the assembly of ecological communities.

## Invasions change the structure of mutualistic networks

After analysing the structures that make an ecological network more or less susceptible to invasions, I then follow by investigating the consequences for the community when the invasion is successful (Figure 4). On one hand, I will explore the feedbacks on the structure of mutualistic network after an successful invasion; on the other hand I will explore when the invasions lead to a large number of secondary extinctions and ultimately ecosystem collapse (Figure 2c).

Successful invasive species have been shown to have pollination effects in the native species trough changes on pollinator populations and behavior<sup>38,57-59</sup>. Although, as mentioned, they can have facilitative effects by increasing the total visitation rates<sup>40,58</sup>, they can also have negative effects by decreasing con-specific or increasing hetero-specific pollen transfer<sup>59</sup>. Therefore, changes on the structure of networks are expected.

Although there are few empirical studies that measure the consequences of ecological invasions in network structure, evidence suggest that invasive species have indeed the ability to modify the structure of mutualistic networks. In particular, invaded communities seem to be more nested<sup>22,42</sup> and less compartmentalised<sup>55</sup> than their un-invaded counterparts, while also having altered visitation patterns<sup>38</sup>. However results are somewhat contradictory<sup>60</sup>, and the limited amount of evidence precludes the generalisation of the patterns observed.

I will use the theoretical results obtained from the population dynamics model to fill the gap on empirical data. Specifically I will quantify the changes on network structure (nestedness,




Figure 4: The network structure determines the invasibility of an ecological community. However, successful invasions also modify the structure itself, whether through a shift in interaction partners or through secondary extinctions in the community.

and compartmentalisation) induced by the invasive species, and the contribution to structure of both the invasive species, and the species it interacts with<sup>42,56</sup>. Therefore, it will be possible to quantify how an initial successful invasion affects the future invasibility of the community. However a successful invasion can, not only affect future invasibility by causing changes on network structure, but also by causing extinction cascades that disrupt mutualisms in the community<sup>57,61</sup>.

It is also possible to quantify the stability of an ecosystem by measuring the number of species extinctions that follow a successful invasion<sup>54,62</sup>. Paradoxically, the same structures that I propose are the most vulnerable to invasions, have also been shown to be the most robust to cascading extinctions in ecological networks<sup>29,55,63</sup>. The research I propose will help disentangle these two seemingly disparate observations.



## 2 Combined effects of global change

Random or top down, down-top, not functional

### 3 Controlling ecological networks

sdfxc

## Research Plan

asd

## References

- [1] M Scheffer, S Carpenter, J Foley, C Folke, and B Walker. Catastrophic shifts in ecosystems. *Nature*, 413(6856):591–596, 2001. ISSN 0028-0836. doi:[10.1038/35098000](https://doi.org/10.1038/35098000).
- [2] Terence P Hughes, David R Bellwood, Carl Folke, Robert S Steneck, and James Wilson. New paradigms for supporting the resilience of marine ecosystems. *Trends in Ecology & Evolution*, 20(7), 2005. doi:[10.1016/j.tree.2005.03.022](https://doi.org/10.1016/j.tree.2005.03.022).
- [3] Jason M. Tylianakis, Raphael K. Didham, Jordi Bascompte, and David a. Wardle. Global change and species interactions in terrestrial ecosystems. *Ecology Letters*, 11(12):1351–1363, 2008. ISSN 1461023X. doi:[10.1111/j.1461-0248.2008.01250.x](https://doi.org/10.1111/j.1461-0248.2008.01250.x).
- [4] C S Holling. Resilience and Stability of Ecological Systems. *Annual Review of Ecology and Systematics*, 4(1):1–23, 1973. ISSN 0066-4162. doi:[10.1146/annurev.es.04.110173.000245](https://doi.org/10.1146/annurev.es.04.110173.000245).
- [5] Lance H Gunderson. Ecological resilience - in theory and application. *Annual Review of Ecology and Systematics*, 31(1):425–439, 2000. ISSN 0066-4162. doi:[10.1146/annurev.ecolsys.31.1.425](https://doi.org/10.1146/annurev.ecolsys.31.1.425).
- [6] Jordi Bascompte, Pedro Jordano, and Jens M Olesen. Asymmetric Coevolutionary Networks Facilitate Biodiversity Maintenance. *Science*, 312(April):431–433, 2006. ISSN 1095-9203. doi:[10.1126/science.1123412](https://doi.org/10.1126/science.1123412).
- [7] Andrew Dobson, David Lodge, Jackie Alder, Graeme S. Cumming, Juan Keymer, Jacquie McGlade, Hal Mooney, James a. Rusak, Osvaldo Sala, Volkmar Wolters, Diana Wall, Rachel Winfree, and Marguerite a. Xenopoulos. Habitat loss, trophic collapse, and the decline of ecosystem services. *Ecology*, 87(8):1915–1924, 2006. ISSN 00129658. doi:[10.1890/0012-9658\(2006\)87\[1915:HLTCAT\]2.0.CO;2](https://doi.org/10.1890/0012-9658(2006)87[1915:HLTCAT]2.0.CO;2).
- [8] Julia Reiss, Jon R. Bridle, José M. Montoya, and Guy Woodward. Emerging horizons in biodiversity and ecosystem functioning research. *Trends in Ecology & Evolution*, 24(9):505–514, 2009. ISSN 01695347. doi:[10.1016/j.tree.2009.03.018](https://doi.org/10.1016/j.tree.2009.03.018).
- [9] Jordi Bascompte and Pedro Jordano. Plant-Animal Mutualistic Networks: The Architecture of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, 38(1):567–593, 2007. ISSN 1543-592X. doi:[10.1146/annurev.ecolsys.38.091206.095818](https://doi.org/10.1146/annurev.ecolsys.38.091206.095818).
- [10] Alexandra Maria Klein, Bernard E Vaissière, James H Cane, Ingolf Steffan-Dewenter, Saul A Cunningham, Claire Kremen, and Teja Tscharntke. Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B: Biological Sciences*, 274(1608):303–313, 2007. ISSN 0962-8452. doi:[10.1098/rspb.2006.3721](https://doi.org/10.1098/rspb.2006.3721).
- [11] Paul Alan Cox and Thomas Elmqvist. Pollinator extinction in the Pacific Islands. *Conservation Biology*, 14(5):1237–1239, 2000. ISSN 08888892. doi:[10.1046/j.1523-1739.2000.00017.x](https://doi.org/10.1046/j.1523-1739.2000.00017.x).
- [12] Peter M. Vitousek, Carla M. D’Antonio, Lloyd L. Loope, Marcel Rejmánek, and Randy Westbrooks. Introduced species: A significant component of human-caused global change. *New Zealand Journal of Ecology*, 21(1):1–16, 1997. ISSN 01106465.
- [13] a. D. Wilton and I. Breitwieser. Composition of the New Zealand seed plant flora. *New Zealand Journal of Botany*, 38(4):537–549, 2000. ISSN 0028-825X. doi:[10.1080/0028825X.2000.9512703](https://doi.org/10.1080/0028825X.2000.9512703).
- [14] David G. Lloyd. Progress in understanding the natural history of New Zealand plants. *New Zealand Journal of Botany*, 23(4):707–722, 1985. ISSN 0028-825X. doi:[10.1080/0028825X.1985.10434239](https://doi.org/10.1080/0028825X.1985.10434239).
- [15] Linda Newstrom and Alastair Robertson. Progress in understanding pollination systems in New Zealand. *New Zealand Journal of Botany*, 43(1):1–59, 2005. ISSN 0028-825X. doi:[10.1080/0028825X.2005.9512943](https://doi.org/10.1080/0028825X.2005.9512943).
- [16] Sandra H. Anderson. The relative importance of birds and insects as pollinators of the New Zealand flora. *New Zealand Journal of Ecology*, 27(2):83–94, 2003. ISSN 01106465.
- [17] Alastair W Robertson, Dave Kelly, Jenny J Ladley, and Ashley D Sparrow. Effects Mistletoes of Pollinator Loss on Endemic New Zealand (Loranthaceae). *Conservation Biology*, 13(3):499–508, 1999.
- [18] Ugo Bastolla, Miguel a Fortuna, Alberto Pascual-García, Antonio Ferrera, Bartolo Luque, and Jordi Bascompte. The architecture of mutualistic networks minimizes competition and increases biodiversity. *Nature*, 458(7241):1018–1020, 2009. ISSN 0028-0836. doi:[10.1038/nature07950](https://doi.org/10.1038/nature07950).
- [19] Javier Garcia-Algarra, Javier Galeano, Juan Manuel Pastor, Jose Maria Iriondo, and Jose J. Ramasco. Rethinking the logistic approach for population dynamics of mutualistic interactions. *Journal of Theoretical Biology*, 363:13, 2013. ISSN 10958541. doi:[10.1016/j.jtbi.2014.08.039](https://doi.org/10.1016/j.jtbi.2014.08.039).

- [20] Scott Bennett, Thomas Wernberg, Euan S. Harvey, Julia Santana-Garcon, and Benjamin J. Saunders. Tropical herbivores provide resilience to a climate-mediated phase shift on temperate reefs. *Ecology Letters*, 18(7):714–723, 2015. ISSN 1461023X. doi:[10.1111/ele.12450](https://doi.org/10.1111/ele.12450).
- [21] Sean P Cornelius, William L Kath, and Adilson E Motter. Realistic control of network dynamics. *Nature Communications*, 4:1942, 2013. ISSN 2041-1723. doi:[10.1038/ncomms2939](https://doi.org/10.1038/ncomms2939).
- [22] Ignasi Bartomeus, Montserrat Vilà, and Luísa Santamaría. Contrasting effects of invasive plants in plant-pollinator networks. *Oecologia*, 155(4):761–770, 2008. ISSN 00298549. doi:[10.1007/s00442-007-0946-1](https://doi.org/10.1007/s00442-007-0946-1).
- [23] D M Richardson, N Allsopp, C M D’Antonio, S J Milton, and M Rejmánek. Plant invasions—the role of mutualisms. *Biological reviews of the Cambridge Philosophical Society*, 75(1):65–93, 2000. ISSN 1464-7931. doi:[10.1111/j.1469-185X.1999.tb00041.x](https://doi.org/10.1111/j.1469-185X.1999.tb00041.x).
- [24] Daniel Simberloff, Daniel Simberloff, Betsy Von Holle, and Betsy Von Holle. Positive interactions of nonindigenous species: invasional meltdown? *Biological Invasions*, pages 21–32, 1999. ISSN 13873547. doi:[10.1023/a:1010086329619](https://doi.org/10.1023/a:1010086329619).
- [25] Daniel Simberloff. Invasional meltdown 6 years later: Important phenomenon, unfortunate metaphor, or both? *Ecology Letters*, 9(8):912–919, 2006. ISSN 1461023X. doi:[10.1111/j.1461-0248.2006.00939.x](https://doi.org/10.1111/j.1461-0248.2006.00939.x).
- [26] Kirsten M Prior, Jennifer M Robinson, S. A. Meadley Dunphy, and Megan E Frederickson. Mutualism between co-introduced species facilitates invasion and alters plant community structure. *Proceedings of the Royal Society B: Biological Sciences*, 282(1800):20142846, 2014. ISSN 0962-8452. doi:[10.1098/rspb.2014.2846](https://doi.org/10.1098/rspb.2014.2846).
- [27] Neil Rooney, Kevin McCann, Gabriel Gellner, and John C Moore. Structural asymmetry and the stability of diverse food webs. *Nature*, 442(7100):265–269, 2006. ISSN 0028-0836. doi:[10.1038/nature04887](https://doi.org/10.1038/nature04887).
- [28] Toshinori Okuyama and J. Nathaniel Holland. Network structural properties mediate the stability of mutualistic communities. *Ecology Letters*, 11(3):208–216, 2008. ISSN 1461023X. doi:[10.1111/j.1461-0248.2007.01137.x](https://doi.org/10.1111/j.1461-0248.2007.01137.x).
- [29] Jason M. Tylianakis, Etienne Laliberté, Anders Nielsen, and Jordi Bascompte. Conservation of species interaction networks. *Biological Conservation*, 143(10):2270–2279, 2010. ISSN 00063207. doi:[10.1016/j.biocon.2009.12.004](https://doi.org/10.1016/j.biocon.2009.12.004). URL <http://dx.doi.org/10.1016/j.biocon.2009.12.004>.
- [30] Elisa Thébault and Colin Fontaine. Stability of ecological communities and the architecture of mutualistic and trophic networks. *Science*, 329(5993):853–856, 2010. ISSN 0036-8075. doi:[10.1126/science.1188321](https://doi.org/10.1126/science.1188321).
- [31] Rudolf P Rohr, Serguei Saavedra, and Jordi Bascompte. On the structural stability of mutualistic systems. *Science*, 345(6195):1253497, 2014. ISSN 0036-8075. doi:[10.1126/science.1253497](https://doi.org/10.1126/science.1253497).
- [32] Alix M C Sauve, Colin Fontaine, and Elisa Thébault. Structure-stability relationships in networks combining mutualistic and antagonistic interactions. *Oikos*, 123(3):378–384, 2014. ISSN 00301299. doi:[10.1111/j.1600-0706.2013.00743.x](https://doi.org/10.1111/j.1600-0706.2013.00743.x).
- [33] Laura a Burkle, John C Marlin, and Tiffany M Knight. Plant-pollinator interactions over 120 years: loss of species, co-occurrence, and function. *Science (New York, N. Y.)*, 339(6127):1611–5, 2013. ISSN 1095-9203. doi:[10.1126/science.1232728](https://doi.org/10.1126/science.1232728).
- [34] Jane Memmott, Paul G. Craze, Nickolas M. Waser, and Mary V. Price. Global warming and the disruption of plant-pollinator interactions. *Ecology Letters*, 10(8):710–717, 2007. ISSN 1461023X. doi:[10.1111/j.1461-0248.2007.01061.x](https://doi.org/10.1111/j.1461-0248.2007.01061.x).
- [35] Stein Joar Hegland, Anders Nielsen, Amparo Lázaro, Anne Line Bjerknes, and Ørjan Totland. How does climate warming affect plant-pollinator interactions? *Ecology Letters*, 12(2):184–195, 2009. ISSN 1461023X. doi:[10.1111/j.1461-0248.2008.01269.x](https://doi.org/10.1111/j.1461-0248.2008.01269.x).
- [36] Jens M. Olesen, Louise I. Eskildsen, and Shadila Venkatasamy. Invasion of pollination networks on oceanic islands: Importance of invader complexes and endemic super generalists. *Diversity and Distributions*, 8(3):181–192, 2002. ISSN 13669516. doi:[10.1046/j.1472-4642.2002.00148.x](https://doi.org/10.1046/j.1472-4642.2002.00148.x).
- [37] Marcelo A Aizen, Carolina L Morales, and Juan M Morales. Invasive mutualists erode native pollination webs. *PLoS Biology*, 6(2):0396–0403, 2008. ISSN 15449173. doi:[10.1371/journal.pbio.0060031](https://doi.org/10.1371/journal.pbio.0060031).
- [38] Montserrat Vilà, Ignasi Bartomeus, Anke C Dietzsch, Theodora Petanidou, Ingolf Steffan-Dewenter, Jane C Stout, and Thomas Tscheulin. Invasive plant integration into native plant-pollinator networks across Europe. *Proceedings. Biological sciences / The Royal Society*, 276(1674):3887–3893, 2009. ISSN 0962-8452. doi:[10.1098/rspb.2009.1076](https://doi.org/10.1098/rspb.2009.1076).

- [39] Anna Traveset, Ruben Heleno, Susana Chamorro, Pablo Vargas, Conley K McMullen, Manuel Nogales, Henri W Herrera, Jens M Olesen, and K McMullen. Invaders of pollination networks in the Galápagos Islands : emergence of novel communities. *Proceedings of the Royal Society B*, 280(iii):20123040, 2013. ISSN 14712954. doi:[10.1098/rspb.2012.3040](https://doi.org/10.1098/rspb.2012.3040).
- [40] Risa D. Sargent and David D. Ackerly. Plant-pollinator interactions and the assembly of plant communities. *Trends in Ecology and Evolution*, 23(3):123–130, 2008. ISSN 01695347. doi:[10.1016/j.tree.2007.11.003](https://doi.org/10.1016/j.tree.2007.11.003).
- [41] David a. Moeller. Facilitative interactions among plants via shared pollinators. *Ecology*, 85(12):3289–3301, 2004. ISSN 00129658. doi:[10.1890/03-0810](https://doi.org/10.1890/03-0810).
- [42] Daniel B Stouffer, Alyssa R Cirtwill, and Jordi Bascompte. How exotic plants integrate into pollination networks. *Journal of Ecology*, 102:1442–1450, 2014. doi:[10.1111/1365-2745.12310](https://doi.org/10.1111/1365-2745.12310).
- [43] Tamara N Romanuk, Yun Zhou, Ulrich Brose, Eric L Berlow, Richard J Williams, and Neo D Martinez. Predicting invasion success in complex ecological networks. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 364(1524):1743–1754, 2009. ISSN 0962-8436. doi:[10.1098/rstb.2008.0286](https://doi.org/10.1098/rstb.2008.0286).
- [44] Núria Galiana, Miguel Lurgi, José M. Montoya, and Bernat C. López. Invasions cause biodiversity loss and community simplification in vertebrate food webs. *Oikos*, 123(6):721–728, 2014. ISSN 16000706. doi:[10.1111/j.1600-0706.2013.00859.x](https://doi.org/10.1111/j.1600-0706.2013.00859.x).
- [45] Benjamin Baiser, Gareth J. Russell, and Julie L. Lockwood. Connectance determines invasion success via trophic interactions in model food webs. *Oikos*, 119(12):1970–1976, 2010. ISSN 00301299. doi:[10.1111/j.1600-0706.2010.18557.x](https://doi.org/10.1111/j.1600-0706.2010.18557.x).
- [46] Martha E. Lopezaraiza-Mikel, Richard B. Hayes, Martin R. Whalley, and Jane Memmott. The impact of an alien plant on a native plant-pollinator network: An experimental approach. *Ecology Letters*, 10(7):539–550, 2007. ISSN 1461023X. doi:[10.1111/j.1461-0248.2007.01055.x](https://doi.org/10.1111/j.1461-0248.2007.01055.x).
- [47] Randall J. Mitchell, Rebecca J. Flanagan, Beverly J. Brown, Nickolas M. Waser, and Jeffrey D. Karron. New frontiers in competition for pollination. *Annals of botany*, 103(9):1403–1413, 2009. ISSN 10958290. doi:[10.1093/aob/mcp062](https://doi.org/10.1093/aob/mcp062).
- [48] Carolina L. Morales and Anna Traveset. Interspecific Pollen Transfer: Magnitude, Prevalence and Consequences for Plant Fitness. *Critical Reviews in Plant Sciences*, 27(4):221–238, 2008. ISSN 0735-2689. doi:[10.1080/07352680802205631](https://doi.org/10.1080/07352680802205631).
- [49] Jaboury Ghazoul. Floral diversity and the facilitation of pollination. *Journal of Ecology*, 94(2):295–304, 2006. ISSN 00220477. doi:[10.1111/j.1365-2745.2006.01098.x](https://doi.org/10.1111/j.1365-2745.2006.01098.x).
- [50] Ignasi Bartomeus, Jordi Bosch, and Montserrat Vilà. High invasive pollen transfer, yet low deposition on native stigmas in a *Carpobrotus*-invaded community. *Annals of Botany*, 102(3):417–424, 2008. ISSN 03057364. doi:[10.1093/aob/mcn109](https://doi.org/10.1093/aob/mcn109).
- [51] Petr Pyšek, Jarošík Chytrý, Vojtěch Milan, Jiří Danihelka, Ingolf Kühn, Jan Pergl, Lubomír Tichý, Jacobus C. Biesmeijer, Willem N. Ellis, William E. Kunin, and Josef Settele. Successful invaders co-opt pollinators of native flora and accumulate insect pollinators with increasing residence time. *Ecological Monographs*, 81(2):277–293, 2011. ISSN 00129615. doi:[10.1890/10-0630.1](https://doi.org/10.1890/10-0630.1).
- [52] Ruben Alarcón. Congruence between visitation and pollen-transport networks in a California plant-pollinator community. *Oikos*, 119(1):35–44, 2010. ISSN 00301299. doi:[10.1111/j.1600-0706.2009.17694.x](https://doi.org/10.1111/j.1600-0706.2009.17694.x).
- [53] Alejandro a. Muñoz and Lohengrin a. Cavieres. The presence of a showy invasive plant disrupts pollinator service and reproductive output in native alpine species only at high densities. *Journal of Ecology*, 96(3):459–467, 2008. ISSN 00220477. doi:[10.1111/j.1365-2745.2008.01361.x](https://doi.org/10.1111/j.1365-2745.2008.01361.x).
- [54] Anthony R Ives and Stephen R Carpenter. Stability and diversity of ecosystems. *Science (New York, N.Y.)*, 317(5834):58–62, 2007. ISSN 0036-8075. doi:[10.1126/science.1133258](https://doi.org/10.1126/science.1133258).
- [55] Matthias Albrecht, Benigno Padrón, Ignasi Bartomeus, and Anna Traveset. Consequences of plant invasions on compartmentalization and species roles in plantpollinator networks. *Proceedings of the Royal Society - B*, 281:20140773, 2014. ISSN 1471-2954. doi:[10.1098/rspb.2014.0773](https://doi.org/10.1098/rspb.2014.0773).
- [56] Serguei Saavedra, Daniel B. Stouffer, Brian Uzzi, and Jordi Bascompte. Strong contributors to network persistence are the most vulnerable to extinction. *Nature*, 478(7368):233–235, 2011. ISSN 0028-0836. doi:[10.1038/nature10433](https://doi.org/10.1038/nature10433).
- [57] C E Christian. Consequences of a biological invasion reveal the importance of mutualism for plant communities. *Nature*, 413(6856):635–639, 2001. ISSN 00280836. doi:[10.1038/35098093](https://doi.org/10.1038/35098093).

- [58] Anne Line Bjerknes, Ørjan Totland, Stein Joar Hegland, and Anders Nielsen. Do alien plant invasions really affect pollination success in native plant species? *Biological Conservation*, 138(1-2):1–12, 2007. ISSN 00063207. doi:[10.1016/j.biocon.2007.04.015](https://doi.org/10.1016/j.biocon.2007.04.015).
- [59] Carolina Laura Morales and Anna Traveset. A meta-analysis of impacts of alien vs. native plants on pollinator visitation and reproductive success of co-flowering native plants. *Ecology Letters*, 12(7):716–728, 2009. ISSN 1461023X. doi:[10.1111/j.1461-0248.2009.01319.x](https://doi.org/10.1111/j.1461-0248.2009.01319.x).
- [60] Gilberto M. Gilberto, Cândida M L Aguiar, Julieta Genini, Celso F. Martins, Fernando C V Zanella, and Marco a R Mello. Invasive Africanized honeybees change the structure of native pollination networks in Brazil. *Biological Invasions*, 14(11):2369–2378, 2012. ISSN 13873547. doi:[10.1007/s10530-012-0235-8](https://doi.org/10.1007/s10530-012-0235-8).
- [61] Mariano a Rodriguez-Cabal, M Noelia Barrios-Garcia, Guillermo C Amico, Marcelo a Aizen, and Nathan J Sanders. Node-by-node disassembly of a mutualistic interaction web driven by species introductions. *Proceedings of the National Academy of Sciences of the United States of America*, 110(41):16503–7, 2013. ISSN 1091-6490. doi:[10.1073/pnas.1300131110](https://doi.org/10.1073/pnas.1300131110).
- [62] W.M. Post and S.L. Pimm. Community assembly and food web stability. *Mathematical Biosciences*, 64(2):169–192, 1983. ISSN 00255564. doi:[10.1016/0025-5564\(83\)90002-0](https://doi.org/10.1016/0025-5564(83)90002-0).
- [63] Daniel B Stouffer and Jordi Bascompte. Compartmentalization increases food-web persistence. *Proceedings of the National Academy of Sciences of the United States of America*, 108(9):3648–52, March 2011. ISSN 1091-6490. doi:[10.1073/pnas.1014353108](https://doi.org/10.1073/pnas.1014353108).