

The pollination trade-off

Fernando Cagua¹, Hugo Marrero², Jason Tylianakis¹, Daniel Stouffer¹

¹ Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury, Private Bag 4800, Christchurch 8041, New Zealand

² Centro de Recursos Naturales Renovables de las Zonas Semiáridas, CONICET, Camino de la Carrindanga Km. 7, 8000 Bahía Blanca, Argentina

Author for correspondence: E. Fernando Cagua (efc29@uclive.ac.nz) - Centre for Integrative Ecology, School of Biological Sciences, University of Canterbury, Private Bag 4800, Christchurch 8140, New Zealand

Introduction

Animal pollination systems play a disproportionately important role in food production and maintenance of global biodiversity (Bascompte & Jordano 2007; Klein *et al.* 2007; Ollerton *et al.* 2011). Alongside the direct mutualisms between plants and their pollinators, the support to biodiversity is substantially achieved as a result of inter-species facilitation—the positive feedback loops that exist between plants that share pollinators or pollinators that share plants [HelpWithRefPlease]. Facilitation is able to promote species coexistence because it offsets the effects of direct competition for resources (Stachowicz 2001). Indeed, studies that are fundamental to our current knowledge of mutualism predict that an upper limit to biodiversity is achieved when the number of mutualistic partners is maximised in a community (Bastolla *et al.* 2009). And yet, not every species of pollinator in natural communities interacts with every species of plant.

One of many possible explanations for why this is true is because sharing a mutualistic partner also has costs (Waser 1978). As soon as there are costs plant start competing for mutualistic partners, even if this competition is not strong enough to produce a **net** negative effect in fitness. The possible detrimental effects of competition for pollinators have been long recognised (Robertson 1895) and widely documented both experimentally (Lewis 1961) and theoretically (Levin & Anderson 1970). Multiple factors (like the number of shared pollinators, the plants’ relative abundance, visitor fidelity, and traits) have been shown to modulate the strength of competition and in turn the fitness outcome (Bobisud & Neuhaus 1975; Campbell 1985; Feinsinger 1987; Morales & Traveset 2008; Mitchell *et al.* 2009). It is currently clear that the effects of these factors on fitness have the potential to drive the evolution of flower phenologies, morphologies and reproductive strategies, as well as ecological differentiation (Caruso 2000; Mitchell *et al.* 2009).

The main two mechanisms through which these ecological factors can affect plant fertilization are (i) by reducing the number of conspecific pollen grains on stigmas or (ii) by increasing the deposition of heterospecific pollen (Morales & Traveset 2008). For example, plants with more attractive flowers might reduce the number of visits—and hence the amount of conspecific pollen—for those less attractive (Yang *et al.* 2011). We denote this as the *quantity* of the pollination service. More subtly, when plants share pollination partners, even receiving a visit might not translate into fertilization due to interspecific pollen transfer (Campbell & Motten 1985). Interspecific pollen transfer affects the pollination service because a focal plant might receive pollen from a different species, or conversely, pollen from the focal plant might be lost to different species. Generally speaking, the higher the proportion of conspecific relative to heterospecific pollen the higher the *quality* of the pollination service. With few exceptions (Rathcke 1988; Lopezaraiza-Mikel *et al.* 2007; Hegland *et al.* 2009; Aizen & Rovere 2010; Tur *et al.* 2016), most of what we know about competition vs. facilitation in pollination systems and its relationship to the quantity and quality of pollination is based mostly on two-plant-species systems. However, competitive interactions between species pairs do not occur in isolation.

Instead, plants often occur in communities in which multiple intertwined factors operate simultaneously and lead to emergent phenomena not observed at smaller scales. For instance, recent empirical evidence suggests that species with original traits generally have fewer interaction partners (Coux *et al.* 2016). This evidence is aligned with the notion that a species that interacts with few species benefits strongly from each of them; in contrast, a species that interacts with a large number of species does so comparatively weakly. If evolutionary specialisation occurs by changing traits to focus on fewer but better partners (Caruso 2000), we should expect a reduction of competition for pollination in species with original traits and an increase in species with a large number of interaction partners (Gibson *et al.* 2012). Alternatively, it might also be the case that flower/pollen density is the dominant force driving pollen transfer (Seifan *et al.* 2014). Abundant plant species might experience a dilution of the available pollinators (Feinsinger 1987) but might also receive more effective visits by capitalising on a larger share of both the pollen being transported by pollinators and animal visits compared to other species. In this case, a potential reduction in the quantity of pollination (the amount of conspecific pollen received) could be compensated by an increase in the quality of pollination (the amount of conspecific relative to heterospecific pollen received). To complicate things more, autonomous self-pollination can also influence a plant sensitivity to competition for pollination because it can ensure reproduction even in the absence of outcross pollen (Kalisz & Vogler 2003). As multiple, potentially correlated, factors may act simultaneously, understanding their relative importance and how plant species harness them to minimise competition for pollination within their communities requires extensive data describing both the structure of the interactions and the pollen flows that occur within it.

Here, we explore competition for pollination at the community scale using comprehensive empirical data from eleven plant-pollinator communities in the Argentinian Pampas. First, we ask the extent to which the outcome of animal-mediated pollination is facilitation or competition. If facilitation is the primary outcome we would expect (i) a larger amount of conspecific pollen deposited in stigmas when flowers are open to animal pollinators than that due to self-pollination and (ii) an overall positive relationship between the amount of conspecific pollen deposited in stigmas and that that would be achieved (Tur *et al.* 2016). Second, we investigate the relative contribution that four ecological factors, that together, describe the role of the plant species and the ecological context, have on competition for pollination. We hypothesize that competition for pollination at the community level should increase as the number of pollination partners a plant has increases (species degree). However, other factors like the plant’s functional originality, its relative abundance, and the visit effectiveness, should have the potential to compensate this increase in competition by improving either the quality or the quantity of pollination. Third, and finally, we explore how plants differ in how they use these factors to minimise competition. Although some plant species might be widespread across multiple locations, competition for pollinators occurs at the community level and are hence defined by the ecological context. If plant species adopt comparable roles across different communities we should expect plant strategies to be more similar within species than across species in the study. On the contrary, if species role is flexible, we should expect plants to be able to adopt different strategies in each community such that competition is more optimally minimised depending on the context.

References

- Aizen, M.A. & Rovere, A.E. (2010). Reproductive interactions mediated by flowering overlap in a temperate hummingbird-plant assemblage. *Oikos*, 119, 696–706.
- Bascompte, J. & Jordano, P. (2007). Plant-Animal Mutualistic Networks: The Architecture of Biodiversity. *Annual Review of Ecology, Evolution, and Systematics*, 38, 567–593.
- Bastolla, U., Fortuna, M.A., Pascual-García, A., Ferrera, A., Luque, B. & Bascompte, J. (2009). The architecture of mutualistic networks minimizes competition and increases biodiversity. *Nature*, 458, 1018–1020.
- Bobisud, L.E. & Neuhaus, R.J. (1975). Pollinator constancy and survival of rare species. *Oecologia*, 21, 263–272.
- Campbell, D.R. (1985). Pollinator Sharing and Seed Set of *Stellaria pubera*: Competition for Pollination.

88 *Ecology*, 66, 544–553.

89 Campbell, D.R. & Motten, A.F. (1985). The Mechanism of Competition for Pollination between Two Forest
90 Herbs. *Ecology*, 66, 554–563.

91 Caruso, C.M. (2000). Competition for Pollination Influences Selection on Floral Traits of *Ipomopsis aggregata*.
92 *Evolution*, 54, 1546–1557.

93 Coux, C., Rader, R., Bartomeus, I. & Tylianakis, J.M. (2016). Linking species functional roles to their
94 network roles. *Ecology Letters*, 19, 762–770.

95 Feinsinger, P. (1987). Effects of plant species on each others pollination: Is community structure influenced?,
96 2, 4.

97 Gibson, M.R., Richardson, D.M. & Pauw, A. (2012). Can floral traits predict an invasive plant’s impact on
98 native plant-pollinator communities? *Journal of Ecology*, 100, 1216–1223.

99 Hegland, S.J., Grytnes, J.-A. & Totland, Ø. (2009). The relative importance of positive and negative
100 interactions for pollinator attraction in a plant community. *Ecological Research*, 24, 929–936.

101 Kalisz, S. & Vogler, D.W. (2003). Benefits of autonomus selfing under unpredictable pollinator environments.
102 *Ecology*, 84, 2928–2942.

103 Klein, A.-M., Vaissiere, B.E., Cane, J.H., Steffan-Dewenter, I., Cunningham, S.A. & Kremen, C. *et al.*
104 (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B:*
105 *Biological Sciences*, 274, 303–313.

106 Levin, D.A. & Anderson, W.W. (1970). Competition for pollinators between simultaneously flowering species.
107 *The American Naturalist*, 104, 455–467.

108 Lewis, H. (1961). Experimental Sympatric Populations of *Clarkia*. *The American Naturalist*, 95, 155–168.

109 Lopezaraiza-Mikel, M.E., Hayes, R.B., Whalley, M.R. & Memmott, J. (2007). The impact of an alien plant
110 on a native plant–Pollinator network: An experimental approach. *Ecology Letters*, 10, 539–550.

111 Mitchell, R.J., Flanagan, R.J., Brown, B.J., Waser, N.M. & Karron, J.D. (2009). New frontiers in competition
112 for pollination. *Annals of Botany*, 103, 1403–1413.

113 Morales, C.L. & Traveset, A. (2008). Interspecific Pollen Transfer: Magnitude, Prevalence and Consequences
114 for Plant Fitness. *Critical Reviews in Plant Sciences*, 27, 221–238.

115 Ollerton, J., Winfree, R. & Tarrant, S. (2011). How many flowering plants are pollinated by animals? *Oikos*,

116 120, 321–326.

117 Rathcke, B. (1988). Interactions for Pollination among Coflowering Shrubs. *Ecology*, 69, 446–457.

118 Robertson, C. (1895). The Philosophy of Flower Seasons, and the Phaenological Relations of the Ento-
 119 mophilous Flora and the Anthophilous Insect Fauna. *The American Naturalist*, 29, 97–117.

120 Seifan, M., Hoch, E.-M., Hanoteaux, S. & Tielbörger, K. (2014). The outcome of shared pollination services
 121 is affected by the density and spatial pattern of an attractive neighbour. *Journal of Ecology*, 102, 953–962.

122 Stachowicz, J.J. (2001). Mutualism, Facilitation, and the Structure of Ecological Communities. *BioScience*,
 123 51, 235.

124 Tur, C., Sáez, A., Traveset, A. & Aizen, M.A. (2016). Evaluating the effects of pollinator-mediated interactions
 125 using pollen transfer networks: Evidence of widespread facilitation in south Andean plant communities.
 126 *Ecology Letters*, 19, 576–586.

127 Waser, N.M. (1978). Interspecific pollen transfer and competition between co-occurring plant species.
 128 *Oecologia*, 36, 223–236.

129 Yang, S., Ferrari, M.J. & Shea, K. (2011). Pollinator Behavior Mediates Negative Interactions between Two
 130 Congeneric Invasive Plant Species. *The American Naturalist*, 177, 110–118.