

Survival Analysis in R

The survival analysis or time-to-event is a clinical course duration variable for each subject having a beginning and an end anywhere along the time line of the complete study. In this chronic kidney disease (CKD) data, it begins when the patient is enrolled into the study or when the treatment begins, and ends when the event of interest, the progression of CKD, is reached or the patient is censored from the study. This report focuses on the survival analysis on the patients' progression of chronic kidney disease.

Load CKD data

First of all, I use `read.csv()` function to load all csv files.

```
creatinine <- read.csv("/Users/yunzhao/Desktop/CKDdata/T_creatinine.csv")
DBP <- read.csv("/Users/yunzhao/Desktop/CKDdata/T_DBP.csv")
demo <- read.csv("/Users/yunzhao/Desktop/CKDdata/T_demo.csv")
glucose <- read.csv("/Users/yunzhao/Desktop/CKDdata/T_glucose.csv")
HGB <- read.csv("/Users/yunzhao/Desktop/CKDdata/T_HGB.csv")
ldl <- read.csv("/Users/yunzhao/Desktop/CKDdata/T_ldl.csv")
meds <- read.csv("/Users/yunzhao/Desktop/CKDdata/T_meds.csv")
SBP <- read.csv("/Users/yunzhao/Desktop/CKDdata/T_SBP.csv")
stage <- read.csv("/Users/yunzhao/Desktop/CKDdata/T_stage.csv")
```

Import libraries

This block of code loads the required packages for survival analysis.

```
library(tidyverse)
```

```
## — Attaching packages ————— tidyverse 1.3.0 —
```

```
## ✓ ggplot2 3.3.3      ✓ purrr 0.3.4
## ✓ tibble 3.1.0       ✓ dplyr 1.0.5
## ✓ tidyr 1.1.3        ✓ stringr 1.4.0
## ✓ readr 1.3.1        ✓ forcats 0.5.0
```

```
## — Conflicts ————— tidyverse_conflicts() —
## x dplyr::filter() masks stats::filter()
## x dplyr::lag()     masks stats::lag()
```

```
library(survival)
library(survminer)
```

```
## Loading required package: ggpubr
```

```
library(ggfortify)
library(ranger)
library(caTools)
library(kableExtra)
```

```
##
## Attaching package: 'kableExtra'
```

```
## The following object is masked from 'package:dplyr':
##
##      group_rows
```

Data pre-processing

Before the analysis, I join all csv files with `left_join` function to create the full CKD data set.

```
# Join csv data
CKDdata <- left_join(creatinine,DBP,by=c("id","time")) %>%
  left_join(glucose, by=c("id","time")) %>%
  left_join(HGB, by=c("id","time")) %>%
  left_join(ldl, by=c("id","time")) %>%
  left_join(SBP, by=c("id","time")) %>%
  left_join(demo,by=c("id")) %>%
  left_join(stage,by=c("id")) %>%
  left_join(meds,by=c("id"))

# Rename columns
names(CKDdata) <- c("id","creatinine","time","DBP","glucose",
  "HGB","ldl","SBP","race","gender","age",
  "stage","drug","dosage","start_day","end_day")

# stage-True = 1: the patient would progress to CKD
# stage-False = 0: the patient would not progress to CKD
CKDdata$stage <- ifelse(CKDdata$stage=="True",1,0)

# Exclude the "id" column
CKDdata$id <- NULL

head(CKDdata)
```

##	creatinine	time	DBP	glucose	HGB	ldl	SBP	race	gender	age	stage
## 1	1.29	0	95.32	6.24	13.51	161.49	134.11	Unknown	Male	70	1
## 2	1.29	0	95.32	6.24	13.51	161.49	134.11	Unknown	Male	70	1
## 3	1.29	0	95.32	6.24	13.51	161.49	134.11	Unknown	Male	70	1
## 4	1.29	0	95.32	6.24	13.51	161.49	134.11	Unknown	Male	70	1
## 5	1.29	0	95.32	6.24	13.51	161.49	134.11	Unknown	Male	70	1
## 6	1.29	0	95.32	6.24	13.51	161.49	134.11	Unknown	Male	70	1
##	drug	dosage	start_day	end_day							
## 1	atorvastatin	10	19	109							
## 2	atorvastatin	10	117	207							
## 3	losartan	100	19	289							
## 4	losartan	100	403	493							
## 5	losartan	100	587	677							
## 6	metformin	1000	19	109							

Kaplan-Meier Model

1. Kaplan-Meier Model

The Kaplan-Meier method is the most common statistical method to estimate survival times and probabilities. In preparing Kaplan-Meier survival analysis, each subject is characterized by two variables: (1) serial time; (2) status at the end of the serial time, which means the event of interest (CKD progression) occurrence or censored.

In the first step, I use Surv() to build the standard survival object: (1) The variable ‘time’ records all survival time; (2) ‘stage’ indicates whether a patient’s progression of CKD was observed (stage = 1) or that survival time was censored (stage = 0); (3) a “+” after the time in the print out of ‘kmf’ indicates censoring.

```
kmf <- with(CKDdata, Surv(time, stage))
head(kmf, 100)
```

##	[1]	0	0	0	0	0	0	0	0	0	0	0	107	107	107	107
##	[16]	107	107	107	107	107	107	107	286	286	286	286	286	286	286	286
##	[31]	286	286	286	382	382	382	382	382	382	382	382	382	382	382	580
##	[46]	580	580	580	580	580	580	580	580	580	580	688	688	688	688	688
##	[61]	688	688	688	688	688	688	0+	0+	0+	0+	68+	68+	68+	68+	289
+																
##	[76]	289+	289+	289+	387+	387+	387+	387+	470+	470+	470+	470+	0	0	184	184
##	[91]	430	430	502	502	621	621	0+	0+	0+	0+					

Then I use Surv(time,stage) ~ 1 and the survfit() function to produce the Kaplan-Meier estimates of the probability of survival over time.

```

kmf_fit <- survfit(Surv(time, stage) ~ 1, data = CKDdata)

# Plot kmf_fit
ggsurvplot(kmf_fit,
            surv.median.line = "hv", # add medians survival
            pval = TRUE, # add p-value and tervals
            conf.int = TRUE,
            risk.table = TRUE, # add risk table
            tables.height = 0.2,
            tables.theme = theme_cleantable(),
            ggtheme = theme_bw(),
            xlab = "Days",
            ylab = "Overall survival probability",
            title="Kaplan-Meier Plot")


```

```

## Warning in .pvalue(fit, data = data, method = method, pval = pval, pval.coord =
pval.coord, : There are no survival curves to be compared.
## This is a null model.

```

Kaplan-Meier Plot




The plot produced by ggsurvplot shows the step function (solid line) with associated confidence bands (shaded area). The horizontal x-axis represents the survival duration for the interval. The vertical y-axis is the change in cumulative probability. Censored observations, indicated by tick marks, reduce the cumulative survival between intervals. The plotting result suggests that the overall patients' survival probability decreases with the increase in time.

Next, I look at survival curves by treatment (drug and dosage), gender and race, respectively. For age, I do a little data munging to look at survival by age (the age is lower than 60 years old and over 60 years old).



```
# drug
kmf_drug_fit <- survfit(Surv(time, stage) ~ drug, data = CKDdata)
ggsurvplot(kmf_drug_fit,
  surv.median.line = "hv", # add medians survival
  pval = TRUE, # add p-value and tervals
  conf.int = TRUE,
  tables.height = 0.2, tables.theme = theme_cleantable(),
  ggtheme = theme_bw(),
  xlab = "Days",
  title="Kaplan-Meier Plot (Drug)")
```

Kaplan-Meier Plot (Drug)



```
# dosage
kmf_dosage_fit <- survfit(Surv(time, stage) ~ dosage, data = CKDdata)
ggsurvplot(kmf_dosage_fit,
  surv.median.line = "hv", # add medians survival
  pval = TRUE, # add p-value and tervals
  conf.int = TRUE,
  tables.height = 0.2, tables.theme = theme_cleantable(),
  ggtheme = theme_bw(),
  xlab = "Days",
  title="Kaplan-Meier Plot (Dosage)")
```


Kaplan-Meier Plot (Dosage)



gender

```
kmf_gender_fit <- survfit(Surv(time, stage) ~ gender, data = CKDdata)
ggsurvplot(kmf_gender_fit,
  surv.median.line = "hv", # add medians survival
  pval = TRUE, # add p-value and tervals
  conf.int = TRUE,
  risk.table = TRUE, # add risk table
  tables.height = 0.2,
  tables.theme = theme_cleantable(),
  ggtheme = theme_bw(),
  xlab = "Days",
  title="Kaplan-Meier Plot (Gender)")
```

Kaplan-Meier Plot (Gender)



Number at risk

gender=Female

5428	3313	1869	562	0
------	------	------	-----	---

gender=Male


5222	3023	1788	501	0
------	------	------	-----	---

race

```
kmf_race_fit <- survfit(Surv(time, stage) ~ race, data = CKDdata)
ggsurvplot(kmf_race_fit,
  surv.median.line = "hv", # add medians survival
  pval = TRUE, # add p-value and tervals
  conf.int = TRUE,
  risk.table = TRUE, # add risk table
  tables.height = 0.2,
  tables.theme = theme_cleantable(),
  ggtheme = theme_bw(),
  xlab = "Days",
  title="Kaplan-Meier Plot (Race)")
```

Kaplan-Meier Plot (Race)

Strata + race=Asian + race=Black + race=Hispanic + race=Unknown + race=White




Number at risk

race=Asian	617	351	208	58	0
race=Black	882	527	285	84	0
race=Hispanic	193	109	48	6	0
race=Unknown	1000	604	318	103	0
race=White	7958	4745	2798	812	0

age

```
dat_age <- mutate(CKDdata, ag=ifelse((age < 60), "lower60", "over60"),
  ag=factor(ag))
kmf_age_fit <- survfit(Surv(time, stage) ~ ag, data = dat_age)
ggsurvplot(kmf_age_fit,
  surv.median.line = "hv", # add medians survival
  pval = TRUE, # add p-value and tervals
  conf.int = TRUE,
  risk.table = TRUE, # add risk table
  tables.height = 0.2,
  tables.theme = theme_cleantable(),
  ggtheme = theme_bw(),
  xlab = "Days",
  title="Kaplan-Meier Plot (Age)")
```


Kaplan-Meier Plot (Age)



Through all plotting results, it only can be concluded that female patients clearly have a better chance of surviving than male patients. Curves among other groups appear to overlap.

2. Estimating x-year survival

One quantity of interest in a survival analysis is the probability of surviving beyond a certain number (x) of years. In this CKD data, the aim is to estimate the probability of patients' surviving to one year before they progress to CKD. I use `summary()` function with "time" argument (The "time" variables are in days, so I use `times=365.25`).

```
summary(survfit(Surv(time, stage) ~ 1, data=CKDdata), times = 365.25)
```

```
## Call: survfit(formula = Surv(time, stage) ~ 1, data = CKDdata)
##
##   time n.risk n.event survival std.err lower 95% CI upper 95% CI
##   365   4465   2371    0.72  0.00504    0.71      0.73
```

The summary result suggests that the one-year probability of survival before CKD progression in this dataset is 72%.

3. Estimating median survival time

The second quantity of interest in a survival analysis is the average survival time, which I quantify using the median. Survival times are not expected to be normally distributed, so the mean is not an appropriate summary. I obtain the median survival time of 558 days from `survfit` object. The median survival time is the time corresponding to a survival probability of 0.5.

```
survfit(Surv(time, stage) ~ 1, data = CKDdata)
```

```
## Call: survfit(formula = Surv(time, stage) ~ 1, data = CKDdata)
##
##           n  events  median 0.95LCL 0.95UCL
##    10650    4153    558    553    563
```

4. Comparing survival times between groups

I conduct the between-group significance test using a log-rank test. The log-rank test equally weights observations over the entire follow-up time and is the most common way to compare survival times between groups. I get the log-rank p-value using the `survdif` function according to gender, race, age, and treatment (drug and dosage) in the CKD data set.

```
# Survival times between groups
survdif(Surv(time, stage) ~ gender, data=CKDdata)
```

```
## Call:
## survdif(formula = Surv(time, stage) ~ gender, data = CKDdata)
##
##           N Observed Expected (O-E)^2/E (O-E)^2/V
## gender=Female 5428    1457    2157    227    486
## gender=Male  5222    2696    1996    245    486
##
##  Chisq= 486  on 1 degrees of freedom, p= <2e-16
```

```
survdif(Surv(time, stage) ~ race, data=CKDdata)
```

```
## Call:
## survdif(formula = Surv(time, stage) ~ race, data = CKDdata)
##
##           N Observed Expected (O-E)^2/E (O-E)^2/V
## race=Asian    617    169    225.0    13.9219    15.0787
## race=Black    882    287    335.4     6.9898     7.8249
## race=Hispanic  193     44     58.5     3.6106     3.7579
## race=Unknown 1000    515    389.5    40.4631    45.9193
## race=White   7958   3138   3144.6     0.0139     0.0589
##
##  Chisq= 66.8  on 4 degrees of freedom, p= 1e-13
```

```
survdif(Surv(time, stage) ~ ag, data = dat_age)
```

```
## Call:
## survdiff(formula = Surv(time, stage) ~ ag, data = dat_age)
##
##           N Observed Expected (O-E)^2/E (O-E)^2/V
## ag=lower60 1510      655      652  0.01400  0.0171
## ag=over60  9140     3498     3501  0.00261  0.0171
##
##  Chisq= 0   on 1 degrees of freedom, p= 0.9
```

```
survdiff(Surv(time, stage) ~ drug, data=CKDdata)
```

```
## Call:
## survdiff(formula = Surv(time, stage) ~ drug, data = CKDdata)
##
## n=10516, 134 observations deleted due to missingness.
##
##           N Observed Expected (O-E)^2/E (O-E)^2/V
## drug=atenolol      524      263   203.02  1.77e+01  1.91e+01
## drug=atorvastatin 1649      453   645.12  5.72e+01  6.97e+01
## drug=bisoprolol     66       30    29.98  1.04e-05  1.07e-05
## drug=canagliflozin  41        3    16.60  1.11e+01  1.14e+01
## drug=carvedilol    284      132   102.79  8.30e+00  8.72e+00
## drug=dapagliflozin  16        0     6.22  6.22e+00  6.36e+00
## drug=irbesartan     68       60    30.71  2.79e+01  2.87e+01
## drug=labetalol     15        0     3.88  3.88e+00  3.99e+00
## drug=losartan     1182      393   443.53  5.76e+00  6.62e+00
## drug=lovastatin     213       83    84.32  2.07e-02  2.18e-02
## drug=metformin     2357      965   943.18  5.05e-01  6.74e-01
## drug=metoprolol    1031      579   375.67  1.10e+02  1.24e+02
## drug=nebivolol      35       35    17.20  1.84e+01  1.89e+01
## drug=olmesartan    137       45    38.26  1.19e+00  1.24e+00
## drug=pitavastatin   21        6     8.57  7.73e-01  7.91e-01
## drug=pravastatin   466      240   198.75  8.56e+00  9.23e+00
## drug=propranolol    59        0    25.28  2.53e+01  2.71e+01
## drug=rosuvastatin   500      117   182.36  2.34e+01  2.51e+01
## drug=simvastatin   1305      445   520.04  1.08e+01  1.27e+01
## drug=telmisartan    34       29    17.15  8.18e+00  8.62e+00
## drug=valsartan     513      233   218.37  9.81e-01  1.07e+00
##
##  Chisq= 356   on 20 degrees of freedom, p= <2e-16
```

```
survdiff(Surv(time, stage) ~ dosage, data=CKDdata)
```

```
## Call:
## survdiff(formula = Surv(time, stage) ~ dosage, data = CKDdata)
##
## n=10516, 134 observations deleted due to missingness.
##
##           N Observed Expected (O-E)^2/E (O-E)^2/V
## dosage=2      21         6      8.57  7.73e-01  7.91e-01
## dosage=5     137        71     59.11  2.39e+00  2.48e+00
## dosage=6.25    17         5      5.09  1.63e-03  1.67e-03
## dosage=10     827       244    297.34  9.57e+00  1.06e+01
## dosage=12.5   121        41     48.68  1.21e+00  1.26e+00
## dosage=20    1492       493    591.29  1.63e+01  1.97e+01
## dosage=25     414       193    145.28  1.57e+01  1.66e+01
## dosage=40    1714       529    689.08  3.72e+01  4.59e+01
## dosage=50    1158       573    402.72  7.20e+01  8.21e+01
## dosage=80     314       144    129.01  1.74e+00  1.84e+00
## dosage=100   1240       510    492.04  6.56e-01  7.64e-01
## dosage=150     16         8      5.76  8.67e-01  8.88e-01
## dosage=160    160        37     70.73  1.61e+01  1.69e+01
## dosage=200     87        45     38.80  9.89e-01  1.03e+00
## dosage=300     77        55     33.93  1.31e+01  1.35e+01
## dosage=320    237       107     98.98  6.49e-01  6.85e-01
## dosage=500    844       234    293.19  1.19e+01  1.32e+01
## dosage=600     15         0      3.88  3.88e+00  3.99e+00
## dosage=640    112        85     47.52  2.96e+01  3.07e+01
## dosage=750     51        16     31.09  7.32e+00  7.76e+00
## dosage=850     61        36     25.57  4.26e+00  4.38e+00
## dosage=1000   757       304    304.32  3.28e-04  3.66e-04
## dosage=1500   128       128     58.87  8.12e+01  8.41e+01
## dosage=1700    96        54     44.67  1.95e+00  2.08e+00
## dosage=2000   396       193    173.16  2.27e+00  2.43e+00
## dosage=2550    24         0     12.33  1.23e+01  1.26e+01
##
## Chisq= 353 on 25 degrees of freedom, p= <2e-16
```

Cox Regression

Cox proportional hazards model

The Cox proportional hazards model is a semi-parametric model that can be used to fit univariable and multivariable regression models that have survival outcomes. In this part: (1) I fit a Cox proportional hazards model that makes use of all the covariates in the data. I split the CKD data set into training and test subsets to make prediction and visualize the results. I also compute Confusion Matrix and ROC curves to assess the model performance. (2) Hazard ratios The quantity of interest from a Cox regression is a hazard ratio (HR). The HR represents the ratio of hazards between two groups at any particular point in time. The regression parameter (coef) from column estimate in the coxph, then $HR = \exp(\text{coef})$.

```
table(CKDdata$stage)
```

```
##
##      0      1
## 6497 4153
```

```
# Split CKD data set
split_data <- sample(1:nrow(CKDdata), 0.8 * nrow(CKDdata), FALSE)
training_set <- CKDdata[split_data,]
test_set <- CKDdata[-split_data,]

# check 1 by 1 for infinite coefficients
training_set$id <- NULL
names(training_set)
```

```
## [1] "creatinine" "time"      "DBP"      "glucose"   "HGB"
## [6] "ldl"        "SBP"      "race"     "gender"   "age"
## [11] "stage"      "drug"     "dosage"
```

```
coxph(Surv(time,stage) ~ DBP, data=training_set)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ DBP, data = training_set)
##
##              coef exp(coef) se(coef)      z      p
## DBP 0.001478    1.001479 0.002937 0.503 0.615
##
## Likelihood ratio test=0.25  on 1 df, p=0.6151
## n= 2233, number of events= 882
## (6287 observations deleted due to missingness)
```

```
coxph(Surv(time,stage) ~ glucose, data=training_set)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ glucose, data = training_set)
##
##               coef exp(coef) se(coef)      z      p
## glucose 0.14782    1.15930  0.01215 12.17 <2e-16
##
## Likelihood ratio test=137.1  on 1 df, p=< 2.2e-16
## n= 5817, number of events= 2119
## (2703 observations deleted due to missingness)
```

```
coxph(Surv(time,stage) ~ HGB, data=training_set)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ HGB, data = training_set)
##
##               coef exp(coef) se(coef)      z      p
## HGB -0.11703    0.88956  0.01381 -8.474 <2e-16
##
## Likelihood ratio test=70.88  on 1 df, p=< 2.2e-16
## n= 4651, number of events= 1812
## (3869 observations deleted due to missingness)
```

```
coxph(Surv(time,stage) ~ ldl, data=training_set)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ ldl, data = training_set)
##
##               coef exp(coef) se(coef)      z      p
## ldl 0.0030588 1.0030634 0.0008004 3.821 0.000133
##
## Likelihood ratio test=14.2  on 1 df, p=0.0001644
## n= 5984, number of events= 2151
## (2536 observations deleted due to missingness)
```

```
coxph(Surv(time,stage) ~ SBP, data=training_set)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ SBP, data = training_set)
##
##               coef exp(coef) se(coef)      z      p
## SBP -0.001438  0.998563  0.002271 -0.633 0.527
##
## Likelihood ratio test=0.4  on 1 df, p=0.5261
## n= 2221, number of events= 872
## (6299 observations deleted due to missingness)
```

```
coxph(Surv(time,stage) ~ race, data=training_set)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ race, data = training_set)
##
##               coef exp(coef) se(coef)      z      p
## raceBlack      0.06584    1.06806  0.10974  0.600  0.54851
## raceHispanic  -0.09574    0.90870  0.18999 -0.504  0.61431
## raceUnknown    0.57662    1.78001  0.09955  5.792 6.94e-09
## raceWhite      0.28187    1.32560  0.08870  3.178  0.00148
##
## Likelihood ratio test=60.76  on 4 df, p=2.011e-12
## n= 8520, number of events= 3344
```

```
coxph(Surv(time,stage) ~ gender, data=training_set)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ gender, data = training_set)
##
##               coef exp(coef) se(coef)      z      p
## genderMale 0.68438    1.98254  0.03626 18.88 <2e-16
##
## Likelihood ratio test=374.4  on 1 df, p=< 2.2e-16
## n= 8520, number of events= 3344
```

```
coxph(Surv(time,stage) ~ age, data=training_set)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ age, data = training_set)
##
##               coef exp(coef) se(coef)      z      p
## age 0.000606    1.000606  0.001795  0.338  0.736
##
## Likelihood ratio test=0.11  on 1 df, p=0.7356
## n= 8520, number of events= 3344
```

```
coxph(Surv(time,stage) ~ drug, data=training_set)
```

```
## Warning in fitter(X, Y, istrat, offset, init, control, weights = weights, :
## Loglik converged before variable 5,7,16 ; coefficient may be infinite.
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ drug, data = training_set)
##
##               coef exp(coef) se(coef)      z      p
## drugatorvastatin -6.211e-01 5.374e-01 8.688e-02 -7.149 8.75e-13
## drugbisoprolol   -1.310e-01 8.772e-01 2.115e-01 -0.619 0.53562
## drugcanagliflozin -1.773e+00 1.698e-01 5.814e-01 -3.050 0.00229
## drugcarvedilol   -5.624e-03 9.944e-01 1.186e-01 -0.047 0.96216
## drugdapagliflozin -1.528e+01 2.319e-07 7.483e+02 -0.020 0.98371
## drugirbesartan    4.988e-01 1.647e+00 1.599e-01  3.120 0.00181
## druglabetalol    -1.528e+01 2.321e-07 9.551e+02 -0.016 0.98724
## druglosartan     -3.704e-01 6.905e-01 8.898e-02 -4.162 3.15e-05
## druglovastatin   -2.701e-01 7.633e-01 1.418e-01 -1.905 0.05676
## drugmetformin    -2.027e-01 8.165e-01 7.735e-02 -2.620 0.00878
## drugmetoprolol    2.541e-01 1.289e+00 8.254e-02  3.079 0.00208
## drugnebivolol     4.228e-01 1.526e+00 2.118e-01  1.997 0.04587
## drugolmesartan    1.205e-02 1.012e+00 1.806e-01  0.067 0.94678
## drugpitavastatin -4.971e-01 6.083e-01 4.525e-01 -1.099 0.27194
## drugpravastatin  -2.930e-02 9.711e-01 1.003e-01 -0.292 0.77016
## drugpropranolol  -1.530e+01 2.272e-07 4.041e+02 -0.038 0.96981
## drugrosuvastatin -6.554e-01 5.192e-01 1.213e-01 -5.403 6.55e-08
## drugsimvastatin  -3.737e-01 6.882e-01 8.652e-02 -4.319 1.57e-05
## drugtelmisartan   5.406e-01 1.717e+00 2.288e-01  2.363 0.01812
## drugvalsartan    -5.472e-02 9.467e-01 9.875e-02 -0.554 0.57947
##
## Likelihood ratio test=336.7 on 20 df, p=< 2.2e-16
## n= 8406, number of events= 3308
## (114 observations deleted due to missingness)
```

```
coxph(Surv(time,stage) ~ dosage, data=training_set)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ dosage, data = training_set)
##
##               coef exp(coef) se(coef)      z      p
## dosage 0.0001094 1.0001094 0.0000316 3.461 0.000538
##
## Likelihood ratio test=11.51 on 1 df, p=0.000693
## n= 8406, number of events= 3308
## (114 observations deleted due to missingness)
```

```
# Cox regression model_1st round
cox1 <- coxph(Surv(time, stage) ~ ., data=training_set)
```

```
## Warning in fitter(X, Y, istrat, offset, init, control, weights = weights, :
## Loglik converged before variable 17,19,28 ; coefficient may be infinite.
```

```
summary(cox1)
```



```
## Call:
## coxph(formula = Surv(time, stage) ~ ., data = training_set)
##
##      n= 1871, number of events= 705
##      (6649 observations deleted due to missingness)
##
##              coef    exp(coef)    se(coef)      z  Pr(>|z|)
## creatinine      2.064e-02  1.021e+00  1.360e-01  0.152 0.879371
## DBP             5.600e-03  1.006e+00  4.337e-03  1.291 0.196588
## glucose         2.310e-02  1.023e+00  2.725e-02  0.848 0.396662
## HGB            -9.234e-02  9.118e-01  3.042e-02 -3.035 0.002404 **
## ldl            -9.274e-03  9.908e-01  1.588e-03 -5.841 5.20e-09 ***
## SBP            -7.469e-03  9.926e-01  3.388e-03 -2.205 0.027467 *
## raceBlack      -7.558e-01  4.696e-01  2.503e-01 -3.020 0.002531 **
## raceHispanic   -7.042e-02  9.320e-01  3.733e-01 -0.189 0.850363
## raceUnknown     7.746e-01  2.170e+00  2.277e-01  3.401 0.000671 ***
## raceWhite       1.871e-01  1.206e+00  1.969e-01  0.950 0.341961
## genderMale      8.610e-01  2.366e+00  1.046e-01  8.231 < 2e-16 ***
## age            -2.132e-02  9.789e-01  4.330e-03 -4.923 8.53e-07 ***
## drugatorvastatin -4.715e-01  6.241e-01  2.015e-01 -2.340 0.019273 *
## drugbisoprolol   9.660e-02  1.101e+00  4.854e-01  0.199 0.842246
## drugcanagliflozin -1.796e+00  1.659e-01  1.017e+00 -1.766 0.077461 .
## drugcarvedilol  -1.501e-01  8.606e-01  2.906e-01 -0.517 0.605469
## drugdapagliflozin -1.629e+01  8.397e-08  3.234e+03 -0.005 0.995981
## drugirbesartan   8.641e-01  2.373e+00  4.049e-01  2.134 0.032844 *
## druglabetalol   -1.587e+01  1.279e-07  1.446e+03 -0.011 0.991245
## druglosartan    -4.888e-01  6.134e-01  2.162e-01 -2.261 0.023780 *
## druglovastatin  -8.701e-02  9.167e-01  3.395e-01 -0.256 0.797762
## drugmetformin   -5.928e-01  5.528e-01  2.395e-01 -2.475 0.013307 *
## drugmetoprolol   2.942e-01  1.342e+00  2.021e-01  1.456 0.145483
## drugnebivolol   -3.165e-01  7.287e-01  3.585e-01 -0.883 0.377372
## drugolmesartan  -4.731e-01  6.231e-01  3.523e-01 -1.343 0.179295
## drugpitavastatin 9.264e-02  1.097e+00  1.021e+00  0.091 0.927674
## drugpravastatin  6.581e-01  1.931e+00  2.261e-01  2.911 0.003608 **
## drugpropranolol -1.615e+01  9.693e-08  1.076e+03 -0.015 0.988030
## drugrosuvastatin -8.067e-01  4.463e-01  2.821e-01 -2.860 0.004239 **
## drugsimvastatin -4.738e-01  6.226e-01  2.145e-01 -2.209 0.027148 *
## drugtelmisartan  1.073e+00  2.925e+00  4.863e-01  2.207 0.027325 *
## drugvalsartan    4.099e-02  1.042e+00  2.295e-01  0.179 0.858263
## dosage          4.486e-04  1.000e+00  1.372e-04  3.271 0.001072 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
##              exp(coef) exp(-coef) lower .95 upper .95
## creatinine      1.021e+00  9.796e-01  0.78197  1.3327
## DBP             1.006e+00  9.944e-01  0.99710  1.0142
## glucose         1.023e+00  9.772e-01  0.97014  1.0795
## HGB             9.118e-01  1.097e+00  0.85902  0.9678
## ldl             9.908e-01  1.009e+00  0.98769  0.9939
## SBP             9.926e-01  1.007e+00  0.98599  0.9992
## raceBlack       4.696e-01  2.129e+00  0.28753  0.7670
## raceHispanic    9.320e-01  1.073e+00  0.44841  1.9371
## raceUnknown     2.170e+00  4.609e-01  1.38850  3.3902
```

```
## raceWhite      1.206e+00  8.294e-01  0.81974  1.7734
## genderMale     2.366e+00  4.227e-01  1.92702  2.9038
## age            9.789e-01  1.022e+00  0.97064  0.9873
## drugatorvastatin 6.241e-01  1.602e+00  0.42049  0.9263
## drugbisoprolol  1.101e+00  9.079e-01  0.42541  2.8517
## drugcanagliflozin 1.659e-01  6.028e+00  0.02259  1.2186
## drugcarvedilol  8.606e-01  1.162e+00  0.48685  1.5212
## drugdapagliflozin 8.397e-08  1.191e+07  0.00000  Inf
## drugirbesartan  2.373e+00  4.214e-01  1.07301  5.2474
## druglabetalol   1.279e-07  7.820e+06  0.00000  Inf
## druglosartan     6.134e-01  1.630e+00  0.40151  0.9371
## druglovastatin   9.167e-01  1.091e+00  0.47119  1.7833
## drugmetformin    5.528e-01  1.809e+00  0.34569  0.8839
## drugmetoprolol   1.342e+00  7.451e-01  0.90310  1.9943
## drugnebivolol    7.287e-01  1.372e+00  0.36090  1.4714
## drugolmesartan   6.231e-01  1.605e+00  0.31237  1.2428
## drugpitavastatin 1.097e+00  9.115e-01  0.14842  8.1090
## drugpravastatin  1.931e+00  5.178e-01  1.23978  3.0079
## drugpropranolol  9.693e-08  1.032e+07  0.00000  Inf
## drugrosuvastatin 4.463e-01  2.240e+00  0.25679  0.7758
## drugsimvastatin  6.226e-01  1.606e+00  0.40896  0.9479
## drugtelmisartan  2.925e+00  3.419e-01  1.12757  7.5872
## drugvalsartan    1.042e+00  9.598e-01  0.66437  1.6338
## dosage          1.000e+00  9.996e-01  1.00018  1.0007
##
## Concordance= 0.749 (se = 0.012 )
## Likelihood ratio test= 362.9 on 33 df, p=<2e-16
## Wald test            = 297.4 on 33 df, p=<2e-16
## Score (logrank) test = 379.8 on 33 df, p=<2e-16
```

Coefficients of drugs: dapagliflozin, labetalol, and propranolol were infinite, so these drugs are removed in the second round of modeling.

```
# Cox regression model_2nd round
CKDdata_1 <- CKDdata %>% filter(drug != "dapagliflozin") %>%
  filter(drug != "labetalol") %>%
  filter(drug != "propranolol")

CKDdata_1$id <- NULL

split_data_1 <- sample(1:nrow(CKDdata_1), 0.8 * nrow(CKDdata_1), FALSE)
training_set_1 <- CKDdata_1[split_data_1,]
test_set_1 <- CKDdata_1[-split_data_1,]

cox2 <- coxph(Surv(time, stage) ~ ., data=training_set_1)
```

```
## Warning in fitter(X, Y, istrat, offset, init, control, weights = weights, :
## Loglik converged before variable 24 ; coefficient may be infinite.
```

```
summary(cox2)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ ., data = training_set_1)
##
##      n= 1817, number of events= 710
##      (6523 observations deleted due to missingness)
##
##              coef exp(coef) se(coef)      z Pr(>|z|)
## creatinine      -5.166e-03  9.948e-01  1.346e-01 -0.038 0.969379
## DBP              9.152e-03  1.009e+00  4.280e-03  2.138 0.032497 *
## glucose          2.820e-02  1.029e+00  2.701e-02  1.044 0.296497
## HGB             -8.564e-02  9.179e-01  3.067e-02 -2.793 0.005227 **
## ldl             -8.971e-03  9.911e-01  1.575e-03 -5.697 1.22e-08 ***
## SBP            -9.070e-03  9.910e-01  3.309e-03 -2.741 0.006122 **
## raceBlack       -5.051e-01  6.034e-01  2.403e-01 -2.102 0.035569 *
## raceHispanic    -3.934e-01  6.748e-01  4.519e-01 -0.871 0.383993
## raceUnknown      8.375e-01  2.311e+00  2.231e-01  3.755 0.000174 ***
## raceWhite        2.680e-01  1.307e+00  1.903e-01  1.408 0.159093
## genderMale       8.648e-01  2.375e+00  1.045e-01  8.272 < 2e-16 ***
## age            -2.201e-02  9.782e-01  4.388e-03 -5.015 5.30e-07 ***
## drugatorvastatin -4.182e-01  6.582e-01  2.020e-01 -2.070 0.038438 *
## drugbisoprolol   2.671e-01  1.306e+00  4.835e-01  0.552 0.580654
## drugcanagliflozin -1.861e+00  1.555e-01  1.018e+00 -1.827 0.067652 .
## drugcarvedilol   -5.718e-02  9.444e-01  2.790e-01 -0.205 0.837608
## drugirbesartan    1.078e+00  2.939e+00  4.056e-01  2.658 0.007851 **
## druglosartan     -4.564e-01  6.335e-01  2.166e-01 -2.107 0.035094 *
## druglovastatin    3.159e-02  1.032e+00  3.214e-01  0.098 0.921698
## drugmetformin    -4.624e-01  6.298e-01  2.369e-01 -1.951 0.051027 .
## drugmetoprolol    2.772e-01  1.319e+00  2.031e-01  1.365 0.172219
## drugnebivolol    -3.372e-01  7.138e-01  3.410e-01 -0.989 0.322845
## drugolmesartan   -2.759e-01  7.589e-01  3.549e-01 -0.777 0.436893
## drugpitavastatin -1.256e+01  3.505e-06  6.667e+02 -0.019 0.984968
## drugpravastatin   6.537e-01  1.923e+00  2.344e-01  2.789 0.005285 **
## drugrosuvastatin -9.813e-01  3.748e-01  2.931e-01 -3.348 0.000813 ***
## drugsimvastatin  -4.339e-01  6.480e-01  2.145e-01 -2.022 0.043139 *
## drugtelmisartan   1.211e+00  3.357e+00  4.506e-01  2.688 0.007184 **
## drugvalsartan    -8.765e-03  9.913e-01  2.330e-01 -0.038 0.969990
## dosage           3.469e-04  1.000e+00  1.331e-04  2.607 0.009141 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
##              exp(coef) exp(-coef) lower .95 upper .95
## creatinine      9.948e-01  1.005e+00  0.76418  1.2951
## DBP             1.009e+00  9.909e-01  1.00076  1.0177
## glucose          1.029e+00  9.722e-01  0.97556  1.0845
## HGB             9.179e-01  1.089e+00  0.86437  0.9748
## ldl             9.911e-01  1.009e+00  0.98802  0.9941
## SBP            9.910e-01  1.009e+00  0.98456  0.9974
## raceBlack       6.034e-01  1.657e+00  0.37675  0.9665
## raceHispanic    6.748e-01  1.482e+00  0.27829  1.6361
## raceUnknown     2.311e+00  4.328e-01  1.49231  3.5777
## raceWhite       1.307e+00  7.649e-01  0.90029  1.8986
## genderMale      2.375e+00  4.211e-01  1.93460  2.9145
## age            9.782e-01  1.022e+00  0.96986  0.9867
```

```
## drugatorvastatin 6.582e-01 1.519e+00 0.44303 0.9780
## drugbisoprolol 1.306e+00 7.656e-01 0.50632 3.3697
## drugcanagliflozin 1.555e-01 6.429e+00 0.02114 1.1446
## drugcarvedilol 9.444e-01 1.059e+00 0.54664 1.6317
## drugirbesartan 2.939e+00 3.402e-01 1.32746 6.5080
## druglosartan 6.335e-01 1.578e+00 0.41437 0.9686
## druglovastatin 1.032e+00 9.689e-01 0.54974 1.9377
## drugmetformin 6.298e-01 1.588e+00 0.39583 1.0021
## drugmetoprolol 1.319e+00 7.579e-01 0.88622 1.9644
## drugnebivolol 7.138e-01 1.401e+00 0.36583 1.3927
## drugolmesartan 7.589e-01 1.318e+00 0.37851 1.5215
## drugpitavastatin 3.505e-06 2.853e+05 0.00000 Inf
## drugpravastatin 1.923e+00 5.201e-01 1.21451 3.0439
## drugrosuvastatin 3.748e-01 2.668e+00 0.21103 0.6657
## drugsimvastatin 6.480e-01 1.543e+00 0.42557 0.9867
## drugtelmisartan 3.357e+00 2.979e-01 1.38832 8.1192
## drugvalsartan 9.913e-01 1.009e+00 0.62787 1.5650
## dosage 1.000e+00 9.997e-01 1.00009 1.0006
##
## Concordance= 0.741 (se = 0.012 )
## Likelihood ratio test= 334.1 on 30 df, p=<2e-16
## Wald test = 318.8 on 30 df, p=<2e-16
## Score (logrank) test = 347.7 on 30 df, p=<2e-16
```

The coefficient of drug canagliflozin is infinite, so it is further removed in the third round of modeling.

```
# Cox regression model_3rd round
CKDdata_2 <- CKDdata_1 %>% filter(drug != "canagliflozin")

CKDdata_2$id <- NULL

split_data_2 <- sample(1:nrow(CKDdata_2), 0.8 * nrow(CKDdata_2), FALSE)
training_set_2 <- CKDdata_2[split_data_2,]
test_set_2 <- CKDdata_2[-split_data_2,]

cox3 <- coxph(Surv(time, stage) ~ ., data=training_set_2)
summary(cox3)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ ., data = training_set_2)
##
## n= 1800, number of events= 680
## (6508 observations deleted due to missingness)
##
##              coef exp(coef) se(coef)      z Pr(>|z|)
## creatinine    0.0880512  1.0920440  0.1371982  0.642  0.52102
## DBP           0.0037049  1.0037118  0.0043181  0.858  0.39089
## glucose       0.0170356  1.0171815  0.0278441  0.612  0.54066
## HGB          -0.0544790  0.9469784  0.0306406 -1.778  0.07540 .
## ldl          -0.0085827  0.9914540  0.0016195 -5.300 1.16e-07 ***
## SBP          -0.0117194  0.9883490  0.0034369 -3.410  0.00065 ***
## raceBlack    -0.5305784  0.5882646  0.2685789 -1.976  0.04821 *
## raceHispanic  0.1510768  1.1630860  0.3852515  0.392  0.69495
```

```

## raceUnknown      1.1476517   3.1507852   0.2476037   4.635  3.57e-06 ***
## raceWhite        0.5028371   1.6534055   0.2190386   2.296  0.02170 *
## genderMale       0.9864832   2.6817867   0.1065609   9.257  < 2e-16 ***
## age             -0.0255756   0.9747487   0.0043466  -5.884  4.00e-09 ***
## drugatorvastatin -0.3265845   0.7213834   0.2044777  -1.597  0.11023
## drugbisoprolol   -0.3114622   0.7323753   0.6071663  -0.513  0.60797
## drugcarvedilol    0.1095011   1.1157214   0.2866500   0.382  0.70246
## drugirbesartan    0.9698376   2.6375162   0.4525334   2.143  0.03210 *
## druglosartan     -0.4054258   0.6666929   0.2200910  -1.842  0.06546 .
## druglovastatin    0.0329942   1.0335446   0.3417684   0.097  0.92309
## drugmetformin     -0.4168363   0.6591288   0.2395880  -1.740  0.08189 .
## drugmetoprolol    0.2938363   1.3415642   0.2073118   1.417  0.15638
## drugnebivolol     -0.2108045   0.8099324   0.3816879  -0.552  0.58075
## drugolmesartan    -0.5114964   0.5995977   0.3446601  -1.484  0.13779
## drugpitavastatin  0.0394841   1.0402740   1.0204232   0.039  0.96913
## drugpravastatin   0.6053877   1.8319622   0.2362488   2.563  0.01039 *
## drugrosuvastatin  -0.7853018   0.4559821   0.2884735  -2.722  0.00648 **
## drugsimvastatin   -0.3525527   0.7028916   0.2143937  -1.644  0.10009
## drugtelmisartan    0.8847006   2.4222591   0.5391208   1.641  0.10080
## drugvalsartan     0.1644554   1.1787510   0.2337780   0.703  0.48176
## dosage            0.0003606   1.0003607   0.0001357   2.658  0.00785 **
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
##               exp(coef) exp(-coef) lower .95 upper .95
## creatinine      1.0920      0.9157      0.8346      1.4290
## DBP             1.0037      0.9963      0.9953      1.0122
## glucose         1.0172      0.9831      0.9632      1.0742
## HGB             0.9470      1.0560      0.8918      1.0056
## ldl             0.9915      1.0086      0.9883      0.9946
## SBP            0.9883      1.0118      0.9817      0.9950
## raceBlack       0.5883      1.6999      0.3475      0.9958
## raceHispanic    1.1631      0.8598      0.5466      2.4748
## raceUnknown     3.1508      0.3174      1.9394      5.1189
## raceWhite       1.6534      0.6048      1.0763      2.5400
## genderMale      2.6818      0.3729      2.1763      3.3047
## age            0.9747      1.0259      0.9665      0.9831
## drugatorvastatin 0.7214      1.3862      0.4832      1.0770
## drugbisoprolol  0.7324      1.3654      0.2228      2.4074
## drugcarvedilol  1.1157      0.8963      0.6361      1.9568
## drugirbesartan  2.6375      0.3791      1.0864      6.4031
## druglosartan    0.6667      1.4999      0.4331      1.0263
## druglovastatin  1.0335      0.9675      0.5290      2.0195
## drugmetformin   0.6591      1.5172      0.4121      1.0542
## drugmetoprolol  1.3416      0.7454      0.8936      2.0141
## drugnebivolol   0.8099      1.2347      0.3833      1.7114
## drugolmesartan  0.5996      1.6678      0.3051      1.1782
## drugpitavastatin 1.0403      0.9613      0.1408      7.6866
## drugpravastatin 1.8320      0.5459      1.1530      2.9108
## drugrosuvastatin 0.4560      2.1931      0.2591      0.8026
## drugsimvastatin 0.7029      1.4227      0.4617      1.0700
## drugtelmisartan 2.4223      0.4128      0.8420      6.9682
## drugvalsartan   1.1788      0.8484      0.7455      1.8639
## dosage          1.0004      0.9996      1.0001      1.0006

```

```
##
## Concordance= 0.753 (se = 0.012 )
## Likelihood ratio test= 352.8 on 29 df, p=<2e-16
## Wald test = 343.7 on 29 df, p=<2e-16
## Score (logrank) test = 369.8 on 29 df, p=<2e-16
```

The coefficient of drug pitavastatin is infinite, so it is further removed in the fourth round of modeling.

```
# Cox regression model_4th round
CKDdata_3 <- CKDdata_2 %>% filter(drug != "pitavastatin")

CKDdata_3$id <- NULL

split_data_3 <- sample(1:nrow(CKDdata_3), 0.8 * nrow(CKDdata_3), FALSE)
training_set_3 <- CKDdata_3[split_data_3,]
test_set_3 <- CKDdata_3[-split_data_3,]

cox4 <- coxph(Surv(time, stage) ~ ., data=training_set_3)
summary(cox4)
```

```
## Call:
## coxph(formula = Surv(time, stage) ~ ., data = training_set_3)
##
## n= 1809, number of events= 686
## (6482 observations deleted due to missingness)
##
##              coef exp(coef) se(coef)      z Pr(>|z|)
## creatinine      0.0627815  1.0647941  0.1394662  0.450 0.652598
## DBP             0.0050122  1.0050248  0.0043986  1.139 0.254496
## glucose         0.0562277  1.0578385  0.0272019  2.067 0.038730 *
## HGB            -0.0667149  0.9354619  0.0307586 -2.169 0.030084 *
## ldl            -0.0082383  0.9917955  0.0015972 -5.158 2.5e-07 ***
## SBP            -0.0114686  0.9885969  0.0034036 -3.370 0.000753 ***
## raceBlack      -0.6781487  0.5075558  0.2483001 -2.731 0.006311 **
## raceHispanic   -0.2727672  0.7612700  0.4054815 -0.673 0.501139
## raceUnknown     0.8106716  2.2494181  0.2306889  3.514 0.000441 ***
## raceWhite       0.1926585  1.2124687  0.1971745  0.977 0.328521
## genderMale      0.8930695  2.4426157  0.1063635  8.396 < 2e-16 ***
## age            -0.0236176  0.9766591  0.0044966 -5.252 1.5e-07 ***
## drugatorvastatin -0.5290989  0.5891356  0.2032403 -2.603 0.009233 **
## drugbisoprolol  -0.1239203  0.8834503  0.4863877 -0.255 0.798896
## drugcarvedilol  -0.2524897  0.7768642  0.2782479 -0.907 0.364181
## drugirbesartan   1.0548905  2.8716607  0.4253158  2.480 0.013129 *
## druglosartan    -0.5444712  0.5801485  0.2137100 -2.548 0.010843 *
## druglovastatin  -0.0443809  0.9565895  0.3503911 -0.127 0.899209
## drugmetformin   -0.3710300  0.6900232  0.2385713 -1.555 0.119894
## drugmetoprolol   0.2373848  1.2679289  0.2011274  1.180 0.237893
## drugnebivolol   -0.3191170  0.7267905  0.3679718 -0.867 0.385815
## drugolmesartan  -0.4620584  0.6299855  0.3332718 -1.386 0.165615
## drugpravastatin  0.5059007  1.6584787  0.2348686  2.154 0.031242 *
## drugrosuvastatin -0.9399611  0.3906430  0.2924886 -3.214 0.001311 **
## drugsimvastatin -0.4964376  0.6086952  0.2138145 -2.322 0.020243 *
## drugtelmisartan  1.1584680  3.1850499  0.5376269  2.155 0.031179 *
```

```
## drugvalsartan      -0.1259861  0.8816271  0.2418180 -0.521  0.602370
## dosage             0.0002609  1.0002609  0.0001427  1.828  0.067553 .
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
##
##               exp(coef) exp(-coef) lower .95 upper .95
## creatinine       1.0648      0.9391   0.8101   1.3995
## DBP              1.0050      0.9950   0.9964   1.0137
## glucose          1.0578      0.9453   1.0029   1.1158
## HGB              0.9355      1.0690   0.8807   0.9936
## ldl              0.9918      1.0083   0.9887   0.9949
## SBP              0.9886      1.0115   0.9820   0.9952
## raceBlack        0.5076      1.9702   0.3120   0.8257
## raceHispanic     0.7613      1.3136   0.3439   1.6853
## raceUnknown      2.2494      0.4446   1.4312   3.5354
## raceWhite        1.2125      0.8248   0.8238   1.7845
## genderMale       2.4426      0.4094   1.9830   3.0088
## age              0.9767      1.0239   0.9681   0.9853
## drugatorvastatin 0.5891      1.6974   0.3956   0.8774
## drugbisoprolol   0.8835      1.1319   0.3405   2.2919
## drugcarvedilol   0.7769      1.2872   0.4503   1.3403
## drugirbesartan   2.8717      0.3482   1.2477   6.6094
## druglosartan     0.5801      1.7237   0.3816   0.8820
## druglovastatin   0.9566      1.0454   0.4814   1.9010
## drugmetformin    0.6900      1.4492   0.4323   1.1014
## drugmetoprolol   1.2679      0.7887   0.8549   1.8806
## drugnebivolol    0.7268      1.3759   0.3533   1.4950
## drugolmesartan   0.6300      1.5873   0.3278   1.2106
## drugpravastatin  1.6585      0.6030   1.0466   2.6280
## drugrosuvastatin 0.3906      2.5599   0.2202   0.6930
## drugsimvastatin  0.6087      1.6429   0.4003   0.9255
## drugtelmisartan  3.1850      0.3140   1.1104   9.1358
## drugvalsartan    0.8816      1.1343   0.5488   1.4162
## dosage           1.0003      0.9997   1.0000   1.0005
##
## Concordance= 0.738 (se = 0.013 )
## Likelihood ratio test= 317.2 on 28 df,  p=<2e-16
## Wald test              = 315.7 on 28 df,  p=<2e-16
## Score (logrank) test = 337.2 on 28 df,  p=<2e-16
```

Prediction

```
# Remove NA
test_set_3_n <- na.omit(test_set_3)

# Make prediction
pred <- predict(cox4, newdata=test_set_3_n, type="survival")
observed_result <- test_set_3_n$stage
predict_result <- ifelse(pred>0.5,0,1)
table(predict_result, observed_result)
```

```
##              observed_result
## predict_result    0      1
##              0 258   91
##              1  33   84
```

Confusion Matrix and ROC

```
# Confusion Matrix
MLmetrics::Accuracy(predict_result, test_set_3_n$stage)
```

```
## [1] 0.7339056
```

```
MLmetrics::Specificity(predict_result, test_set_3_n$stage)
```

```
## [1] 0.7179487
```

```
MLmetrics::Sensitivity(predict_result, test_set_3_n$stage)
```

```
## [1] 0.739255
```

```
# ROC
require(ROCR)
```

```
## Loading required package: ROCR
```

```
require(pROC)
```

```
## Loading required package: pROC
```

```
## Type 'citation("pROC")' for a citation.
```

```
##
## Attaching package: 'pROC'
```

```
## The following objects are masked from 'package:stats':
##
##      cov, smooth, var
```




```
rocplot <- function(pred, truth, ...) {
  predob = prediction(pred, truth)
  perf = performance(predob, "tpr", "fpr")
  plot(perf, ...)
  area <- auc(truth, pred)
  area <- format(round(area, 4), nsmall = 4)
  text(x=0.8, y=0.1, labels = paste("AUC =", area))

  # the reference x=y line
  segments(x0=0, y0=0, x1=1, y1=1, col="gray", lty=2)
}

rocplot(predict_result, observed_result, col="blue")
```


```
## Setting levels: control = 0, case = 1
```

```
## Setting direction: controls < cases
```



Plot test set results

```
ggplot() +
  geom_point(aes(x = test_set_3_n$time, y = test_set_3_n$stage),
             colour = 'red') +
  geom_line(aes(x = test_set_3_n$time, y = predict(cox4, newdata = test_set_3_n)),
            colour = 'blue') +
  ggtitle('Cox regression (Test set)') +
  xlab('Days') +
  ylab('Survival probability')
```



Based on the results, glucose, HGB, LDL, SBP, race, gender, age, some drugs, and drug doseage are significant risk factors ($p < 0.05$) contributing to the progression of CKD. The Cox regression model is a good predictor in this CKD data set. A significant advantage of this Cox model is its ease of use. The algorithm automatically calculates simultaneously the effect of several risk factors on survival time.

Random Forest Model

Finally, I use the `ranger()` function to fit a Random Forest Ensemble model to the CKD dataset. `ranger()` builds a model for each observation in the dataset. The next block of code plots random curves in the training and test sets, along with a curve that represents the global average for all of the patients.

```
# Split dataset into training and test sets
split_data <- sample(1:nrow(CKDdata), 0.8 * nrow(CKDdata), FALSE)
training_set <- CKDdata[split_data,]
test_set <- CKDdata[-split_data,]

# Build random forest model
r_fit <- ranger(Surv(time, stage) ~ creatinine+gender+race+age, data=training_set,
mtry = 4, importance="permutation", splitrule="extratrees",verbose=TRUE)
```

```
## Computing permutation importance.. Progress: 35%. Estimated remaining time: 58
seconds.
## Computing permutation importance.. Progress: 69%. Estimated remaining time: 28
seconds.
```


```
prob_pred <- predict(r_fit, data=test_set, mtry = 4, importance="permutation", spl
itrule="extratrees",verbose=TRUE)

# Average the survival models
CKDprogress_time <- r_fit$unique.death.times
surv_probability <- data.frame(r_fit$survival)
avg_probability <- sapply(surv_probability, mean)

# Plot the survival models
# Training set
plot(r_fit$unique.death.times,r_fit$survival[1,],
     type = "l",
     ylim = c(0,1),
     col = "red",
     xlab = "Days",
     ylab = "Survival Probability",
     main = "Patient Survival Curves (Training set)")

cols <- colors()
for (n in sample(c(2:dim(training_set)[1]), 20)){
  lines(r_fit$unique.death.times, r_fit$survival[n,], type = "l", col = cols[n])
}
lines(CKDprogress_time, avg_probability, lwd = 2)
legend(100, 0.4, legend = c('Average = black'))
```


Patient Survival Curves (Training set)



```
# Test set
plot(r_fit$unique.death.times, r_fit$survival[1,],
     type = "l",
     ylim = c(0,1),
     col = "red",
     xlab = "Days",
     ylab = "Survival Probability",
     main = "Patient Survival Curves (Test set)")

cols <- colors()
for (n in sample(c(2:dim(test_set)[1]), 20)){
  lines(r_fit$unique.death.times, r_fit$survival[n,], type = "l", col = cols[n])
}
lines(CKDprogress_time, avg_probability, lwd = 2)
legend(100, 0.4, legend = c('Average = black'))
```

Patient Survival Curves (Test set)



The next block of illustrates how ranger () ranks variable importance.

```
# Variable importance
vi <- data.frame(sort(round(r_fit$variable.importance, 4), decreasing = TRUE))
names(vi) <- "importance"
head(vi)
```

```
##           importance
## age             0.3348
## creatinine      0.2762
## gender          0.2242
## race            0.1649
```

```
cat("Prediction Error = 1 - Harrell's c-index = ", r_fit$prediction.error)
```

```
## Prediction Error = 1 - Harrell's c-index = 0.1076468
```

I notice that ranger () flags age, creatinine, gender, and race are the most important factors of CKD progression. Age, gender, and race are the same variables with the p-values less than 0.05 in the Cox regression model. Furthermore, ranger() also computes Harrell's c-index. This is a generalization of the ROC curve, which reduces to the Wilcoxon-Mann-Whitney statistics for binary variables, which in turn, is

equivalent to computing the area under the ROC curve. Here, the prediction error is 0.11 and the ROC value of 0.89 would normally be pretty good for a first try. But ranger () doesn't do anything to address the time varying coefficients, which is apparently a challenge.

In summary, for this CKD data set, I would choose a carefully constructed Cox Regression model that takes into account all varying coefficients. Tree-based models for survival analysis will be useful in dealing with very large data sets.