

Self-propelled particles as an active matter system


Étienne Fodor,^{1,2} Cesare Nardini,¹ Elsen Tjhung,¹ Michael E. Cates,¹
Julien Tailleur,² Paolo Visco,² Frédéric van Wijland²

1. Department of Applied Mathematics and Theoretical Physics, University of Cambridge
2. Laboratoire Matière et Systèmes Complexes, Université Paris Diderot

School of Mathematical Sciences
Queen Mary University of London

Introduction


Active matter: interacting self-propelled particles



A. Walther and A. H. E. Müller, Soft Matter 4, 663 (2008)


Introduction

Light-induced clusters of colloids



J. Palacci *et al.*, Science **339**, 936 (2013)


Simulated interacting active colloids



G. S. Redner *et al.*,
Phys. Rev. Lett. **110**, 055701 (2013)

Introduction

Run-and-tumble and active Brownian particles



M. E . Cates and J. Tailleur, Ann. Rev. CMP 6, 219 (2015)

Introduction

Many-body dynamics

$$0 = -\nabla_i U - \gamma \frac{dx_i}{dt} + \xi_i$$

Equilibrium fluctuations ξ_i

No memory and Gaussian

Boltzmann distribution

$$P_S \sim e^{-U/T}$$

Introduction

Many-body dynamics

$$\frac{dx_i}{dt} = -\mu \nabla_i U + v_i$$

Nonequilibrium self-propulsion v_i

Memory and non-Gaussian

Non-Boltzmann distribution

Introduction

Main questions of interest

- Phase diagram: equilibrium mapping

J. Tailleur *et al.*, Phys. Rev. Lett. **100**, 218103 (2008)

- Equation of state: pressure

A. P. Solon *et al.*, Phys. Rev. Lett. **114**, 198301 (2015)

- Nonequilibrium properties

Outline

1 Interacting active particles

- Minimal model of self-propulsion
- How far from equilibrium?
- Effective equilibrium regime

Outline

1 Interacting active particles

- Minimal model of self-propulsion
- How far from equilibrium?
- Effective equilibrium regime

2 Active field theory

- Coarse-grained dynamics
- Spatial structure of nonequilibrium

Outline

1 Interacting active particles

- Minimal model of self-propulsion
- How far from equilibrium?
- Effective equilibrium regime

2 Active field theory

- Coarse-grained dynamics
- Spatial structure of nonequilibrium

Interacting active particles

Many-body dynamics

$$\frac{dx_i}{dt} = -\mu \nabla_i U + v_i$$

Nonequilibrium self-propulsion v_i

Memory and Gaussian

G. Szamel et al., Phys. Rev. E **91**, 062304 (2015)


C. Maggi et al., Sci. Rep. **5**, 10724 (2015)

T. F. F. Farage et al., Phys. Rev. E **91**, 042310 (2015)

Persistence time $\tau \rightarrow$ Equilibrium $\tau = 0$

Interacting active particles

Interacting repulsive particles



ÉF et al., Phys. Rev. Lett. 117, 038103 (2016)

Interacting active particles

Comparing forward and backward dynamics

Quantifying irreversibility

$$\mathcal{S} = \lim_{t \rightarrow \infty} \frac{1}{t} \ln \frac{\mathcal{P}}{\mathcal{P}^R}$$

Entropy production rate

Equilibrium $\mathcal{S} = 0$

Interacting active particles

Passive Brownian particles + non-conservative force F_i

$$S = \frac{\left\langle F_i \frac{dx_i}{dt} \right\rangle}{T} = \frac{\text{power of } F_i}{\text{temperature}}$$

Self-propelled particles

$$S = \frac{\mu \tau^2}{2T} \left\langle \nabla_i^3 U \left(\frac{dx_i}{dt} \right)^3 \right\rangle$$

Interacting active particles

Péclet number

$$Pe = \frac{\text{persistence length}}{\text{interaction range}}$$

$$Pe = \sqrt{\frac{\text{persistence time}}{\text{relaxation time}}}$$

Competition self-propulsion/interaction

Interacting active particles

Perturbative treatment in Péclet number

- $P_S \sim e^{-U/T} [1 + \mathcal{O}(\text{Pe}^2)]$

- $\left\langle \nabla_i^3 U \left(\frac{dx_i}{dt} \right)^3 \right\rangle = \mathcal{O}(\text{Pe}^3)$


Effective equilibrium regime

$\left\{ \begin{array}{l} \text{Non-Boltzmann statistics} \\ \text{Time reversal symmetry} \end{array} \right.$

Interacting active particles

Effective potential

$$U_{\text{eff}} = U + \tau \left[\frac{(\nabla_i U)^2}{2} - T \nabla_i^2 U \right]$$



Original pair-wise potential

$$U = \frac{1}{2} \sum_{\{i,j\}=1}^N \phi(r_i - r_j)$$

$$\phi \sim \frac{1}{x^{12}}$$

Interacting active particles

External perturbation $f_i(t)$

$$\langle \delta x_i(t) \rangle = \int_0^t R(t-s) f_i(s) ds$$

Response function $R(t)$

Passive Brownian particles

$$R(t) = -\frac{1}{T} \frac{d}{dt} \langle x_i(t) x_i(0) \rangle$$

Fluctuation-dissipation theorem

Interacting active particles


Fluctuation-dissipation relation

$$R(t) = -\frac{1}{T} \frac{d}{dt} \left\langle x_i(t)x_i(0) + \tau^2 \frac{dx_i(t)}{dt} \frac{dx_i(0)}{dt} \right\rangle$$

Probing effective equilibrium

Interacting active particles

Numerical simulations: 3 values of T



Susceptibility

$$\chi(t) = \frac{1}{N} \int_0^t R(t-s) ds$$

Correlation

$$C_{\text{eff}}(t) = \frac{1}{N} \left\langle x_i^2(t) + \tau^2 \left[\frac{dx_i(t)}{dt} \right]^2 \right\rangle - \frac{1}{N} \left\langle x_i(t)x_i(0) + \tau^2 \frac{dx_i(t)}{dt} \frac{dx_i(0)}{dt} \right\rangle$$

Outline

1 Interacting active particles

- Minimal model of self-propulsion
- How far from equilibrium?
- Effective equilibrium regime

2 Active field theory

- Coarse-grained dynamics
- Spatial structure of nonequilibrium

Active field theory

Fluctuating density field

$$\rho(x, t) = \sum_{i=1}^N \delta[x - x_i(t)]$$

Active model B

$$\partial_t \rho = \nabla \left[\nabla \left(\frac{\delta \mathcal{F}}{\delta \rho} + \mu_A \right) + \Lambda \right]$$

Equilibrium fluctuations Λ

Active chemical potential μ_A

Active field theory

Phase separation

$$\mathcal{F} = \int \left[\frac{a}{2} \rho^2 + \frac{b}{4} \rho^4 + \frac{\kappa}{2} |\nabla \rho|^2 \right] dx$$


Cahn-Hilliard free energy

Active chemical potential

$$\mu_A = \lambda |\nabla \rho|^2$$

Active field theory

Shift of coexisting phase densities



R. Wittkowski *et al.*, Nat. Com. **5**, 4351 (2014)

Active field theory

Entropy production rate

$$\mathcal{S} = \frac{1}{T} \int \left\langle \mu_A \partial_t \rho \right\rangle dx$$

Spatial density of entropy production


$$\sigma = \frac{\lambda}{T} \left\langle |\nabla \rho|^2 \partial_t \rho \right\rangle$$

Probing nonequilibrium from field profile


Active field theory

Spatial structure of entropy production rate


Field profile



Entropy density



Entropy density



C. Nardini *et al.*, arXiv:1610.06112

Active field theory

External perturbation $h(x, t)$

$$\langle \delta\rho(x, t) \rangle = \int R(x - y, t - s) h(y, s) dy ds$$

Response function $R(x, t)$

Equilibrium dynamics


$$R(x - y, t) = -\frac{1}{T} \frac{d}{dt} \langle \rho(x, t) \rho(y, 0) \rangle$$

Fluctuation-dissipation theorem

Active field theory

Violation of fluctuation-dissipation

$$\omega C(k, \omega) / 2T R(k, \omega)$$



Response



$$R(k, \omega) = \int R(x, t) e^{ikx} \sin(\omega t) dt dx$$

Correlation

$$C(k, \omega) = \int \langle \rho(x, t) \rho(0, 0) \rangle e^{i(kx + \omega t)} dt dx$$

Active field theory

Violation of fluctuation-dissipation



Active field theory

Spectral decomposition of entropy production

$$\mathcal{S} = \int \sigma(k, \omega) \frac{dk d\omega}{(2\pi)^{d+1}}$$

Violation of fluctuation-dissipation

$$\sigma(k, \omega) = \frac{\omega}{Tk^2} [\omega C(k, \omega) - 2TR(k, \omega)]$$

Conclusion

Nonequilibrium properties of active systems

- Interacting active particles
 - Effective equilibrium regime
- Active field theory
 - Spatial structure of nonequilibrium

References: Phys. Rev. Lett. **117**, 038103 (2016) | [arXiv:1610.06112](https://arxiv.org/abs/1610.06112)

Conclusion

Future directions

- Other models of self-propulsion
 - Non-Gaussian without memory
- Hydrodynamic description
 - Dynamics of passive tracer

Interacting active particle

Density waves around an obstacle

