

Self-propelled particles as an active matter system

Étienne Fodor,^{1,2} Cesare Nardini,³ Paolo Visco,¹
Julien Tailleur,¹ Frédéric van Wijland^{1,2}

1. Laboratoire Matière et Systèmes Complexes, Université Paris Diderot


2. Yukawa Institute of Theoretical Physics, University of Kyoto

3. Department of Applied Mathematics and Theoretical Physics, University of Cambridge

Yukawa Institute of Theoretical Physics
University of Kyoto


Introduction

Light-induced clusters of colloids




J. Palacci, S. Sacanna, A. P. Steinberg,
D. J. Pine, and P. M. Chaikin,
Science **339**, 936 (2013)

Simulated interacting active colloids



G. S. Redner, M. F. Hagan, and A. Baskaran,
Phys. Rev. Lett. **110**, 055701 (2013)


Dense suspension of active colloids: glassy dynamics



L. Berthier, Phys. Rev. Lett. **112**, 220602 (2014)

Introduction

Run-and-tumble and active Brownian particles



M. E. Cates and J. Tailleur, Ann. Rev. CMP **6**, 219 (2015)


Competition self-propulsion/interaction

Péclet number

$$Pe = \frac{\text{persistence length}}{\text{interaction range}}$$

$$Pe = \sqrt{\frac{\text{persistence time}}{\text{relaxation time}}}$$

Introduction



T. F. F. Farage, P. Krinninger, and J. M. Brader, Phys. Rev. E **91**, 042310 (2015)

Brownian particles

$$\gamma \dot{\mathbf{r}}_i = - \sum_j \nabla_i \phi(\mathbf{r}_i - \mathbf{r}_j) + \boldsymbol{\xi}_i$$

Mobility $\mu = \gamma^{-1}$

Equilibrium fluctuations

$$\langle \boldsymbol{\xi}_i(t) \boldsymbol{\xi}_j(0) \rangle = 2\gamma T \delta_{ij} \delta(t)$$

$$P_S = \exp \left[-\frac{1}{2T} \sum_{i,j} \phi(\mathbf{r}_i - \mathbf{r}_j) \right]$$

Self-propelled particles

$$\dot{\mathbf{r}}_i = -\mu \sum_j \nabla_i \phi(\mathbf{r}_i - \mathbf{r}_j) + \mathbf{v}_i$$

Nonequilibrium fluctuations \mathbf{v}

- Random direction with fixed norm
- Memory and non-Gaussian

Main questions of interest

- Equation of state: temperature, pressure
- Phase diagram: stationary measure
- Structural relaxation: time scales

- 1 Searching for a minimal model of self-propulsion

- 1 Searching for a minimal model of self-propulsion
- 2 Quantifying nonequilibrium

- 1 Searching for a minimal model of self-propulsion
- 2 Quantifying nonequilibrium
- 3 Effective equilibrium regime

- 1 Searching for a minimal model of self-propulsion
- 2 Quantifying nonequilibrium
- 3 Effective equilibrium regime

Searching for a minimal model of self-propulsion

Overdamped dynamics

$$\dot{\mathbf{r}}_i = -\mu \nabla_i U + \mathbf{v}_i$$

External potential + pair-wise interactions

$$U = \sum_i V_{\text{ext}}(\mathbf{r}_i) + \frac{1}{2} \sum_{i,j} \phi(\mathbf{r}_i - \mathbf{r}_j)$$

Searching for a minimal model of self-propulsion

Overdamped dynamics

$$\dot{\mathbf{r}}_i = -\mu \nabla_i U + \mathbf{v}_i$$

Gaussian fluctuations

$$\langle \mathbf{v}_i(t) \mathbf{v}_j(0) \rangle = \delta_{ij} \frac{\mu T}{\tau} e^{-|t|/\tau}$$

Equilibrium reference

$$\langle \mathbf{v}_i(t) \mathbf{v}_j(0) \rangle \xrightarrow{\tau \rightarrow 0} 2\mu T \delta_{ij} \delta(t)$$

Searching for a minimal model of self-propulsion

No interaction: harmonic oscillator

$$\dot{r} = -\kappa\mu r + v$$

$$\langle v(t)v(0) \rangle = \frac{\mu T}{\tau} e^{-|t|/\tau}$$

Stationary distribution

$$P_S = \exp \left[- (1 + \tau\kappa\mu) \frac{\kappa r^2}{2T} \right]$$

Searching for a minimal model of self-propulsion

No interaction: harmonic oscillator

$$\dot{r} = -\kappa\mu r + v$$

$$\langle v(t)v(0) \rangle = \frac{\mu T}{\tau} e^{-|t|/\tau}$$

Stationary distribution

$$P_s = \exp \left[- \left(1 + \text{Pe}^2 \right) \frac{\kappa r^2}{2T} \right]$$

Searching for a minimal model of self-propulsion

No interaction: harmonic oscillator

$$\dot{r} = -\kappa\mu r + v$$

$$\dot{v} = -\frac{v}{\tau} + \xi$$

$$\langle \xi(t)\xi(0) \rangle = \frac{2\mu T}{\tau^2} \delta(t)$$

Equilibrium underdamped dynamics

$$\dot{r} = p$$

$$\dot{p} = -\frac{p}{\tau} - \frac{\mu\kappa}{\tau} \left(1 + \frac{\kappa\tau}{\gamma} \right) r + \xi$$

Searching for a minimal model of self-propulsion

Interacting particles

$$\dot{r}_i = -\mu \nabla_i U + v_i$$

$$\dot{v}_i = -\frac{v_i}{\tau} + \xi_i$$

$$\langle \xi_i(t) \xi_j(0) \rangle = \frac{2\mu T}{\tau^2} \delta_{ij} \delta(t)$$

Nonequilibrium underdamped dynamics

$$\dot{r}_i = p_i$$

$$\dot{p}_i = -\frac{p_i}{\tau} - \mu p_k \partial_{ik}^2 U - \frac{\mu}{\tau} \nabla_i U + \xi_i$$

Searching for a minimal model of self-propulsion

Previous attempt: neglecting inertia

$$\dot{r}_i = p_i$$


$$\dot{p}_i = -\frac{p_i}{\tau} - \mu p_k \partial_{ik}^2 U - \frac{\mu}{\tau} \nabla_i U + \xi_i$$

Stationary measure

$$P_S = \exp \left[-\frac{U}{T} - \frac{\tau^2}{2T} (\partial_i U)^2 \right] |\det \mathbb{M}|$$

$$\mathbb{M}_{ij} = \delta_{ij} + \tau \mu \partial_{ij}^2 U$$

Searching for a minimal model of self-propulsion



C. Maggi, U. Marini Bettolo Marconi, N. Gnan, and R. Di Leonardo, Sci. Rep. **5**, 10724 (2015)

Searching for a minimal model of self-propulsion

Previous attempt: neglecting inertia

$$\dot{r}_i = p_i$$

$$\dot{p}_i = -\frac{p_i}{\tau} - \mu p_k \partial_{ik}^2 U - \frac{\mu}{\tau} \nabla_i U + \xi_i$$

Mapping to equilibrium dynamics

Outline

- 1 Searching for a minimal model of self-propulsion
- 2 Quantifying nonequilibrium
- 3 Effective equilibrium regime

Entropy production rate

$$\sigma = \lim_{t \rightarrow \infty} \frac{1}{t} \ln \frac{\mathcal{P}}{\mathcal{P}^R}$$

Brownian particles + non-conservative force

$$\gamma \dot{r}_i = F_i - \partial_i U + \xi_i$$

$$\langle \xi_i(t) \xi_j(0) \rangle = 2\gamma T \delta_{ij} \delta(t)$$

$$\sigma = \frac{1}{T} \langle \dot{r}_i F_i \rangle$$

Self-propelled particles

$$\dot{r}_i = -\mu \partial_i U + v_i$$

$$\langle v_i(t) v_j(0) \rangle = \delta_{ij} \frac{\mu T}{\tau} e^{-|t|/\tau}$$

Entropy production

$$\sigma = \frac{\tau^2}{2T} \sum_{ijk} \left\langle \dot{r}_i \dot{r}_j \dot{r}_k \partial_{ijk}^3 U \right\rangle$$

Quantifying nonequilibrium

Péclet number

$$\text{Pe} = \frac{\text{persistence length}}{\text{interaction range}} = \sqrt{\frac{\tau}{\tau_R}}$$

Relaxation time

$$\tau_R = \frac{(\text{interaction range})^2}{\mu T}$$

Self-propelled particles: scaled dynamics

- space unit = interaction range
- time unit = τ/Pe
- energy unit = T

Self-propelled particles: scaled dynamics

$$\dot{r}_i = p_i$$

$$\dot{p}_i = -\frac{1}{\text{Pe}} \left(p_i + \text{Pe}^2 p_j \partial_{ij}^2 U \right) - \partial_i U + \xi_i$$

$$\langle \xi_i(t) \xi_i(0) \rangle = \frac{2}{\text{Pe}} \delta_{ij} \delta(t)$$

Self-propelled particles: scaled dynamics

$$\dot{r}_i = p_i$$

$$\dot{p}_i = -\frac{1}{\text{Pe}} \left(p_i + \text{Pe}^2 p_j \partial_{ij}^2 U \right) - \partial_i U + \xi_i$$

$$\langle \xi_i(t) \xi_i(0) \rangle = \frac{2}{\text{Pe}} \delta_{ij} \delta(t)$$

Maggi *et al.*: Pe finite and $\text{Pe} \rightarrow 0$

Entropy production

$$\sigma = \frac{\text{Pe}^2}{2} \sum_{ijk} \left\langle p_i p_j p_k \partial_{ijk}^3 U \right\rangle$$

Small Pe expansion

$$\sum_{ijk} \left\langle p_i p_j p_k \partial_{ijk}^3 U \right\rangle = c_1 \text{Pe} + c_2 \text{Pe}^2 + c_3 \text{Pe}^3 + \mathcal{O}(\text{Pe}^4)$$

Stationary measure

$$P_S = e^{-U - \frac{p_i^2}{2}} \left\{ 1 - \frac{\text{Pe}^2}{2} \left[(\partial_i U)^2 + \mathcal{L}U \right] + \frac{\text{Pe}^3}{6} p_i \partial_i \mathcal{L}U + \mathcal{O}(\text{Pe}^4) \right\}$$

$$\mathcal{L}U = (p_j p_k - 3\delta_{jk}) \partial_{jk}^2 U$$

- Position-velocity coupling
- Term Pe^3 odd in velocity

Entropy production

$$\sigma = \frac{\text{Pe}^5}{2\mathcal{Z}} \int_r (\partial_{ijk}^3 U)^2 e^{-U} + \mathcal{O}(\text{Pe}^6)$$

No contribution from the Pe^2 term of the measure

Outline


- 1 Searching for a minimal model of self-propulsion
- 2 Quantifying nonequilibrium
- 3 Effective equilibrium regime

Stationary distribution


$$P_S = \exp \left\{ -U - \text{Pe}^2 \left[\frac{(\partial_i U)^2}{2} - \partial_{ii}^2 U \right] + \mathcal{O}(\text{Pe}^4) \right\}$$

- Three-body interactions
- Attractive effects from bare repulsion


Effective equilibrium regime




Effective equilibrium regime



Effective equilibrium regime



Effective equilibrium regime



Brownian particles: external perturbation

$$\gamma \dot{r}_i = f_i - \partial_i U + \xi_i$$

Response function

$$R(t) = \left. \frac{\delta \langle r_i(t) \rangle}{\delta f_i(0)} \right|_{f=0}$$

Fluctuation-dissipation theorem

$$R(t) = -\frac{1}{T} \frac{d}{dt} \langle r_i(t) r_i(0) \rangle$$

Self-propelled particles: external perturbation

$$\dot{r}_i = \mu (f_i - \partial_i U) + v_i$$

Scaled dynamics

$$\dot{p}_i = -\frac{1}{\text{Pe}} (p_i + \text{Pe}^2 p_j \partial_{ij}^2 U) + f_i + \text{Pe} \dot{f}_i - \partial_i U + \xi_i$$

$$\langle \xi_i(t) \xi_i(0) \rangle = \frac{2}{\text{Pe}} \delta_{ij} \delta(t)$$

Fluctuation-dissipation relation with $Pe = 0$

$$R(t) = -\frac{d}{dt} \underbrace{\left[\langle r_i(t)r_i(0) \rangle + Pe^2 \langle \dot{r}_i(t)\dot{r}_i(0) \rangle \right]}_{C(t)}$$

Susceptibility $\chi(t) = \int_0^t R(s)ds$

Correlation $\Delta C(t) = C(0) - C(t)$



Fluctuation-dissipation relation when $Pe \rightarrow 0$

$$R(t) = -\frac{d}{dt} \underbrace{\left[\langle r_i(t)r_i(0) \rangle + Pe^2 \langle \dot{r}_i(t)\dot{r}_i(0) \rangle \right]}_{C(t)} + \mathcal{O}(Pe^4)$$



Susceptibility $\chi(t) = \int_0^t R(s)ds$

Correlation $\Delta C(t) = C(0) - C(t)$

Effective equilibrium regime



Effective equilibrium regime



Minimal model of self-propulsion

Gaussian exponentially correlated fluctuations

- Stationary measure in position-velocity
- Entropy production
- Fluctuation dissipation relation

Future directions

- Collective modes
Hydrodynamic equations, tracer dynamics
- Non-Gaussian white fluctuations | H. Hayakawa
Static structure, phase separation