



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

On the number of crossing-free perfect matchings with designated left and right endpoints

Master Thesis

Robin Leroy

January 19, 2038

Advisors: Prof. Dr. Emo Welzl, Manuel Wettstein
Department of Computer Science, ETH Zürich

Abstract

We give upper bounds on the number of crossing-free straight-edge perfect matchings on a set of n points with designated left and right endpoints for some left-to-right orderings of the designated endpoints. When all left endpoints lie to the left of the right endpoints (matchings across a line), we give an upper bound dependent on the number of extreme points. In particular, when half the points are extreme, there are at most $\mathcal{O}(2.6^n)$ crossing-free perfect matchings across a line. When the designated endpoints form four blocks, left-right-left-right, there are at most $\mathcal{O}(5^n)$ crossing-free perfect matchings.

Contents

Contents	iii
1 Background	3
1.1 Geometric graphs	3
1.2 Asymptotics	4
1.3 A remark on the asymptotics of the binomial coefficient . . .	5
1.4 Triangulations of convex point sets and the Catalan numbers	6
1.5 Crossing-free perfect matchings	7
1.6 Bracket expressions and their relations to perfect matchings .	8
1.7 An optimal lower bound for the number of crossing-free perfect matchings	10
1.8 Matchings across a line	11
2 Analysing the overcounting in the upper bound for matchings across a line	15
3 Highly convex matchings across a line	21
4 Three changes of bracket direction	25
4.1 Improving upon the binomial bound	27
4.2 A recurrence	29
4.3 A better recurrence, and a solution	30
4.3.1 A combinatorial interpretation of spc	32
4.3.2 Tying up some loose ends	33
4.4 Asymptotics	34
5 Outlook	37
A Experimental asymptotics of $C_a \text{spc}(a, b)$	39
B An upper bound for spc using the ζ function	41

CONTENTS

Bibliography

45

Introduction

In the study of plane embeddings of planar graphs, also known as plane graphs, embeddings with non-crossing straight edges (called crossing-free geometric graphs) are of particular interest. For instance, it is known [16] that any optimal solution to the travelling salesman problem in the Euclidean plane has non-crossing straight edges. In 1935, K. Wagner proved [17] that any planar graph has such an embedding.

A natural enumerative question is then, given a vertex set of n points in the Euclidean plane, to bound the number of plane graphs of a certain sort (cycles, matchings, *etc.*) which can be drawn with non-crossing straight edges. For spanning cycles, the question was introduced by M. Newborn and W. O. J. Moser in 1976 [9], and a super-exponential upper bound was given. In 1980, M. Ajtai, V. Chvátal, M. Newborn, and E. Szemerédi [1] proved that there are only exponentially many crossing-free geometric graphs on a given point set.

Exponential bounds for various sorts of graphs were then improved, in particular for triangulations—the maximal crossing-free geometric graphs. One of the interesting features of triangulations is that finding a bound for them yields a bound for the overall number of crossing-free geometric graphs [1]. In addition, bounds on the number of triangulations are of interest in geometric modeling, see for instance [14].

Some special configurations of points are better understood; in particular, for point sets in convex position, the numbers of crossing-free perfect matchings, triangulations, and spanning cycles are known. In fact, the study of point sets in convex position vastly predates the general problem, with the number of triangulations given by Euler in 1751 [7].

For perfect matchings, an optimal lower bound of $C_{\frac{n}{2}}$ was obtained as early as 1995 by A. García, M. Noy, and J. Tejel [8]; upper bounds have been harder to come by, with the best one yet being asymptotically 10.05^n , given

by M. Sharir and E. Welzl in 2005 [13] In that same work, an upper bound of 5.83^n is given for the number of crossing-free perfect matchings on a point set whose points are designated as left or right endpoints; moreover, an upper bound of 4^n is given when all the left endpoints are to the left of the right endpoints (matchings across a line).

In this thesis, we start by giving an overview of some of the existing results, in particular a way to classify arrangements of designated left and right endpoints, and we restate the proof of the bound for matchings across a line by M. Sharir and E. Welzl (chapter 1). We then study in detail the inequalities in that bound and the objects it actually counts, yielding criteria that will allow us to tighten the bound in specific cases (chapter 2). Using these criteria, we give a bound for matchings across a line that depends on the number of points in the interior of the convex hull (chapter 3). We then give new bounds on the number of crossing-free perfect matchings on a point set with designated left and right endpoints for a specific family of left-to-right orderings of the left and right endpoints (chapter 4).

Chapter 1

Background

We first give some definitions, as well as existing proofs whose ideas we will use to derive our results.

1.1 Geometric graphs

Definition 1.1 (geometric graph) *Given a set of points \mathcal{P} in the Euclidean plane \mathbb{R}^2 , a geometric graph is a collection of straight line segments (edges) whose endpoints are elements of \mathcal{P} .*

A geometric graph can be described as a simple graph (in the combinatorial sense) on the vertices \mathcal{P} , where the edge $\{p, q\}$ corresponds to the segment joining p and q .

Definition 1.2 (crossing-free) *A geometric graph is crossing-free if no two edges share points other than their endpoints; it is called crossing otherwise.*

Note that being crossing-free implies that the corresponding simple graph is planar, and that the geometric graph is a plane embedding.

Definition 1.3 (triangulation) *A triangulation is a maximal crossing-free geometric graph, that is, a geometric graph such that for all p and q in \mathcal{P} that are not joined by a segment, adding the segment joining p and q would result in a crossing geometric graph.*

Note that the faces (in the sense of plane graphs) formed by a triangulation are all triangles, with the possible exception of the outer face (thus this definition is *not* equivalent to that of a triangulation of the 2-sphere).

Since a geometric graph corresponds to a simple graph on the underlying point set, we can also look at geometric graphs that belong to special classes of simple graphs. As we will only be focusing on geometric graphs, we

simply call *perfect matching* a geometric graph which corresponds to a perfect matching in the combinatorial sense.

As outlined in the introduction, we are interested in statements regarding the number of geometric graphs, possibly restricted to a certain type, on a given point set. Evidently, that number would depend on the choice of the point set \mathcal{P} , so instead we seek bounds on that number depending on the cardinality $|\mathcal{P}|$, and possibly restricting \mathcal{P} so that it satisfies certain properties.

An important property that \mathcal{P} can have is general position. We say that a point set in the plane is *in general position* if no three points are collinear. In this thesis we will only study point sets in general position.

In general, if $g(\mathcal{P})$ is the number of geometric graphs of a certain sort on the point set \mathcal{P} , we will look for lower bounds l and upper bounds u of the form

$$\forall n \in \mathbb{N}, \forall \mathcal{P} \text{ such that } |\mathcal{P}| = n, l(n) \leq g(\mathcal{P}) \leq u(n),$$

where \mathcal{P} runs over all point sets that satisfy the relevant properties.

1.2 Asymptotics

In addition, we will want asymptotics on such bounds l and u , to compare them with existing results. Since the bounds are often exponential, we tend to ignore polynomial factors. We will thus write that

$$u(n) \preceq_n f(n)$$

if for some polynomial Π , $\forall n \in \mathbb{N}, u(n) \leq \Pi(n)f(n)$.

In some cases, we may be interested in classes of functions $g_z(\mathcal{P})$ which depend on some parameter $z \in \Omega_n$, where Ω_n is a set of arbitrary objects. For instance, g_z could count crossing-free graphs on \mathcal{P} that contain an independent set of size $z \in [n]$. In this thesis, we will often use classes g_z with z being the left-to-right ordering of designated left and right endpoints—this will be formalized in section 1.6.

When giving exponential bounds in n for these parameterized classes of graphs, we will want to keep track of any dependence on z . We will write that

$$u(n, z) \preceq_n f(n, z) \text{ uniformly in } z \in \Omega_n$$

if for some polynomial Π (that does not depend on z),

$$\forall n \in \mathbb{N}, \forall z \in \Omega_n, u(n, z) \leq \Pi(n)f(n, z).$$

If $f(n, z)$ admits a supremum $F(n)$ in $z \in \Omega_n$ for all n , this implies that $u(n, z) \leq \Pi(n)F(n)$, giving an upper bound truly independent of z . It also makes it possible to get a bound by summation of $f(n, z)$ over $z \in \Omega_n$.

Observe that these bounds up to a polynomial are related to \mathcal{O} notation when the upper bound is exponential; specifically, for $\zeta > 1$,

$$u(n) \preceq_n \zeta^n \implies \forall \varepsilon > 0, u(n) \in \mathcal{O}((\zeta + \varepsilon)^n).$$

Moreover, we can bound a product by the product of the bounds for its factors. Specifically, for positive f_1 and f_2 , if $u_1(a, z) \preceq_a f_1(a, z)$ and $u_2(b, z) \preceq_b f_2(b, z)$, both uniformly in z , we have

$$u_1(a, z)u_2(b, z) \preceq_m f_1(a, z)f_2(m - a, z)$$

uniformly in $b \in [0, m]$ and in z . Indeed, if $u_i(a, z) \leq \Pi_i(a)f_i(a, z)$, where without loss of generality the coefficients of Π_i are positive, we get

$$\begin{aligned} u_1(a, z)u_2(b, z) &\leq \Pi_1(a)\Pi_2(b)f_1(a, z)f_2(b, z) \\ &\leq \Pi_1(a + b)\Pi_2(a + b)f_1(a, z)f_2(b, z). \end{aligned}$$

1.3 A remark on the asymptotics of the binomial coefficient

Unsurprisingly, binomial coefficients will be involved in the bounds, so we now give a uniform exponential upper bound for them. It can be shown, by writing the logarithm of the factorial as a sum of logarithms, and then bounding that sum by integrals, that

$$\forall k \in \mathbb{N}^*, e\left(\frac{k}{e}\right)^k \leq k! \leq ke\left(\frac{k}{e}\right)^k,$$

where $\mathbb{N}^* := \mathbb{N} \setminus \{0\}$. It follows that

$$\begin{aligned} \forall k \in \mathbb{N}^*, \forall \eta \in]0, 1[, \binom{k}{\eta k} &\leq \frac{ke\left(\frac{k}{e}\right)^k}{e\left(\frac{\eta k}{e}\right)^{\eta k} e\left(\frac{(1-\eta)k}{e}\right)^{(1-\eta)k}} \\ &= \frac{k}{e} \frac{k^k}{(\eta k)^{\eta k} ((1-\eta)k)^{(1-\eta)k}} \\ &= \frac{k}{e} \left(\frac{1}{\eta^\eta (1-\eta)^{1-\eta}} \right)^k \\ &= \frac{k}{e} \left(\frac{(1-\eta)^{\eta-1}}{\eta^\eta} \right)^k, \end{aligned}$$

so that, since $\forall k \in \mathbb{N}, \binom{k}{0} = \binom{k}{k} = 1$, we have

$$\binom{k}{\eta k} \preceq_k \left(\frac{(1-\eta)^{\eta-1}}{\eta^\eta} \right)^k \text{ uniformly in } \eta \in [0, 1]. \quad (1.1)$$

1.4 Triangulations of convex point sets and the Catalan numbers

A set of points is said to be *in convex position* if it is in general position and all points are in the boundary of its convex hull.

For points in convex position, the numbers of triangulations and perfect matchings are known. The following recurrence, given by J. A. Segner in 1758 [11], defines the *Catalan numbers*.

Theorem 1.4 (Segner) *Let C_n be the number of triangulations of a set of $n + 2$ points in convex position. Then $C_0 = 1$, and for $n > 0$,*

$$C_n = \sum_{i=0}^{n-1} C_i C_{n-i-1}. \quad (1.2)$$

Proof Consider a set of $n + 2$ points in convex position. For $n = 0$ the result is trivial, the only triangulation is an edge.

Let $n > 0$, and number the points along the boundary of the convex hull, from 0 to $n + 1$, see figure 1.1. The edge e joining points $n + 1$ and n is in the boundary of the convex hull, and thus must be in any triangulation, since no other edge can cross it. Let i be the index of the third point of the triangle containing edge e , which may be any of the others, from 0 to $n - 1$.

Then, restricting the triangulation to the $i + 2$ points $\{0, \dots, i, n + 1\}$ yields a triangulation of those points. Similarly, restricting the triangulation of the whole point set to points i through n yields a triangulation of $n - i + 1$ points. It follows that $C_n \leq \sum_{i=0}^{n-1} C_i C_{n-i-1}$.

Moreover, given a triangulation of $\{0, \dots, i, n + 1\}$ and a triangulation of $\{i, \dots, n\}$, adding the edge e forms a triangulation of the whole point set, proving equality. \square

An explicit expression for C_n was given earlier by Euler [7], along with the generating function. A convenient form is

$$C_n = \frac{1}{n+1} \binom{2n}{n}, \quad (1.3)$$

yielding asymptotically

$$C_n \sim \frac{4^n}{n^{\frac{3}{2}} \sqrt{\pi}} \preceq_n 4^n. \quad (1.4)$$




Figure 1.1 – Segner's recurrence.




Figure 1.2 – Partitioning a convex point set to construct a crossing-free perfect matching.

1.5 Crossing-free perfect matchings

We will call $\text{PM}_{\mathcal{P}}$ the set of perfect matchings on the point set \mathcal{P} , and $\text{CFPM}_{\mathcal{P}}$ the set of crossing-free perfect matchings on the point set \mathcal{P} . When discussing perfect matchings, we will denote the unique edge incident to point p by e_p .

If \mathcal{P} is a set of $n = 2k$ points in convex position, it was shown¹ by A. Errera in 1930 [6] that

$$|\text{CFPM}_{\mathcal{P}}| = C_k.$$

Indeed, given a crossing-free perfect matching, number the points from 0 to $2k - 1$ along the boundary of the convex hull, as in figure 1.2. Let j be the point to which 0 is matched by edge e_0 . Observe that e_0 partitions the point set. It follows that the restriction of the matching to either $\{j + 1, \dots, 2k - 1\}$ or $\{1, \dots, j - 1\}$ is a crossing-free perfect matching. As a consequence, these sets have even size, so $j = 2i + 1$ for some i . We thus have Segner's recurrence for the number of crossing-free perfect matchings on a convex point

¹In fact, Errera counts the number of figures that can be formed in the upper half-plane by k non-intersecting semicircles with their centres on the boundary. Since these are geodesics of the hyperbolic plane in the Poincaré half-plane model, switching to the Beltrami-Klein model turns the half-plane into the unit disk, and the semicircles into chords, so that the problem becomes that of crossing-free perfect matchings in convex position.

set. Since there is only one—fairly boring—crossing-free perfect matching on the empty set, we get the Catalan numbers.

For a set \mathcal{P} of n points in general position, it was proved by Micha Sharir and Emo Welzl in 2005 [13] that $|\text{CFPM}_{\mathcal{P}}| \preceq_n 10.05^n$.

1.6 Bracket expressions and their relations to perfect matchings

We say that a point set is *in general position with respect to the horizontal* if it is in general position and no two points lie on a vertical line. Note that any point set in general position can be put in general position with respect to the horizontal by an arbitrarily small rotation. Moreover, note that points in general position with respect to the horizontal are ordered from left to right. For a point set \mathcal{P} in general position with respect to the horizontal, we call its leftmost point $\text{lm } \mathcal{P}$.

This ordering can be used to classify perfect matchings on the point set; to this end, we introduce bracket expressions. A *bracket expression* of length n is a sequence of n opening brackets \langle or closing brackets \rangle . It is a *well-formed prefix* if, when read from left to right, the number of closing brackets encountered never exceeds the number of opening brackets encountered. A *well-formed bracket expression* is a well-formed prefix with the same number of opening and closing brackets.

It is a well-known result, shown² by E. C. Catalan in 1838 [3], that the number of well-formed bracket expressions of length $2k$ is the Catalan number C_k . In fact, the recurrence can readily be seen from a grammatical definition of well-formed bracket expressions,³

$$\begin{aligned} \text{wfbe} ::= & \langle \text{wfbe} \rangle \text{wfbe} \\ & | \text{empty}, \end{aligned}$$

where the sum in the recurrence ranges over the lengths of the component bracket expressions.

Note that inserting an opening bracket, followed by a closing bracket anywhere after the inserted opening bracket, into a well-formed bracket expression, results in a well-formed bracket expression.

Well-formed bracket expressions can be used to define classes of perfect matchings on a given point set. Given a point set \mathcal{P} of size n in general

²Specifically, Catalan showed that the recurrence (1.2) counts the number of parenthesizings of $n + 1$ factors.

³We have not enclosed nonterminals in angle brackets here, as would be usual in Backus-Naur form, since that would result in hopeless confusion in a grammar describing nothing but sequences of angle brackets.




Figure 1.3 – A well-formed bracket expression constructed from a perfect matching.

position with respect to the horizontal and a perfect matching μ on \mathcal{P} , any point in \mathcal{P} is either a left or right endpoint of an edge, since it is incident to exactly one edge, and that edge is not vertical.

Construct a bracket expression $\mathfrak{B}_{\mathcal{P}}(\mu)$ of length n as follows: order \mathcal{P} from left to right; the i th bracket is opening if the i th point of \mathcal{P} is a left endpoint of μ , and it is closing otherwise. This bracket expression is well-formed, since it can be constructed by starting from the (well-formed) empty bracket expression, by successively inserting both brackets corresponding to each edge, where the closing bracket will be inserted to the right of the opening bracket.

Given a well-formed bracket expression B and a point set \mathcal{P} , we will say that a perfect matching μ is *consistent with B* if $\mathfrak{B}_{\mathcal{P}}(\mu) = B$; moreover, we will refer to the points of \mathcal{P} corresponding to opening brackets of B as *left-points* (since they will be left endpoints of any perfect matching consistent with B), and similarly we will refer to the points of \mathcal{P} corresponding to closing brackets of B as *right-points*.

One approach to bounding the size of $\text{CFPM}_{\mathcal{P}}$ is to bound the size of $\mathfrak{B}_{\mathcal{P}}^{-1}(B) \cap \text{CFPM}_{\mathcal{P}}$ for bracket expressions B of length n , *i.e.*, the number of crossing-free perfect matchings on \mathcal{P} consistent with B . We thus define

$$\nu_{\mathcal{P}}(B) := \left| \mathfrak{B}_{\mathcal{P}}^{-1}(B) \cap \text{CFPM}_{\mathcal{P}} \right|.$$

In order to concisely refer to bracket expressions, we will use the notations \langle^k for k successive opening brackets, and \rangle^k for k successive closing brackets, for instance,

$$\langle^2 \rangle \langle^2 \rangle^3 = \langle \langle \rangle \langle \rangle \rangle \rangle.$$

A remarkable property of ν is its behaviour with respect to concatenation of well-formed bracket expressions. If B_1 and B_2 are well-formed bracket expressions of lengths k_1 and $k_2 = n - k_1$, \mathcal{P}_1 is the set of the leftmost k_1 points of \mathcal{P} , and $\mathcal{P}_2 = \mathcal{P} \setminus \mathcal{P}_1$ is the set of the rightmost k_2 points, we have

$$\nu_{\mathcal{P}}(B_1 B_2) = \nu_{\mathcal{P}_1}(B_1) \nu_{\mathcal{P}_2}(B_2).$$




Figure 1.4 – Untangling in a perfect matching. Replacing the thick edges by the thin ones reduces the total edge length, while preserving left and right endpoints.

Indeed, no edge may join \mathcal{P}_1 and \mathcal{P}_2 , since half the points of \mathcal{P}_1 are right-points, and \mathcal{P}_1 lies to the left of \mathcal{P}_2 . The problem thus decomposes in two independent parts.

The function $\nu_{\mathcal{P}}$ has been studied before; though they did not explicitly phrase it in terms of brackets, M. Sharir and E. Welzl showed in 2005 [13] that

$$\nu_{\mathcal{P}}(B) \preceq_n 5.3793^n \quad (1.5)$$

uniformly in B . Taking the sum of this bound over bracket expressions B yields an upper bound for the overall number of crossing-free perfect matchings,

$$|\text{CFPM}_{\mathcal{P}}| \preceq_n C_{\frac{n}{2}} 5.3793^n \preceq_n 10.76^n,$$

but it is worse than the one mentioned in the previous section.

1.7 An optimal lower bound for the number of crossing-free perfect matchings

The following result, and its proof are, due to A. J. Ruiz-Vargas and E. Welzl [10].

Theorem 1.5 (Ruiz-Vargas–Welzl) *Let \mathcal{P} be a point set of size $n = 2k$ in general position with respect to the horizontal, and let B be a well-formed bracket expression of length n . Then there exists a crossing-free perfect matching consistent with B ; in other words, $\nu_{\mathcal{P}}(B) \geq 1$.*

Proof Let μ_0 be a perfect matching on \mathcal{P} consistent with B . This is always possible, for instance, parsing the bracket expression, match the point corresponding to an opening bracket and the point corresponding to the matching closing bracket.

Define $\Lambda(\mu)$ for a perfect matching μ on \mathcal{P} to be the sum of the lengths of the edges of μ .

Then, repeat the following procedure, starting at $i = 0$. If there is no crossing in μ_i , we have found a perfect matching with the desired properties. If there is a crossing, let p, p', q , and q' be the points involved, with p and p' being the left endpoints and q and q' being the right endpoints, so that the edge pq crosses the edge $p'q'$. Remove these edges, and replace them by pq' and $p'q$, thus untangling them. This yields another perfect matching μ_{i+1} . By the triangle inequality (see figure 1.4), $\Lambda(\mu_{i+1}) < \Lambda(\mu_i)$.

If this procedure did not terminate, it would yield a sequence $(\mu_i)_{i \in \mathbb{N}}$ of crossing perfect matchings on \mathcal{P} on which Λ is strictly decreasing, thus an infinite sequence of distinct graphs on \mathcal{P} . Since there are only finitely many graphs on \mathcal{P} , this is a contradiction, so we eventually⁴ find a crossing-free perfect matching. \square

This immediately yields a lower bound for the number of crossing-free perfect matchings, since there are $C_{\frac{n}{2}}$ well-formed bracket expressions of length n . Note that this lower bound had already been proved by A. García, M. Noy, and J. Tejel in 1995, by a recurrence that split the point set, yielding the Segner recurrence with inequality, rather than via well-formed bracket expressions [8].

Corollary 1.6 *Let \mathcal{P} be a point set of even size n in general position. There are at least $C_{\frac{n}{2}}$ distinct crossing-free perfect matchings on \mathcal{P} , i.e., $|\text{CFPM}_{\mathcal{P}}| \geq C_{\frac{n}{2}}$.*

Moreover, this lower bound is optimal, since it is attained if \mathcal{P} is in convex position (see section 1.5).

Now that we have a tight uniform lower bound for $\nu_{\mathcal{P}}$, we will start looking at upper bounds dependent on the bracket expression.

1.8 Matchings across a line

Again we consider $n = 2k$ points in general position with respect to the horizontal. The matchings consistent with bracket expressions with k opening brackets followed by k closing brackets, $\langle^k \rangle^k$, are called *matchings across a line*. Indeed, any edge in such a matching will cross any vertical line that separates the left-points from the right-points.

The following result, and its proof, were given by M. Sharir and E. Welzl in 2005 [13].

Theorem 1.7 (Sharir–Welzl) *Let \mathcal{P} be a set of $n = 2k$ points in general position with respect to the horizontal. Then there are at most $C_{\frac{n}{2}}^2$ crossing-free perfect matchings across a line on \mathcal{P} , i.e., $\nu_{\mathcal{P}}(\langle^k \rangle^k) \leq C_k^2$.*

⁴The untangling procedure itself predates this proof, and was shown to terminate in $\mathcal{O}(n^3)$ steps by J. van Leeuwen and A. A. Schoone in 1980 [16].




Figure 1.5 – Constructing the numberings λ and ρ of the left- and right-points from a crossing-free perfect matching across a line; here $\iota(\{p, q\}) = 2$, and thus $\lambda(p) = \rho(q) = 2$.

The idea of the proof is as follows. First, pick a vertical line that separates the left-points from the right-points; we will call it *the vertical line*. Further, let us call the set of left-points \mathcal{L} and the set of right-points \mathcal{R} .

A perfect matching consistent with a given well-formed bracket expression (and in particular, a perfect matching across a line) is uniquely defined by a bijection $\mu : \mathcal{L} \rightarrow \mathcal{R}$ from the left-points to the right-points. Consider such a perfect matching across a line, and let E be the set of its edges. Now, number the intersections between the edges of the perfect matching and the vertical line from top to bottom. This yields a numbering $\iota : E \rightarrow [k]$ of the edges.

Define $\lambda(p) := \iota(e_p)$ mapping a left-point to the intersection number of its edge, and similarly $\rho(q) := \iota(e_q)$ for the right-points. We have $\mu = \rho^{-1} \circ \lambda$. The bijection λ (respectively ρ) determines the order in which the left points (respectively right points) reach the vertical line.

If the matching is crossing-free, we will show that λ and ρ have to be in sets of size C_k , thus that there are at most C_k^2 functions $\mu = \rho^{-1} \circ \lambda$, i.e., that there can be at most C_k^2 crossing-free perfect matchings across a line.

Since we are going to reuse these concepts in subsequent chapters, we will formalize and name the properties of λ and ρ that we will consider.

Definition 1.8 (crossing-free alignment) *Let S be a set of k points in general position. Let V be a vertical line to the right of S . A bijection λ from S to $[k]$ is a crossing-free left-alignment of S on V if there exists a set $A = \{\sigma_p \mid p \in S\}$ of k line segments such that the following hold:*

1. *for every $p \in S$, the left endpoint of σ_p is p , and its right endpoint is on V ;*
2. *no two segments cross, and no two segments share any endpoints;*




Figure 1.6 – A crossing-free left-alignment of five points. Once the index of the leftmost point is chosen (thick segment), the rest consists of two crossing-free left-alignments of i and $k - i - 1$ points each—here $k = 5$ and $i = 2$.

3. let $\iota : A \rightarrow [k]$ the numbering of the right endpoints of the segments in A from top to bottom, then $\lambda(p) = \iota(\sigma_p)$.

Such a set of segments is said to realize λ .

Correspondingly, we define a crossing-free right-alignment, where “right” and “left” are swapped in the definition.

A crossing-free left-alignment of five points, with a realizing set of segments—drawn as arrows—is shown in figure 1.6.

Lemma 1.9 Let \mathcal{S} be a set of k points in general position, and let V be a vertical line to the right of \mathcal{S} . There are at most C_k crossing-free left-alignments of \mathcal{S} on V .

Proof Let λ be a crossing-free left-alignment of \mathcal{S} on V . Let $A = \{\sigma_p \mid p \in \mathcal{S}\}$ be a set of non-crossing line segments realizing λ . For a point $p \in \mathcal{S}$, we will call σ_p the segment of p .

$\lambda(\text{lm } \mathcal{S})$, being the index on the vertical line V of the right endpoint of the segment $\sigma_{\text{lm } \mathcal{S}}$ of the leftmost point, is equal to 1 plus the number of points of \mathcal{S} that are above $\sigma_{\text{lm } \mathcal{S}}$. Indeed, the segments of points above $\sigma_{\text{lm } \mathcal{S}}$ must themselves reach the vertical line V above $\sigma_{\text{lm } \mathcal{S}}$, otherwise they would cross $\sigma_{\text{lm } \mathcal{S}}$, and correspondingly for points below $\sigma_{\text{lm } \mathcal{S}}$, so that there are as many edges reaching the vertical line V below $\sigma_{\text{lm } \mathcal{S}}$ as there are points below $\sigma_{\text{lm } \mathcal{S}}$.

Moreover, as the oriented angle between $\sigma_{\text{lm } \mathcal{S}}$ and the horizontal increases, points are only added to the set of points below $\sigma_{\text{lm } \mathcal{S}}$, so that choosing the number of points below $\sigma_{\text{lm } \mathcal{S}}$ determines the sets of points below and above $\sigma_{\text{lm } \mathcal{S}}$.

Further, since points above $\sigma_{\text{lm } \mathcal{S}}$ must reach the vertical line V above $\sigma_{\text{lm } \mathcal{S}}$, and correspondingly for points below, if the point p is above $\sigma_{\text{lm } \mathcal{S}}$, then $\lambda(p) < \lambda(\text{lm } \mathcal{S})$, and if it is below, $\lambda(p) > \lambda(\text{lm } \mathcal{S})$. Thus λ restricted to the points above $\sigma_{\text{lm } \mathcal{S}}$ is a crossing-free left-alignment of the $\lambda(\text{lm } \mathcal{S}) - 1$




Figure 1.7 – Composing crossing-free alignments, only to get a crossing perfect matching.




Figure 1.8 – Composing crossing-free alignments, and getting a crossing-free perfect matching, but one that does not decompose to the original alignments.

points above $\sigma_{\text{lm } \mathcal{S}}$, and $\lambda - \lambda(\text{lm } \mathcal{S})$ restricted to the points below $\sigma_{\text{lm } \mathcal{S}}$ is a crossing-free left-alignment of the $k - \lambda(\text{lm } \mathcal{S})$ points below $\sigma_{\text{lm } \mathcal{S}}$, where the appropriate subsets of A provide the set of segments required in the definition.

It follows that λ is determined by the choice of $\lambda(\text{lm } \mathcal{S})$ and crossing-free left-alignments of $\lambda(\text{lm } \mathcal{S}) - 1$ and $k - \lambda(\text{lm } \mathcal{S})$ points. Letting $i := \lambda(\text{lm } \mathcal{S}) - 1$, these are crossing-free alignments of i and $k - i - 1$ points (see figure 1.6). Therefore, if ω_j is an upper bound for the number of crossing-free left-alignments of j points when $j < k$, we can give an upper bound on the number of crossing-free left-alignments of k points,

$$\omega_k := \sum_{i=0}^{k-1} \omega_i \omega_{k-i-1}.$$

We can start the recurrence with $\omega_0 = 1$; this is the recurrence for the Catalan numbers, thus $\omega_k = C_k$. \square

The same result holds for crossing-free right-alignments, simply exchange “left” and “right” in the above lemma and its proof.

Proof (of theorem 1.7) If we have a crossing-free perfect matching across a line, then λ as constructed in figure 1.5 is a crossing-free left-alignment of \mathcal{L} on the vertical line; indeed, letting σ_p be the portion of the edge e_p to the left of the vertical line yields segments that realize λ . Similarly, ρ is a crossing-free right-alignment of \mathcal{R} on the vertical line. \square

Chapter 2

Analysing the overcounting in the upper bound for matchings across a line

The bound given in the previous chapter for the number of matchings across a line is not optimal; an example is given by Sharir and Welzl of a crossing-free left-alignment λ and a crossing-free right-alignment ρ such that $\rho^{-1} \circ \lambda$ is not crossing-free (see figure 1.7).

There is however more to the overcounting than just counting some crossing matchings. To characterize that, it helps to name some functions and sets. Again, we are considering a set \mathcal{P} of $2k$ points in general position with respect to the horizontal and the bracket expression $\langle^k \rangle^k$. We let \mathcal{L} be the set of left-points, and \mathcal{R} be the set of right-points. Moreover, we pick a vertical line V separating \mathcal{L} and \mathcal{R} . In this chapter and the following one, we will use the terms crossing-free left- and right-alignments to mean crossing-free left- and right-alignments on V .

In the previous chapter, given a crossing-free perfect matching μ , we constructed crossing-free left- and right-alignments λ and ρ (see figure 1.5). Let us call these crossing-free alignments the *decomposition* \mathfrak{D} of μ , thus $\mathfrak{D}(\mu) = (\lambda, \rho)$. We remarked that the composition $\rho^{-1} \circ \lambda$ of these alignments yielded the original matching. We thus define the *composition* $\mathfrak{C}(\lambda, \rho) := \rho^{-1} \circ \lambda$ for arbitrary crossing-free left-alignment λ of \mathcal{L} on V and ρ of \mathcal{R} on V . Moreover, let us call $\text{CFLA}_{\mathcal{L}, V}$ the set of crossing-free left-alignments of \mathcal{L} on V , and $\text{CFLA}_{\mathcal{R}, V}$ the set of crossing-free left-alignments of \mathcal{R} on V . Finally, let us name the sets of perfect matchings across a line and of crossing-free perfect matchings across a line,

$$\begin{aligned} \text{PMAL}_{\mathcal{P}} &:= \mathfrak{B}_{\mathcal{P}}^{-1}(\langle^k \rangle^k) \\ \text{CFPMAL}_{\mathcal{P}} &:= \text{PMAL}_{\mathcal{P}} \cap \text{CFPM}_{\mathcal{P}}. \end{aligned}$$

2. ANALYSING THE OVERCOUNTING IN THE UPPER BOUND FOR MATCHINGS ACROSS A LINE

Then the following diagram commutes, *i.e.*, decomposing a crossing-free perfect matching across a line and then composing the resulting alignments yields the initial matching.

$$\begin{array}{ccc}
 \text{CFPMAL}_{\mathcal{P}} & \xrightarrow{\mathfrak{D}} & \text{CFLA}_{\mathcal{L},V} \times \text{CFRA}_{\mathcal{R},V} \\
 \downarrow \subseteq & \swarrow \mathfrak{C} & \\
 \text{PMAL}_{\mathcal{P}} & &
 \end{array}$$

We will call the elements of $\text{CFLA}_{\mathcal{L},V} \times \text{CFRA}_{\mathcal{R},V}$ *pairs of alignments*.

The result of Sharir and Welzl is proved in two steps. First we see from the commutative diagram that $\text{CFPMAL}_{\mathcal{P}}$ lies within the image of \mathfrak{C} , so that

$$\begin{aligned}
 \nu_{\mathcal{P}}(\langle^k \rangle^k) &= |\text{CFPMAL}_{\mathcal{P}}| \\
 &\leq |\mathfrak{C}(\text{CFLA}_{\mathcal{L},V} \times \text{CFRA}_{\mathcal{R},V})| \\
 &\leq |\text{CFLA}_{\mathcal{L},V} \times \text{CFRA}_{\mathcal{R},V}|.
 \end{aligned} \tag{2.1}$$

Then, we derive an upper bound for the number $|\text{CFLA}_{\mathcal{L},V} \times \text{CFRA}_{\mathcal{R},V}|$ of pairs of alignments. We will examine this second step later. Looking at the commutative diagram, we can see that \mathfrak{D} is injective; this is another way of getting (2.1), but more importantly, it means that $\nu_{\mathcal{P}}(\langle^k \rangle^k)$ is equal to the cardinality of the image of \mathfrak{D} .

It follows that the pairs of alignments that are *not* in the image of \mathfrak{D} are responsible for the slackness of the inequality (2.1). We will thus call them *overcounted pairs of alignments*. An overcounted pair of alignments (λ, ρ) falls into one of two families:

1. $\mathfrak{C}(\lambda, \rho)$ can be a crossing perfect matching, as in figure 1.7;
2. $\mathfrak{C}(\lambda, \rho)$ can be a crossing-free perfect matching; in this case, we have $(\lambda, \rho) \neq \mathfrak{D} \circ \mathfrak{C}(\lambda, \rho)$, since (λ, ρ) is overcounted; an example is shown in figure 1.8.

We would like to give a bound on the number of non-overcounted pairs of alignments. In order to do that, we turn to the second part of the proof of Sharir and Welzl, namely the bound on the number of pairs of alignments, and examine it in more detail, with some fresh definitions. Again, we will look at crossing-free left-alignments; the following argument equally applies to crossing-free right-alignments.

For a point set \mathcal{S} in general position with respect to the horizontal, note that we can partition the half-plane to the right of $\text{Im } \mathcal{S}$ into $|\mathcal{S}|$ regions, separated by rays emanating from $\text{Im } \mathcal{S}$ and going through other points of \mathcal{S} , see figure 2.1. We will call those regions, from top to bottom, $\mathcal{R}_{\mathcal{S},1}$ to $\mathcal{R}_{\mathcal{S},|\mathcal{S}|}$. Note that there are i points of \mathcal{S} above the interior of $\mathcal{R}_{\mathcal{S},i+1}$, and




Figure 2.1 – A set \mathcal{S} of five points. The half-plane to the right of the leftmost point $\text{lm } \mathcal{S}$ is partitioned into $|\mathcal{S}|$ cones $k_{\mathcal{S},i}$.




Figure 2.2 – A crossing-free perfect matching μ across a line V . Its decomposition is $(\lambda, \rho) := \mathcal{D}(\mu)$; the crossing-free left-alignment λ is shown here. Since $\lambda(\text{lm } \mathcal{L}) = 3$, $\text{lm } \mathcal{L}$ is matched within the cone $k_{\mathcal{L},3}$ (shaded).

$|\mathcal{S}| - i - 1$ points below. We will use the term *left cone* of \mathcal{L} to mean $k_{\mathcal{S},i}$ for some $\mathcal{S} \subseteq \mathcal{L}$ and some $i \in [\mathcal{S}]$.

Observe that for a crossing-free left-alignment λ of \mathcal{L} with $\lambda(\text{lm } \mathcal{L}) = i + 1$, any realizing segment of λ incident to $\text{lm } \mathcal{L}$ separates the points $\mathcal{L} \setminus \{\text{lm } \mathcal{L}\}$ into sets of points above and below the segment, as in figure 1.6. Furthermore, these sets are the same as the sets of points above and below $k_{\mathcal{L},i+1}$. As a consequence, given a crossing-free left-alignment λ of \mathcal{L} , for any set of segments $A = \{\sigma_p \mid p \in \mathcal{L}\}$ realizing λ , the segment $\sigma_{\text{lm } \mathcal{L}}$ of the leftmost point has to be contained in $k_{\mathcal{L},\lambda(\text{lm } \mathcal{L})}$. As we remarked in section ??, for a crossing-free perfect matching μ , the edges of μ restricted to the left side of V realize the crossing-free left-alignment in $\mathcal{D}(\mu)$. Thus, if $(\lambda, \rho) = \mathcal{D}(\mu)$ is in the image of \mathcal{D} , μ must match $\text{lm } \mathcal{L}$ to a right-point in the cone $k_{\mathcal{L},\lambda(\text{lm } \mathcal{L})}$, see figure 2.2.

To extend this reasoning beyond the leftmost point of \mathcal{L} , we define *left conifers*¹ on \mathcal{L} . A left conifer κ on \mathcal{L} is a map from points in \mathcal{L} to left cones

¹Conifers are constructed by recursive subdivision, so in some sense they are trees with

of \mathcal{L} , satisfying the following properties:

1. $\kappa(\text{lm } \mathcal{L})$ is a left \mathcal{L} -cone;
2. κ restricted to the points above the interior of $\kappa(\text{lm } \mathcal{L})$ is a left conifer on these points;
3. κ restricted to the points below the interior of $\kappa(\text{lm } \mathcal{L})$ is a left conifer on these points.

See figure 2.3 for an example.




Figure 2.3 – The construction of a left conifer on a set \mathcal{L} of five points.

Let $\text{LC}_{\mathcal{L}}$ be the set of left conifers on \mathcal{L} . From the observations above on the number of cones and the size of the sets of points above and below their interiors, we can see that the number of left conifers on a point set depends only on the size of that point set. Specifically, if ω_k is the number of left conifers on k points, we have

$$\omega_k = \sum_{i=0}^{k-1} \omega_i \omega_{k-i-1} \text{ and } \omega_0 = 1,$$

where the summation index i corresponds to the choice $\kappa(\text{lm } \mathcal{L}) = \mathcal{K}_{\mathcal{L},i+1}$. We thus have $|\text{LC}_{\mathcal{L}}| = C_{|\mathcal{L}|}$.

Recall that a crossing-free left-alignment λ of \mathcal{L} on V is determined by the value of $\lambda(\text{lm } \mathcal{L})$ and by crossing-free left alignments λ^- and λ^+ on V of cones, hence the name.



Figure 2.4 – A crossing-free left-alignment λ (the one from figure 1.6), together with the conifer \mathfrak{R}_λ .

subsets \mathcal{S}^- and \mathcal{S}^+ above and below any edge realizing λ incident to $\text{lm } \mathcal{L}$, *i.e.*, subsets above and below $\mathcal{K}_{\mathcal{L}, \lambda(\text{lm } \mathcal{L})}$. In other words, we have

$$\begin{aligned}\mathcal{S}^- &= \lambda^{-1}(\{1, \dots, \lambda(\text{lm } \mathcal{L}) - 1\}), \\ \mathcal{S}^+ &= \lambda^{-1}(\{\lambda(\text{lm } \mathcal{L}) + 1, \dots, |\mathcal{L}|\}), \\ \lambda^-(p) &= \lambda(p) \text{ for } p \in \mathcal{S}^-, \text{ and} \\ \lambda^+(p) &= \lambda(p) - \lambda(\text{lm } \mathcal{L}) \text{ for } p \in \mathcal{S}^+.\end{aligned}$$

Given a crossing-free left-alignment λ of \mathcal{L} on V , this allows us to construct a left conifer \mathfrak{R}_λ as follows:

1. $\mathfrak{R}_\lambda(\text{lm } \mathcal{L}) = \mathcal{K}_{\mathcal{L}, \lambda(\text{lm } \mathcal{L})}$;
2. $\mathfrak{R}_\lambda(p) = \mathfrak{R}_{\lambda^-}(p)$ for $p \in \mathcal{S}^-$;
3. $\mathfrak{R}_\lambda(p) = \mathfrak{R}_{\lambda^+}(p)$ for $p \in \mathcal{S}^+$.

See figure 2.4 for an example.

This gives an injective map

$$\begin{aligned}\mathfrak{R} : \text{CFLA}_{\mathcal{L}, V} &\longrightarrow \text{LC}_{\mathcal{L}} \\ \lambda &\longmapsto \mathfrak{R}_\lambda\end{aligned}$$

from crossing-free left-alignments of \mathcal{L} on V to left conifers on \mathcal{L} . In turn, this yields $|\text{CFLA}_{\mathcal{L}, V}| \leq |\text{LC}_{\mathcal{L}}| = \text{C}_{|\mathcal{L}|}$, which is exactly the Sharir–Welzl bound. Note that \mathfrak{R} is in general not bijective, see figure 2.5.

Now, note that the map \mathfrak{R} has the property that for any set of segments $A = \{\sigma_p \mid p \in \mathcal{L}\}$ realizing λ , σ_p lies inside the cone $\mathfrak{R}_\lambda(p)$. Since the edges of a crossing-free perfect matching μ realize the crossing-free alignments $\mathfrak{D}(\mu) = (\lambda, \rho)$, $\mathfrak{R}_\lambda(p)$ must contain the right-point to which μ matches p . This extends the property illustrated by figure 2.2 to the whole of \mathcal{L} .




Figure 2.5 – It is impossible to construct a vertical line V to the right of these four points and a crossing-free left-alignment λ of these points on V such that \mathfrak{K}_λ is the conifer shown. Indeed, realizing segments σ_p and σ_q would have to lie within the red and green cones respectively, and thus they would cross, a contradiction.

This property allows us to derive a necessary condition that a pair of alignments has to fulfill in order not to be overcounted. Consider a crossing-free left-alignment λ such that, for some $p \in \mathcal{L}$, there is no right-point in $\mathfrak{K}_\lambda(p)$. Then, there is no crossing-free right-alignment ρ such that (λ, ρ) is in the image of \mathfrak{D} , i.e., for all crossing-free right-alignments ρ , (λ, ρ) is overcounted.

This condition motivates some definitions. We will call a crossing-free left-alignment λ *overcounted* if for any crossing-free right-alignment ρ , (λ, ρ) is overcounted. We will call a left cone of \mathcal{L} *vacuous* if it does not contain a point from \mathcal{R} . Moreover, we will write that a conifer κ is *vacuous* if for some $p \in \mathcal{L}$, $\kappa(p)$ is vacuous. Thus, if \mathfrak{K}_λ is vacuous, λ is overcounted. It follows that an upper bound on the number of non-vacuous left conifers is an upper bound on the number of non-overcounted crossing-free left-alignments.

Chapter 3

Highly convex matchings across a line

Again, we are considering crossing-free perfect matchings across a line, with a set \mathcal{L} of left-points and a set \mathcal{R} of right-points, with $|\mathcal{L}| = |\mathcal{R}| = k = \frac{n}{2}$, separated by a vertical line V . We will now examine the properties of the convex hull to give an upper bound on the number of non-vacuous left conifers; as observed at the end of the previous chapter, this will yield an upper bound on the number of non-overcounted crossing-free left-alignments, and by applying the same argument on the right, we will derive an upper bound on the number of non-overcounted pairs of alignment—thus an upper bound on the number of crossing-free perfect matchings across a line.

Let us first define some concepts related to convex hulls. For a point set \mathcal{T} in general position, the *extreme points* of \mathcal{T} are those that are in the boundary of the convex hull of \mathcal{T} . For $\mathcal{S} \subseteq \mathcal{L}$, the *liminary points* of \mathcal{S} are the points




Figure 3.1 – Point sets \mathcal{L} and \mathcal{R} , both of size 8. \mathcal{L} has five liminary points, shown in white. One of them is the leftmost point, the others are separated in two upper liminary points and three lower liminary points. The first two \mathcal{L} -cones and the last three \mathcal{L} -cones (shaded) are vacuous.

3. HIGHLY CONVEX MATCHINGS ACROSS A LINE

of \mathcal{S} which are extreme points of $\mathcal{S} \cup \mathcal{R}$, see figure 3.1 which illustrates the special case $\mathcal{S} = \mathcal{L}$. Note that the liminary points of \mathcal{L} are extreme points of \mathcal{P} , and that for any $\mathcal{S} \subseteq \mathcal{L}$, $\text{lm } \mathcal{S}$ is a liminary point of \mathcal{S} .

Let $\mathcal{S} \subseteq \mathcal{L}$. Observe that, clockwise around $\text{lm } \mathcal{S}$, one subset of the liminary points of \mathcal{S} appears first, followed by all points of \mathcal{R} , followed by the rest of the liminary points of \mathcal{S} (except $\text{lm } \mathcal{S}$). We will call those liminary points of \mathcal{S} which appear before \mathcal{R} the *upper liminary points of \mathcal{S}* , and those that appear after \mathcal{R} the *lower liminary points of \mathcal{S}* . Again, figure 3.1 illustrates the special case $\mathcal{S} = \mathcal{L}$.

If there are Δ_1 upper liminary points of \mathcal{S} , then the cones $\mathcal{K}_{\mathcal{S},1}$ through $\mathcal{K}_{\mathcal{S},\Delta_1}$ are vacuous. Similarly, if there are Δ_2 lower liminary points of \mathcal{S} , then the cones $\mathcal{K}_{\mathcal{S},k-\Delta_2}$ through $\mathcal{K}_{\mathcal{S},k}$ are vacuous. This is also shown in figure 3.1 in the case $\mathcal{S} = \mathcal{L}$.

Moreover, for a non-vacuous cone $\mathcal{K}_{\mathcal{S},i+1}$, consider the points \mathcal{S}^- above $\mathcal{K}_{\mathcal{S},i+1}$ and the points \mathcal{S}^+ below $\mathcal{K}_{\mathcal{S},i+1}$. Observe that any upper liminary points of \mathcal{S} are liminary points of \mathcal{S}^- , and any lower liminary points of \mathcal{S} are liminary points of \mathcal{S}^+ .

We can now define an upper bound $\omega_{k,\Delta}$ on the number of non-vacuous left conifers on k left-points of which at least Δ are liminary. We take the maximum over the (nonnegative) values of Δ_1 and Δ_2 , with $\Delta_1 + \Delta_2 = \Delta - 1$ (since the leftmost point is liminary, but neither upper nor lower liminary). Given the numbers of upper and lower liminary points, we take the sum over the choices of $\kappa(\text{lm } \mathcal{L}) = \mathcal{K}_{\mathcal{L},i+1}$, and excluding values of i which correspond to vacuous cones by the observations above. This yields the recurrence

$$\omega_{k,\Delta} := \max_{\Delta_1 + \Delta_2 = \Delta - 1} \sum_{i=\Delta_1}^{k-\Delta_2-1} \omega_{i,\Delta_1} \omega_{k-i-1,\Delta_2}.$$

Moreover, we can start that recurrence with

$$\begin{aligned} \omega_{k,0} &:= \omega_{k,1} \text{ for } k > 0, \\ \omega_{k,k} &:= 1, \end{aligned}$$

since there is at least one liminary point of \mathcal{L} if it is nonempty (the leftmost one), and if all points are liminary, only one cone is non-vacuous.

This is a bound on the number of non-overcounted crossing-free left-alignments of \mathcal{L} . Since the same argument applies on the right, we get the following upper bound on $\nu_{\mathcal{P}}(\langle^k \rangle^k)$ when \mathcal{P} has Γ extreme points.

$$\nu_{\mathcal{P}}(\langle^k \rangle^k) \leq \max_{\Delta_{\mathcal{L}} + \Delta_{\mathcal{R}} = \Gamma} \omega_{k,\Delta_{\mathcal{L}}} \omega_{k,\Delta_{\mathcal{R}}}$$

We will show in section 4.3.2 that

$$\omega_{k,\Delta} = \frac{\Delta+1}{k+1} \binom{2k-\Delta}{k}.$$

We now examine the expression $\omega_{k,\Delta_{\mathcal{L}}} \omega_{k,\Delta_{\mathcal{R}}}$ to maximize it under $\Delta_{\mathcal{L}} + \Delta_{\mathcal{R}} = \Gamma$.

$$\begin{aligned} \omega_{k,\Delta_{\mathcal{L}}} \omega_{k,\Delta_{\mathcal{R}}} &= \frac{\Delta_{\mathcal{L}}+1}{k+1} \binom{2k-\Delta_{\mathcal{L}}}{k} \frac{\Delta_{\mathcal{R}}+1}{k+1} \binom{2k-\Delta_{\mathcal{R}}}{k} \\ &= \frac{\Delta_{\mathcal{L}}+1}{k+1} \frac{(2k-\Delta_{\mathcal{L}})!}{k!(k-\Delta_{\mathcal{L}})!} \frac{\Delta_{\mathcal{R}}+1}{k+1} \frac{(2k-\Delta_{\mathcal{R}})!}{k!(k-\Delta_{\mathcal{R}})!} \end{aligned}$$

Incrementing $\Delta_{\mathcal{L}}$ and decrementing $\Delta_{\mathcal{R}}$ multiplies the above expression by

$$\begin{aligned} \psi &:= \frac{\Delta_{\mathcal{L}}+2}{\Delta_{\mathcal{L}}+1} \frac{k-\Delta_{\mathcal{L}}}{2k-\Delta_{\mathcal{L}}} \frac{\Delta_{\mathcal{R}}}{\Delta_{\mathcal{R}}+1} \frac{2k-\Delta_{\mathcal{R}}+1}{k-\Delta_{\mathcal{R}}+1} \\ &= \theta_1(\Delta_{\mathcal{L}}+1) \frac{1}{\theta_k(k-\Delta_{\mathcal{L}})} \frac{1}{\theta_1(\Delta_{\mathcal{R}})} \theta_k(k-\Delta_{\mathcal{R}}+1) \\ &= \frac{\theta_1(\Delta_{\mathcal{L}}+1)}{\theta_1(\Delta_{\mathcal{R}})} \frac{\theta_k(k+1-\Delta_{\mathcal{R}})}{\theta_k(k+1-(\Delta_{\mathcal{L}}+1))}, \end{aligned}$$

where $\theta_x(y) := \frac{x+y}{y} = 1 + \frac{x}{y}$. For positive x and y , $\theta_x(y)$ is decreasing in y , so for $\Delta_{\mathcal{L}}+1 \leq \Delta_{\mathcal{R}}$, $\psi \geq 1$. In other words, if $\Delta_{\mathcal{L}}+1 \leq \Delta_{\mathcal{R}}$, $\omega_{k,\Delta_{\mathcal{L}}} \omega_{k,\Delta_{\mathcal{R}}}$ increases when incrementing $\Delta_{\mathcal{L}}$ and decrementing $\Delta_{\mathcal{R}}$. By symmetry, it is maximal when

$$\Delta_{\mathcal{L}} = \left\lfloor \frac{\Gamma}{2} \right\rfloor \text{ and } \Delta_{\mathcal{R}} = \left\lceil \frac{\Gamma}{2} \right\rceil,$$

so that, when the number Γ of extreme points of \mathcal{P} is even,

$$\nu_{\mathcal{P}}(\langle k \rangle^k) \leq \left(\frac{\frac{\Gamma}{2}+1}{k+1} \binom{2k-\frac{\Gamma}{2}}{k} \right)^2 = \left(\frac{\frac{\Gamma}{2}+1}{\frac{n}{2}+1} \binom{n-\frac{\Gamma}{2}}{\frac{n}{2}} \right)^2.$$

Asymptotically, if we let $\Xi := \frac{\Gamma}{n}$ be proportion of the points that are extreme,

$$\begin{aligned} \nu_{\mathcal{P}}(\langle k \rangle^k) &\preceq_n \left(\frac{\left(1 - \frac{1}{2-\Xi}\right)^{\frac{1}{2-\Xi}-1}}{\left(\frac{1}{2-\Xi}\right)^{\frac{1}{2-\Xi}}} \right)^{2n-\Xi n} \\ &= \left(\frac{(2-\Xi)^{\frac{1}{2-\Xi}} (\Xi-1)^{\frac{1-\Xi}{\Xi-2}}}{(\Xi-2)^{\frac{1-\Xi}{\Xi-2}}} \right)^{(2-\Xi)n} \\ &= \left(\frac{(2-\Xi)(\Xi-1)^{\Xi-1}}{(\Xi-2)^{\Xi-1}} \right)^n \\ &= \left(\frac{(1-\Xi)^{\Xi-1}}{(2-\Xi)^{\Xi-2}} \right)^n. \end{aligned} \tag{3.1}$$

3. HIGHLY CONVEX MATCHINGS ACROSS A LINE

See figure 3.2 for a plot. The base of the exponential is 4 for $\Xi = 0$, and it tends to 1 as Ξ tends to 1. For $\Xi = \frac{1}{2}$, *i.e.*, when half the points are extreme, this gives an asymptotic bound of

$$\nu_{\mathcal{P}}(\langle^k \rangle^k) \preceq_n \left(\frac{3\sqrt{3}}{2} \right)^n \approx 2.6^n.$$




Figure 3.2 – Blue: the base of the exponential bound (3.1) as a function of the proportion of extreme points Ξ . Yellow: the base of the pre-existing bound. We have a 4^n bound from the Sharir-Welzl theorem on matchings across a line, and for $\eta = 1$ the bound becomes subexponential—and is tight, and is 1. Green: the bound (1.5) on all of $\nu_{\mathcal{P}}$ for comparison.

Chapter 4

Three changes of bracket direction

We now consider perfect matchings consistent with bracket expressions which have three changes of bracket directions, *i.e.*, bracket expressions of the form $\langle^a \rangle^b \langle^c \rangle^d$, a opening brackets, b closing brackets, c opening brackets, d closing brackets, where $a - b = d - c \geq 0$ and $a + b + c + d =: n$.

In term of points, this means that four sets can be separated by vertical lines, from left to right, a left-points forming the set \mathcal{A} , b right-points forming \mathcal{B} , d left-points forming \mathcal{D} , and c right-points forming \mathcal{C} . We pick a vertical line separating \mathcal{A} and \mathcal{B} and call it V , and we pick a vertical line separating \mathcal{C} and \mathcal{D} and call it W , see figure 4.1.

Given a crossing-free perfect matching on those points, numbering from top to bottom the intersections between V and edges incident to the points in \mathcal{A} , we get a crossing-free left-alignment α of \mathcal{A} on V . $a - b$ of the a edges intersecting V are incident to points in \mathcal{D} ; the other b are incident to points in \mathcal{B} . Numbering those b edges yields a crossing-free right-alignment β of \mathcal{B} on V .

Similarly on the right side, we get a crossing-free right-alignment of δ of \mathcal{D} on W , and numbering the subset of edges joining \mathcal{C} and \mathcal{D} , we get a left-alignment γ of \mathcal{C} on W . Call \mathcal{S} the set of $a - b$ points of \mathcal{A} that are matched with points of \mathcal{D} , and call \mathcal{T} the set of $d - c$ points of \mathcal{D} that are matched to points of \mathcal{A} .

The matching is uniquely determined by α , β , the choice of the subsets $\mathcal{S} \subseteq \mathcal{A}$ and $\mathcal{T} \subseteq \mathcal{D}$, and by β and χ . This gives the following bound for the number of these matchings:

$$\nu_{\mathcal{P}}(\langle^a \rangle^b \langle^c \rangle^d) \leq C_a \binom{a}{b} C_b C_c \binom{d}{c} C_d. \quad (4.1)$$



Figure 4.1 – Four crossing-free alignments.




Figure 4.2 – Blue: the base $4x_\eta$ of the exponential bound (4.1) for $\nu_{\mathcal{P}}(\langle a \rangle^b \langle b \rangle^a)$ as a function of $\eta = \frac{b}{a}$. Yellow: the base of the pre-existing bound. We have a 4^n bound from the Sharir-Welzl theorem on matchings across a line when $\eta = 0$ or $\eta = 1$, otherwise the pre-existing bound is (1.5). The bound is better than existing results for small and large η , but not for all η .

Asymptotically, the factor involving a and b is, for $b = \eta a$,

$$C_a \binom{a}{b} C_b \preccurlyeq_{a+b} 4^{a+b} \binom{a}{b} \preccurlyeq_{a+b} 4^{a+b} \underbrace{\left(\frac{(1-\eta)^{1-\frac{1}{\eta}}}{\eta} \right)^{\frac{\eta}{\eta+1}}}_{x_\eta}^{a+b},$$

uniformly in $\eta \in [0, 1]$. See figure 4.2 for a plot. The base x_η of the exponential bound for the binomial coefficient is maximal¹ when $\eta = \frac{3-\sqrt{5}}{2}$, where it

¹This can be shown by differentiating x_η . One can alternatively study the sequence $\max_b \binom{m-b}{b}$, as in the work of S. M. Tanny and M. Zuker [15].



Figure 4.3 – The points of $\mathcal{S} \subseteq \mathcal{A}$ and $\mathcal{T} \subseteq \mathcal{D}$ are shown in white. The vertical lines V and W bound the strip Σ . The long edges join \mathcal{S} and \mathcal{T} , partitioning the strip Σ in cells Z_i . The crossing-free right-alignments β_i are shown. Note that here \mathcal{B}_4 is empty. In this figure, $a = 7$, $b = 4$, $c = 1$, and $d = 4$.

is $\phi = \frac{1+\sqrt{5}}{2}$. This gives the overall bound

$$\nu_{\mathcal{P}}(\langle^a \rangle^b \langle^c \rangle^d) \preceq_n 4^{a+b} \phi^{a+b} 4^{c+d} \phi^{c+d} = (4\phi)^n \approx 6.472^n$$

uniformly in a , b , c , and d such that $a + b + c + d = n$. Note that this is worse than the existing 5.3793^n bound on $\nu_{\mathcal{P}}(B)$ for an arbitrary well-formed bracket expression B , see (1.5).

4.1 Improving upon the binomial bound

We can however improve upon that bound. Once α , δ , \mathcal{S} , and \mathcal{T} are fixed, the edges of the matching that cross both the left line and the right line—let us call these *long edges*—are determined. As a result, the strip Σ between the vertical lines V and W is partitioned by the long edges in trapezoidal cells Z_1, \dots, Z_{a-b+1} , numbered from top to bottom, where the first and last cells are unbounded, see figure 4.3.

We will consider the crossing-free right-alignment β ; the same argument will hold for γ . Since \mathcal{B} lies within the strip Σ , it is partitioned in $a - b + 1$ subsets $\mathcal{B}_i := \mathcal{B} \cap Z_i$. Moreover, an edge e_p emanating from a point $p \in \mathcal{B}_i$ cannot cross the long edges, and thus the portion $e_p \cap \Sigma$ of e_p that lies within the strip Σ is confined to the cell Z_i .

Let β_i be the crossing-free right-alignment of \mathcal{B}_i on V realized by the segments $\{e_p \cap \Sigma \mid p \in \mathcal{B}_i\}$. Since the set of segments $\{e_p \cap \Sigma \mid p \in \mathcal{B}\}$ realizes β , β is composed of the crossing-free right-alignments β_i . Formally, for



Figure 4.4 – The edges crossing V . The points of \mathcal{S} are shown in white. The crossing-free left-alignment α of \mathcal{A} on V is shown. The elements of $S = \alpha(\mathcal{S})$ are shown in red. We have $\text{cells}([a] \setminus S) = \{\{1\}, \{3, 4\}, \{6\}\}$. Since \mathcal{B}_4 is empty, there is no cell of $[a] \setminus S$ corresponding to it.

$$p \in \mathcal{B}_i,$$

$$\beta(p) = \beta_i(p) + \sum_{j=1}^{i-1} |\mathcal{B}_j|.$$

We now know that β is determined by crossing-free alignments β_i of point sets \mathcal{B}_i . In order to turn this into a bound, we need to know the sizes $|\mathcal{B}_i|$ of these point sets.

Let us look at the edges crossing V , numbered from top to bottom from 1 to a , see figure 4.4. Of these a edges, $a - b$ are long edges; they are the edges emanating from \mathcal{S} . Their numbers are $S = \alpha(\mathcal{S}) \subseteq [a]$. The other edges are the ones that emanate from points of \mathcal{B} . Let s_i be the i th element of S in increasing order, thus $s_1 = \min S$, and $s_{|S|} = s_{a-b} = \max S$.

The numbering of the a edges crossing V , from top to bottom, then starts with the edges emanating from \mathcal{B}_1 , numbered 1 through $s_1 - 1$, the first long edge, numbered s_1 , the edges emanating from \mathcal{B}_2 , numbered $s_1 + 1$ through $s_2 - 1$, the second long edge, numbered s_2 , the edges emanating from \mathcal{B}_3 , numbered from $s_2 + 1$, etc. In other words, the numbers of the edges emanating from any nonempty \mathcal{B}_i forms a maximal set of consecutive elements of $[a] \setminus S$. This motivates the following definition.

Definition 4.1 (cells of a set of integers) *The cells of $S' \subseteq \mathbb{N}$, written $\text{cells}(S')$, is the set of maximal subsets of consecutive elements S' .*

For instance, $\text{cells}(\{2, 3, 5, 6, 7, 10, 13, 14\}) = \{\{2, 3\}, \{5, 6, 7\}, \{10\}, \{13, 14\}\}$.

Then, we can improve the $\binom{a}{b} C_b$ factor in the bound (in which the binomial comes from the choice of \mathcal{S} , and the Catalan number comes from the choice

of β), summing over the choices of S (and thus of S above). The improved factor becomes²

$$\text{spc}(a, b) := \sum_{S \in \binom{[a]}{a-b}} \prod_{\chi \in \text{cells}([a] \setminus S)} C_{|\chi|} = \sum_{S' \in \binom{[a]}{b}} \prod_{\chi \in \text{cells}(S')} C_{|\chi|}, \quad (4.2)$$

and the overall bound becomes

$$\nu_{\mathcal{P}}(\langle a \rangle^b \langle c \rangle^d) \leq C_a \text{spc}(a, b) \text{spc}(d, c) C_d. \quad (4.3)$$

4.2 A recurrence

In order to compute spc efficiently, and eventually, get its asymptotics, it is useful to get rid of the cells function. We can express spc as a recurrence instead. First, we note that $\text{spc}(a, a) = C_a$: there is only one summand, S' is the whole set, so it has only one cell, namely S' itself. Otherwise, $a - b \geq 1$; in the sum over the S , consider the greatest element j of S , which is at least $a - b$, and split the sum over that,

$$\text{spc}(a, b) = \sum_{j=a-b}^a \sum_{\substack{S \in \binom{[a]}{a-b} \\ j = \max S}} \prod_{\chi \in \text{cells}([a] \setminus S)} C_{|\chi|}.$$

For fixed j , all summands (of the sum over S) will have a factor with $\chi = \{j+1, \dots, a\}$, and thus a factor of C_{a-j} . Factoring out this C_{a-j} , we get

$$\text{spc}(a, b) = \sum_{j=a-b}^a C_{a-j} \sum_{\substack{S \in \binom{[a]}{a-b} \\ j = \max S}} \prod_{\chi \in \text{cells}([a] \setminus S) \\ \chi \neq \{j+1, \dots, a\}} C_{|\chi|}.$$

Now, note that choosing a subset S of $[a]$ of size $a - b$ whose maximum is j is equivalent to choosing a subset \tilde{S} of $[j-1]$ of size $a - b - 1$, where $S = \tilde{S} \cup \{j\}$. Moreover, the cells of $[a] \setminus S$ other than $\{j+1, \dots, a\}$ are exactly the cells of $[j-1] \setminus \tilde{S}$, thus

$$\text{spc}(a, b) = \sum_{j=a-b}^a C_{a-j} \sum_{\substack{\tilde{S} \in \binom{[j-1]}{a-b-1}}} \prod_{\chi \in \text{cells}([j-1] \setminus \tilde{S})} C_{|\chi|}.$$

By definition of spc , this means

$$\text{spc}(a, b) = \sum_{j=a-b}^a C_{a-j} \text{spc}(j-1, b+j-a).$$

²The name spc was chosen because we define this as a Sum of Products of Catalan numbers.

1					
1	1				
1	2	2			
1	3	5	5		
1	4	9	14	14	
1	5	14	28	42	42

Figure 4.5 – The first few values of spc ; a vertically from 0 to 5, b horizontally from 0 to a . The values appearing in the $b + 1$ summands of the recurrence (4.4) with $a = 6$ and $b = 3$ are highlighted.

Rewriting this as a sum over $i := a - j$, this gives us the following recurrence for spc :

$$\text{spc}(a, b) = \sum_{i=0}^b C_i \text{spc}(a - i - 1, b - i) \quad \text{for } b < a, \quad (4.4)$$

$$\text{spc}(a, a) = C_a. \quad (4.5)$$

We can use this recurrence directly to get a very slack bound on the asymptotics of (4.3), see appendix B. Since the recurrence allows for memoization, we can also compute spc , and experimentally estimate its asymptotics, see appendix A. The 5^n bound that we can conjecture from these numerical calculations is much better than what we get by attempting to bound this recurrence, so this motivates looking for a better one, to ultimately solve spc .

4.3 A better recurrence, and a solution

We can now turn this recurrence into a simpler and better-known one, namely

$$\text{spc}(a, b) = \text{spc}(a - 1, b) + \text{spc}(a, b - 1) \quad \text{for } a \geq 1, 0 \leq b \leq a, \quad (4.6)$$

$$\text{spc}(a, b) = 0 \quad \text{for } b < 0 \text{ or } b > a, \quad (4.7)$$

$$\text{spc}(0, 0) = 1. \quad (4.8)$$

Since spc has not been formally defined for negative arguments, we extend the definition with $\text{spc}(a, b) = 0$ for $b < 0$; this is consistent with original the definition, since it yields a sum over subsets $S \subseteq [a]$ bigger than a , and it yields an empty sum in the recurrence we just derived. In addition to that, we also let $\text{spc}(a, b) = 0$ when $b > a$. This takes care of (4.7). Moreover, we know that $\text{spc}(0, 0) = C_0 = 1$. There thus remains to prove (4.6). We will do so first for $b = a$, then for $b = a - 1$, and finally for the $0 \leq b < a - 1$.

Note that we have

$$\text{spc}(b, b-1) = \sum_{i=0}^{b-1} C_i \text{spc}(b-i-1, b-i-1) = \sum_{i=0}^{b-1} C_i C_{b-i-1} = C_b, \quad (4.9)$$

and thus, for $a = b > 0$,

$$\text{spc}(a, b) = C_b = 0 + C_b = \text{spc}(a-1, b) + \text{spc}(a, b-1). \quad (4.10)$$

Further, for $a-1 = b$, we get

$$\text{spc}(a, b) = \sum_{i=0}^b C_i C_{b-i} = C_b + \sum_{i=0}^{b-1} C_i C_{b-i}$$

applying (4.5) on the left and (4.9) on the right,

$$\begin{aligned} &= \text{spc}(a-1, b) + \sum_{i=0}^{b-1} C_i \text{spc}(b-i, b-i-1) \\ &= \text{spc}(a-1, b) + \sum_{i=0}^{b-1} C_i \text{spc}(a-i-1, b-i-1) \\ &= \text{spc}(a-1, b) + \text{spc}(a, b-1). \end{aligned} \quad (4.11)$$

Now let $a-1 > b \geq 0$. Assume $\text{spc}(a', b') = \text{spc}(a'-1, b') + \text{spc}(a', b'-1)$ for $0 \leq b' < a' < a$. Then we can apply this assumption to the summands of $\text{spc}(a, b)$:

$$\begin{aligned} \text{spc}(a, b) &= \sum_{i=0}^b C_i \text{spc}(a-i-1, b-i) \\ &= \sum_{i=0}^b C_i (\text{spc}(a-i-2, b-i) + \text{spc}(a-i-1, b-i-1)) \\ &= \sum_{i=0}^b C_i \text{spc}(a-i-2, b-i) + \sum_{i=0}^b C_i \text{spc}(a-i-1, b-i-1) \\ &= \sum_{i=0}^b C_i \text{spc}((a-1)-i-1, b-i) + \sum_{i=0}^{b-1} C_i \text{spc}(a-i-1, (b-1)-i) \\ &\quad + \underbrace{C_b \text{spc}(a-b-1, -1)}_{=0}, \end{aligned}$$

so, by (4.4),

$$= \text{spc}(a-1, b) + \text{spc}(a, b-1). \quad (4.12)$$

This completes the proof of (4.6).

1				
1	1			
1	2	2		
1	3	5	5	
1	4	9	14	14
1	5	14	28	42

Figure 4.6 – The recurrence (4.6).

The recurrence we just proved defines a triangle of numbers sometimes known as the Catalan triangle³, and its solution is known, namely

$$\text{spc}(a, b) = \frac{(a+b)!(a-b+1)}{b!(a+1)!} = \frac{a-b+1}{a+1} \binom{a+b}{a}, \quad (4.13)$$

as shown by L. F. A. Arbogast in 1800 [2, p. 214–217].

4.3.1 A combinatorial interpretation of spc

The Catalan triangle counts well-formed prefixes; specifically, $\text{spc}(a, b)$ is the number of well-formed prefixes with a opening brackets and b closing brackets. This can in fact be seen from all the definitions of spc above, thus providing a more combinatorial solution, as we will now see.

In the original definition (4.2) of spc , the sum is over the choices for the positions of the $a - b$ unmatched opening brackets; between those, well-formed bracket expressions (counted by Catalan numbers) are inserted.

The first recurrence (4.4) corresponds to the following grammatical definition⁴ of well-formed prefixes, where again wfbe denotes a well-formed bracket expression:

$$\begin{aligned} \text{wfprefix} ::= & \text{wfbe} \langle \text{wfprefix} \\ & | \text{wfbe}. \end{aligned}$$

In this case the sum ranges over the length of the well-formed bracket expression preceding the first unmatched bracket.

Finally, (4.6) reflects the fact that a well-formed prefix with a opening brackets and b closing brackets either ends with an opening bracket (preceded by

³This triangle first appears as an example in a book by Arbogast in 1800 [2]. It features prominently in the previously-mentioned work of Errera [6], where it is attributed to Delannoy, who defined it in 1889 [4] as the number of paths by which a rook can reach a square on a triangular chessboard by going only down and right. The association with Catalan’s name seems to come from a 1976 paper by Shapiro [12], where a definition similar to Delannoy’s is independently given. In that paper, the name is justified by (4.5).

⁴This grammar requires infinite look-ahead to parse, but doing a first pass to mark the unmatched opening brackets resolves that.

a well-formed prefix with $a - 1$ opening and b closing brackets), or with a closing bracket (preceded by a well-formed prefix with a opening and $b - 1$ closing brackets).

4.3.2 Tying up some loose ends

We now solve the recurrence ω defined in chapter 3 by a combinatorial argument. Recall that ω is defined by

$$\begin{aligned}\omega_{k,\Delta} &:= \max_{\Delta_1+\Delta_2=\Delta-1} \sum_{i=\Delta_1}^{k-\Delta_2-1} \omega_{i,\Delta_1} \omega_{k-i-1,\Delta_2}, \\ \omega_{k,0} &:= \omega_{k,1} \text{ for } k > 0, \text{ and} \\ \omega_{k,k} &:= 1.\end{aligned}$$

We will show that $\omega(k, \Delta)$ is the number of well-formed prefixes with k opening brackets of which Δ are unmatched, thus that $\omega(k, \Delta) = \text{spc}(k, k - \Delta)$.

Let $\omega_{k,\Delta} := \text{spc}(k, k - \Delta)$ be the number of well-formed prefixes with k opening brackets of which Δ are unmatched. We have $\omega_{k,k} = 1$, since the only such prefix is \langle^k .

Consider $\omega_{k,0}$ where $k > 0$, *i.e.*, the number of well-formed bracket expressions of length k . Note that any nonempty well-formed bracket expression can be turned into a well-formed prefix with one unmatched opening bracket by removing the final closing bracket, and that adding a closing bracket to a well-formed prefix with one unmatched opening bracket yields a well-formed bracket expression. These operations are the inverse of each other, thus they are bijections, so $\omega_{k,0} = \omega_{k,1}$.

Finally, we look at the sum. For any nonnegative Δ_1 and Δ_2 such that $\Delta_1 + \Delta_2 = \Delta - 1$,

$$\sum_{i=\Delta_1}^{k-\Delta_2-1} \omega_{i,\Delta_1} \omega_{k-i-1,\Delta_2}$$

counts the number of ways of constructing a bracket expression with k opening brackets of which Δ are unmatched by chaining

1. a prefix containing the first Δ_1 unmatched opening brackets;
2. an opening bracket, which will be unmatched;
3. a prefix containing the last Δ_2 unmatched opening brackets.

Thus, for any nonnegative Δ_1 and Δ_2 such that $\Delta_1 + \Delta_2 = \Delta - 1$,

$$\omega_{k,\Delta} = \sum_{i=\Delta_1}^{k-\Delta_2-1} \omega_{i,\Delta_1} \omega_{k-i-1,\Delta_2}.$$

In particular,

$$\omega_{k,\Delta} = \max_{\Delta_1+\Delta_2=\Delta-1} \sum_{i=\Delta_1}^{k-\Delta_2-1} \omega_{i,\Delta_1} \omega_{k-i-1,\Delta_2}.$$

It follows that $\omega = \varpi$; this completes the proof. Substituting (4.13), we get

$$\omega_{k,\Delta} = \frac{\Delta+1}{k+1} \binom{2k-\Delta}{k}. \quad (4.14)$$

4.4 Asymptotics

We can now use the expression (4.13) for spc to study the properties of the bound (4.3). First, we note that in the extreme case where $b = c = 0$, so that the problem is that of matchings across a line, the bound becomes C_a^2 , *i.e.*, we get exactly the Sharir–Welzl bound. When $a = b$ and $c = d$, so that the bracket expression being considered is $\langle^a \rangle^a \langle^d \rangle^d$, by (4.5), we get a bound of $C_a^2 C_b^2$. This is the bound that is obtained from matchings across a line by chaining well-formed bracket expressions. For nondegenerate cases, we look at the asymptotics.

We are interested in the asymptotics as a function of the length $n = a + b + c + d$ of the bracket expression. Since the bound is a product of two identical two-parameter factors, we study one of them,

$$\varphi_{a,b} := C_a \text{spc}(a, b),$$

as a function of $m := a + b$. Let $b = \eta a$, thus $0 \leq \eta \leq 1$, the above expression becomes

$$\varphi_{\frac{m}{\eta+1}, \frac{m\eta}{\eta+1}} = C_{\frac{m}{\eta+1}} \binom{m}{\frac{m}{\eta+1}} \frac{m - m\eta + \eta + 1}{m + \eta + 1} \leq C_{\frac{m}{\eta+1}} \binom{m}{\frac{m}{\eta+1}}.$$

Asymptotically, this yields

$$\varphi_{\frac{m}{\eta+1}, \frac{m\eta}{\eta+1}} \lesssim_m 4^{\frac{m}{\eta+1}} \left(\frac{\left(1 - \frac{1}{\eta+1}\right)^{\frac{1}{\eta+1}-1}}{\left(\frac{1}{\eta+1}\right)^{\frac{1}{\eta+1}}} \right)^m = \underbrace{\left(4^{\frac{1}{\eta+1}} \frac{\eta+1}{\eta^{\frac{1}{\eta+1}}} \right)^m}_{y_\eta}$$

uniformly in $\eta \in [0, 1]$. See figure 4.7 for a plot.

With some calculus, one can see that as η ranges from 0 to 1, the base y_η of that exponential is maximal for $\eta = \frac{1}{4}$, where its value is 5. We thus have the following asymptotic bound:

$$\nu_{\mathcal{P}}(\langle^a \rangle^b \langle^c \rangle^d) \leq C_a \text{spc}(a, b) \text{spc}(d, c) C_d \lesssim_n 5^{a+b} 5^{c+d} = 5^n \quad (4.15)$$

uniformly in a, b, c , and d such that $a + b + c + d = n$.




Figure 4.7 – Blue: the base y_η of the exponential bound (4.3) for $\nu_{\mathcal{P}}(\langle^a \rangle^b \langle^b \rangle^a)$ as a function of $\eta = \frac{b}{a}$. Yellow: the base of the pre-existing bound. We have a 4^n bound from the Sharir-Welzl theorem on matchings across a line when $\eta = 0$ or $\eta = 1$, otherwise the pre-existing bound is (1.5).

Chapter 5

Outlook

As previously observed, uniform bounds on $\nu_{\mathcal{P}}$ can be used to bound the number of crossing-free perfect matchings, by summation over bracket expressions. The bound derived in the previous chapter only applies to a limited class of bracket expressions, and so it does not by itself yield such an overall upper bound.

Moreover, while we used cones and conifers to improve the bound on the number of matchings across a line, we did so only by looking at convexity properties, and we did not extend this reasoning to the bracket expressions considered in chapter 4.

We conclude this work by outlining a way in which the bound from chapter 4 may be extended to arbitrary bracket expressions, and by discussing how conifers can be used to further refine bounds regardless of convexity considerations.

Recall that $\text{spc}(a, b)$ counts the choices of the indices on the left side of the strip Σ of the long edges crossing Σ , together with (an upper bound on) right-alignments of the right-points by within that strip, separated by long edges. Looking at an arbitrary well-formed bracket expression, we can separate it into

1. a sequence of opening brackets \langle^a ;
2. a sequence of expressions of the form $\rangle^{b_i} \langle^{c_i}$;
3. a sequence of closing brackets \rangle^d .

The points \mathcal{B}_i and \mathcal{C}_i corresponding to the $\rangle^{b_i} \langle^{c_i}$ lie within vertical strips Σ_i , and an argument similar to section 4.1 can be made as follows.

Working from left to right, we pick a crossing-free right-alignment of the points \mathcal{A} corresponding to \langle^a . Then, looking at a strip Σ_i , where the yet-unmatched points to the left of Σ_i reach its left side in a known vertical

5. OUTLOOK

order, we choose which ones will cross the strip. This determines the size of the sets $\mathcal{B}_{ij} \subseteq \mathcal{B}_i$ of right-points within Σ_i that are separated by strip-crossing edges. Since the points from which the strip-crossing edges emanate are known, the sizes determine the sets. We then pick crossing-free left-alignments of the \mathcal{B}_{ij} . On the right side of the Σ_i , we pick the vertical indices of the strip-crossing edges, and this determines the sizes, of the sets $\mathcal{C}_{ij} \subseteq \mathcal{C}_i$ of right-points in Σ_i separated by strip-crossing edges, and thus the \mathcal{C}_{ij} themselves. We pick crossing-free left-alignments of the \mathcal{C}_{ij} , and move on to strip Σ_{i+1} . When finally we reach the points \mathcal{D} corresponding to \rangle^d , we pick a crossing-free right alignment of them.

This yields a bound of

$$\nu_{\mathcal{P}}(\langle^a \rangle^{b_1} \langle^{c_1} \dots \rangle^{b_r} \langle^{c_r} \rangle^d) \leq C_a C_d \prod_{i=1}^r \text{spc}(a + \tau(i), b_i) \text{spc}(a + \tau(i) - b_i, c_i),$$

where $\tau(i) := \sum_{j=1}^{i-1} (c_j - b_j)$. This bound is super-exponential: for the bracket expression $\langle^a \rangle \langle \dots \rangle \langle^a$, it yields $C_a^2 \text{spc}(a, 1)^{2r} = C_a^2 a^{2r}$. It may however be possible to improve it with overcounting analyses similar to the one we suggest below.

Let us come back to crossing-free perfect matchings across a line. We have, based on conifers,

$$\nu_{\mathcal{P}}(\langle^k \rangle^k) \leq C_k C_k = \left(\sum_{i=0}^{k-1} C_i C_{k-i-1} \right) \left(\sum_{j=0}^{k-1} C_j C_{k-j-1} \right),$$

where in the first sum i corresponds to the choice of the left \mathcal{L} -cone $\mathcal{K}_{\mathcal{L}, i+1}$, and in the second sum j corresponds to the choice of the right \mathcal{R} -cone $\mathcal{K}_{\mathcal{R}, j+1}$. When expanding the product of these sums, if $i = j$, we are counting crossing-free perfect matchings where the leftmost point is matched to the rightmost point. However, recall that to avoid overcounting, a point must be matched within its cone. Since there is a single left-cone of \mathcal{L} that contains the rightmost point, and a single right-cone of \mathcal{R} that contains the leftmost point, all but at most one of the $i = j$ terms are overcounted. We then get the bound

$$\nu_{\mathcal{P}}(\langle^k \rangle^k) \leq \max_{i \in \{0, \dots, k-1\}} (C_i C_{k-i-1})^2 + \sum_{i=0}^{k-1} \sum_{\substack{j=0 \\ j \neq i}}^{k-1} C_i C_{k-i-1} C_j C_{k-j-1}.$$

It may then be possible to extend this reasoning to the rest of the conifers to get an asymptotic improvement.

Appendix A

Experimental asymptotics of $C_a \text{spc}(a, b)$

Here we examine the exponential asymptotic behaviour of $C_a \text{spc}(a, b)$ as a function of $a + b$; this provides the asymptotics of the bound for $\nu_{\mathcal{P}}(\langle a \rangle^b \langle b \rangle^a)$. Specifically, we will be looking at the behaviour for constant $\eta = \frac{b}{a}$.

The definition (4.2) of spc ,

$$\text{spc}(a, b) := \sum_{S \in \binom{[a]}{a-b}} \prod_{\chi \in \text{cells}([a] \setminus S)} C_{|\chi|},$$

has exponentially many terms in the sum for constant η , so it is highly impractical to compute.

However, using recurrence (4.4),

$$\begin{aligned} \text{spc}(a, b) &= \sum_{i=0}^b C_i \text{spc}(a - i - 1, b - i) && \text{for } b < a, \\ \text{spc}(a, a) &= C_a, \end{aligned}$$

we can use memoization to compute $\text{spc}(a, b)$ in cubic time.

We use the following *Mathematica* code:

```
spc[a_, a_] := spc[a, a] = CatalanNumber[a];
spc[a_, b_] := spc[a, b] =
  Sum[spc[i, i] spc[a - i - 1, b - i], {i, 0, b}];
```

Note that we replace the Catalan number C_i in the summands by $\text{spc}(i, i)$ in order to benefit from the memoization of Catalan numbers. While this does

A. EXPERIMENTAL ASYMPTOTICS OF $C_a \text{spc}(a, b)$

not visibly change the order of the asymptotics for the values of i that we are considering, it improves performance by a noticeable constant.

This allows us to draw some plots (figure A.1) to take an educated guess at the behaviour of $C_a \text{spc}(a, b)$. From this we can conjecture that $\nu_{\mathcal{P}}(\langle a \rangle^b \langle b \rangle^a)$ is at most $\mathcal{O}(5^n)$.




Figure A.1 – Plots of $(4^a \text{spc}(a, b))^{\frac{1}{a+b}}$, where $a = \frac{b}{\eta}$, as a function of b , for various values of η . As b tends to infinity, this tends to the base of the exponential bound. We use 4^a rather than C_a because we want to get rid of as many polynomial factors as possible, since we are looking for the exponential behaviour. Note that the topmost curve seems to asymptotically approach 5.

Appendix B

An upper bound for spc using the ζ function

We can use the recurrence (4.4)

$$\begin{aligned}\text{spc}(a, b) &= \sum_{i=0}^b C_i \text{spc}(a - i - 1, b - i) && \text{for } b < a, \\ \text{spc}(a, a) &= C_a,\end{aligned}$$

to derive an exponential bound for $\text{spc}(a, b)$ as a function of $a + b$. We will do so by induction.

We first show an upper bound on the Catalan numbers which we will use in the induction.

Remark B.1 For $a \geq 1$, we have

$$C_a \leq \frac{4^a}{a^{\frac{3}{2}}\sqrt{\pi}} - \left(4\frac{\sqrt{2}}{\sqrt{\pi}} - 2\right). \quad (\text{B.1})$$

Proof We use the upper bound shown by R. D. Dutton and R. C. Brigham in [5],

$$C_a \leq \frac{4^a}{(a+1)\sqrt{a\pi}}.$$

It follows that

$$\frac{4^a}{a^{\frac{3}{2}}\sqrt{\pi}} - C_a \geq \frac{4^a}{a^{\frac{3}{2}}\sqrt{\pi}} - \frac{4^a}{(a+1)\sqrt{a\pi}} = \frac{4^a}{a^{\frac{3}{2}}(a+1)\sqrt{\pi}},$$

which is an increasing function of a for $a > 2$. For $n = 3$, we have

$$\frac{4^a}{a^{\frac{3}{2}}(a+1)\sqrt{\pi}} = \frac{16}{3\sqrt{3\pi}} > 4\frac{\sqrt{2}}{\sqrt{\pi}} - 2.$$

There thus remains to check (B.1) for $n = 1$ and $n = 2$. It holds for both, with equality at $n = 2$. \square

We now go back to bounding $\text{spc}(a, b)$. Let $b < a$. Define

$$\vartheta := 1 + \frac{\zeta(\frac{3}{2})}{\sqrt{\pi}} - \frac{1}{3} \left(4 \frac{\sqrt{2}}{\sqrt{\pi}} - 2 \right),$$

and observe that $\vartheta \approx 2.0767 > 2$. The reason for this definition will become clear at the end of the calculation. Assuming that for all $\text{spc}(a', b')$ involved in the sum for $\text{spc}(a, b)$,

$$\text{spc}(a', b') \leq \vartheta^{a'+b'},$$

we get

$$\begin{aligned} \text{spc}(a, b) &= \sum_{i=0}^b C_i \text{spc}(a-i-1, b-i) \\ &\leq \sum_{i=0}^b C_i \vartheta^{a+b-2i-1} \\ &= \vartheta^{a+b-1} \left(1 + \sum_{i=1}^b C_i \vartheta^{-2i} \right) \end{aligned}$$

using (B.1),

$$\leq \vartheta^{a+b-1} \left(1 + \sum_{i=1}^b \left(\frac{4^i}{i^{\frac{3}{2}} \sqrt{\pi}} - \left(4 \frac{\sqrt{2}}{\sqrt{\pi}} - 2 \right) \right) \vartheta^{-2i} \right)$$

with $\vartheta > 2$,

$$\begin{aligned} &\leq \vartheta^{a+b-1} \left(1 + \sum_{i=1}^b \left(\frac{4^i}{i^{\frac{3}{2}} \sqrt{\pi}} - \left(4 \frac{\sqrt{2}}{\sqrt{\pi}} - 2 \right) \right) 4^{-i} \right) \\ &= \vartheta^{a+b-1} \left(1 + \sum_{i=1}^b \left(\frac{1}{i^{\frac{3}{2}} \sqrt{\pi}} - \left(4 \frac{\sqrt{2}}{\sqrt{\pi}} - 2 \right) 4^{-i} \right) \right) \\ &\leq \vartheta^{a+b-1} \left(1 + \sum_{i=1}^{\infty} \left(\frac{1}{i^{\frac{3}{2}} \sqrt{\pi}} - \left(4 \frac{\sqrt{2}}{\sqrt{\pi}} - 2 \right) 4^{-i} \right) \right) \\ &= \vartheta^{a+b-1} \left(1 + \frac{\zeta(\frac{3}{2})}{\sqrt{\pi}} - \frac{1}{3} \left(4 \frac{\sqrt{2}}{\sqrt{\pi}} - 2 \right) \right) \\ &= \vartheta^{a+b}. \end{aligned}$$

We start the induction for $a = b$ with $\text{spc}(a, a) = C_a \leq 4^a < \vartheta^{a+a}$. This gives us the bound

$$\text{spc}(a, b) \leq \vartheta^{a+b},$$

and so, with $\eta = \frac{b}{a}$,

$$\nu_{\mathcal{P}}(\langle^a \rangle^b \langle^b \rangle^a) \leq (C_a \text{spc}(a, b))^2 \leq (4^{\frac{1}{\eta+1}} \vartheta)^n. \quad (\text{B.2})$$

A plot of that bound is shown in figure B.1.




Figure B.1 – Blue: the base of the exponential bound (B.2) as a function of η . Yellow: pre-existing bounds. Green: the bound (4.1), for comparison. Red: the actual asymptotics of $C_a \text{spc}(a, b)$.

Bibliography

- [1] Miklós Ajtai, Václav Chvátal, Monroe M. Newborn, and Endre Szemerédi. Crossing-free subgraphs. In Peter L. Hammer, Alexander Rosa, Gert Sabidussi, and Jean Turgeon, editors, *Theory and Practice of Combinatorics — A collection of articles honoring Anton Kotzig on the occasion of his sixtieth birthday*, volume 60 of *North-Holland Mathematics Studies*, pages 9–12. North-Holland, 1982.
- [2] Louis François Antoine Arbogast. *Du Calcul des Dérivations*. Levrault, Frères, 1800.
- [3] Eugène Charles Catalan. Note sur une équation aux différences finies. *Journal de Mathématiques Pures et Appliquées*, 1(3):508–516, 1838.
- [4] Henri Auguste Delannoy. Emploi de l'échiquier pour la résolution de divers problèmes de probabilité. In *Compte rendu de la 18^{me} session : congrès de Paris—1889*, pages 43–52. Association française pour l'avancement des sciences, fusionnée avec l'association scientifique de France (fondée par Le Verrier en 1864), 1890.
- [5] Ronald D. Dutton and Robert C. Brigham. Computationally efficient bounds for the catalan numbers. *European Journal of Combinatorics*, 7(3):211–213, 1986.
- [6] Alfred Errera. Analysis situs : un problème d'énumération. *Mémoires de la Classe des Sciences de l'Académie royale de Belgique, série 2, collection in-8^o*, 11(6), 1931.
- [7] Leonhard Euler. Letter to Christian Goldbach. <http://eulerarchive.maa.org/correspondence/letters/000868.pdf> (retrieved 2016-09-10), 1751.

- [8] Alfredo García, Marc Noy, and Javier Tejel. Lower bounds on the number of crossing-free subgraphs of K_N . *Computational Geometry*, 16(4):211–221, 2000.
- [9] Monroe M. Newborn and William Oscar Jules Moser. Optimal crossing-free Hamiltonian circuit drawings of K_n . *Journal of Combinatorial Theory, Series B*, 29(1):13–26, 1980.
- [10] Andres J. Ruiz-Vargas and Emo Welzl. Crossing-free perfect matchings in wheel point sets. To appear.
- [11] János András Segner. Enumeratio modorum quibus figurae planae rectilineae per diagonales diuiduntur in triangula. In *Novi Commentarii Academiae Scientiarum Imperialis Petropolitanae*, volume 7 *pro annis 1758 et 1759*, pages 203–209, 1761.
- [12] Louis W. Shapiro. A Catalan triangle. *Discrete Mathematics*, 14(1):83–90, 1974.
- [13] Micha Sharir and Emo Welzl. On the number of crossing-free matchings, cycles, and partitions. *SIAM Journal on Computing*, 36(3):695–720, 2006.
- [14] Warren Douglas Smith. *Studies in Computational Geometry Motivated by Mesh Generation*. PhD thesis, Princeton, 1989. AAI9002713.
- [15] Stephen M. Tanny and Michael Zuker. On a unimodal sequence of binomial coefficients. *Discrete Mathematics*, 9(1):79–89, 1974.
- [16] Jan van Leeuwen and Anneke A. Schoone. Untangling a travelling salesman tour in the plane. In J. R. Mühlbacher, editor, *Proceedings of the 7th Conference on Graph-theoretic Concepts in Computer Science*, pages 87–98, 1981.
- [17] Klaus Wagner. Bemerkungen zum Vierfarbenproblem. *Jahresbericht der Deutschen Mathematiker-Vereinigung*, 46:26–32, 1936.



Eidgenössische Technische Hochschule Zürich
Swiss Federal Institute of Technology Zurich

Declaration of originality

The signed declaration of originality is a component of every semester paper, Bachelor's thesis, Master's thesis and any other degree paper undertaken during the course of studies, including the respective electronic versions.

Lecturers may also require a declaration of originality for other written papers compiled for their courses.

I hereby confirm that I am the sole author of the written work here enclosed and that I have compiled it in my own words. Parts excepted are corrections of form and content by the supervisor.

Title of work (in block letters):

Authored by (in block letters):

For papers written by groups the names of all authors are required.

Name(s):

First name(s):

With my signature I confirm that

- I have committed none of the forms of plagiarism described in the '[Citation etiquette](#)' information sheet.
- I have documented all methods, data and processes truthfully.
- I have not manipulated any data.
- I have mentioned all persons who were significant facilitators of the work.

I am aware that the work may be screened electronically for plagiarism.

Place, date

Signature(s)

For papers written by groups the names of all authors are required. Their signatures collectively guarantee the entire content of the written paper.