

# Deep Generative Models

## Lecture 7

Roman Isachenko



AI Masters

Autumn, 2022

# Recap of previous lecture

## Gaussian AR NF

$$\mathbf{x} = g(\mathbf{z}, \boldsymbol{\theta}) \quad \Rightarrow \quad \mathbf{x}_j = \sigma_j(\mathbf{x}_{1:j-1}) \cdot \mathbf{z}_j + \mu_j(\mathbf{x}_{1:j-1}).$$

$$\mathbf{z} = f(\mathbf{x}, \boldsymbol{\theta}) \quad \Rightarrow \quad \mathbf{z}_j = (\mathbf{x}_j - \mu_j(\mathbf{x}_{1:j-1})) \cdot \frac{1}{\sigma_j(\mathbf{x}_{1:j-1})}.$$

- ▶ Sampling is sequential, density estimation is parallel.
- ▶ Forward KL is a natural loss.

## Inverse gaussian AR NF

$$\mathbf{x} = g(\mathbf{z}, \boldsymbol{\theta}) \quad \Rightarrow \quad \mathbf{x}_j = \tilde{\sigma}_j(\mathbf{z}_{1:j-1}) \cdot \mathbf{z}_j + \tilde{\mu}_j(\mathbf{z}_{1:j-1})$$

$$\mathbf{z} = f(\mathbf{x}, \boldsymbol{\theta}) \quad \Rightarrow \quad \mathbf{z}_j = (\mathbf{x}_j - \tilde{\mu}_j(\mathbf{z}_{1:j-1})) \cdot \frac{1}{\tilde{\sigma}_j(\mathbf{z}_{1:j-1})}.$$

- ▶ Sampling is parallel, density estimation is sequential.
- ▶ Reverse KL is a natural loss.

## Recap of previous lecture

Let split  $\mathbf{x}$  and  $\mathbf{z}$  in two parts:

$$\mathbf{x} = [\mathbf{x}_1, \mathbf{x}_2] = [\mathbf{x}_{1:d}, \mathbf{x}_{d+1:m}]; \quad \mathbf{z} = [\mathbf{z}_1, \mathbf{z}_2] = [\mathbf{z}_{1:d}, \mathbf{z}_{d+1:m}].$$

### Coupling layer

$$\begin{cases} \mathbf{x}_1 = \mathbf{z}_1; \\ \mathbf{x}_2 = \mathbf{z}_2 \odot \sigma(\mathbf{z}_1, \theta) + \mu(\mathbf{z}_1, \theta). \end{cases} \quad \begin{cases} \mathbf{z}_1 = \mathbf{x}_1; \\ \mathbf{z}_2 = (\mathbf{x}_2 - \mu(\mathbf{x}_1, \theta)) \odot \frac{1}{\sigma(\mathbf{x}_1, \theta)}. \end{cases}$$

Estimating the density takes 1 pass, sampling takes 1 pass!

### Jacobian

$$\det \left( \frac{\partial \mathbf{z}}{\partial \mathbf{x}} \right) = \det \begin{pmatrix} \mathbf{I}_d & \mathbf{0}_{d \times m-d} \\ \frac{\partial \mathbf{z}_2}{\partial \mathbf{x}_1} & \frac{\partial \mathbf{z}_2}{\partial \mathbf{x}_2} \end{pmatrix} = \prod_{j=1}^{m-d} \frac{1}{\sigma_j(\mathbf{x}_1, \theta)}.$$

Coupling layer is a special case of autoregressive flow.

## Recap of previous lecture

	VAE	NF
Objective	ELBO $\mathcal{L}$	Forward KL/MLE
Encoder	stochastic $\mathbf{z} \sim q(\mathbf{z} \mathbf{x}, \phi)$	deterministic $\mathbf{z} = f(\mathbf{x} \theta)$ $q(\mathbf{z} \mathbf{x}, \theta) = \delta(\mathbf{z} - f(\mathbf{x}, \theta))$
Decoder	stochastic $\mathbf{x} \sim p(\mathbf{x} \mathbf{z}, \theta)$	deterministic $\mathbf{x} = g(\mathbf{z} \theta)$ $p(\mathbf{x} \mathbf{z}, \theta) = \delta(\mathbf{x} - g(\mathbf{z}, \theta))$
Parameters	$\phi, \theta$	$\theta \equiv \phi$

### Theorem

MLE for normalizing flow is equivalent to maximization of ELBO for VAE model with deterministic encoder and decoder:

$$p(\mathbf{x}|\mathbf{z}, \theta) = \delta(\mathbf{x} - f^{-1}(\mathbf{z}, \theta)) = \delta(\mathbf{x} - g(\mathbf{z}, \theta));$$

$$q(\mathbf{z}|\mathbf{x}, \theta) = p(\mathbf{z}|\mathbf{x}, \theta) = \delta(\mathbf{z} - f(\mathbf{x}, \theta)).$$

# Outline

## 1. Discrete data vs continuous model

Discretization of continuous distribution  
Dequantization of discrete data

## 2. ELBO surgery

## 3. VAE limitations

VAE prior

VAE posterior

# Outline

## 1. Discrete data vs continuous model

Discretization of continuous distribution  
Dequantization of discrete data

## 2. ELBO surgery

## 3. VAE limitations

VAE prior

VAE posterior

## Discrete data vs continuous model

Let our data  $\mathbf{y}$  comes from discrete distribution  $\Pi(\mathbf{y})$  and we have continuous model  $p(\mathbf{x}|\theta) = \text{NN}(\mathbf{x}, \theta)$ .

- ▶ Images (and not only images) are discrete data, pixels lie in the integer domain  $\{0, 255\}$ .
- ▶ By fitting a continuous density model  $p(\mathbf{x}|\theta)$  to discrete data  $\Pi(\mathbf{y})$ , one can produce a degenerate solution with all probability mass on discrete values.

## Discrete model

- ▶ Use **discrete** model (e.x.  $P(\mathbf{y}|\theta) = \text{Cat}(\pi(\theta))$ ).
- ▶ Minimize any suitable divergence measure  $D(\Pi, P)$ .
- ▶ NF works only with continuous data  $\mathbf{x}$  (there are discrete NF, see papers below).
- ▶ If pixel value is not presented in the train data, it won't be predicted.

---

*Hoogeboom E. et al. Integer discrete flows and lossless compression*


*Tran D. et al. Discrete flows: Invertible generative models of discrete data*

# Discrete data vs continuous model

## Continuous model

- ▶ Use **continuous** model (e.x.  $p(\mathbf{x}|\theta) = \mathcal{N}(\mu_{\theta}(\mathbf{x}), \sigma_{\theta}^2(\mathbf{x}))$ ), but
  - ▶ **discretize** model (make the model outputs discrete): transform  $p(\mathbf{x}|\theta)$  to  $P(\mathbf{y}|\theta)$ ;
  - ▶ **dequantize** data (make the data continuous): transform  $\Pi(\mathbf{y})$  to  $\pi(\mathbf{x})$ .
- ▶ Continuous distribution know numerical relationships.

## CIFAR-10 pixel values distribution





# Outline

## 1. Discrete data vs continuous model

Discretization of continuous distribution

Dequantization of discrete data

## 2. ELBO surgery

## 3. VAE limitations

VAE prior

VAE posterior

# Discretization of continuous distribution

## Model discretization through CDF

$$F(\mathbf{x}|\theta) = \int_{-\infty}^{\mathbf{x}'} p(\mathbf{x}'|\theta) d\mathbf{x}'; \quad P(\mathbf{y}|\theta) = F(\mathbf{y} + 0.5|\theta) - F(\mathbf{y} - 0.5|\theta)$$

## Mixture of logistic distributions

$$p(x|\mu, s) = \frac{\exp^{-(x-\mu)/s}}{s(1 + \exp^{-(x-\mu)/s})^2}; \quad p(x|\pi, \mu, s) = \sum_{k=1}^K \pi_k p(x|\mu_k, s_k).$$

## PixelCNN++

$$p(\mathbf{x}|\theta) = \prod_{j=1}^m p(x_j|\mathbf{x}_{1:j-1}, \theta); \quad p(x_j|\mathbf{x}_{1:j-1}, \theta) = \sum_{k=1}^K \pi_k p(x|\mu_k, s_k).$$

Here,  $\pi_k = \pi_{k,\theta}(\mathbf{x}_{1:j-1})$ ,  $\mu_k = \mu_{k,\theta}(\mathbf{x}_{1:j-1})$ ,  $s_k = s_{k,\theta}(\mathbf{x}_{1:j-1})$ .

For the pixel edge cases of 0, replace  $x - 0.5$  by  $-\infty$ , and for 255 replace  $x + 0.5$  by  $+\infty$ .

---

Salimans T. et al. *PixelCNN++: Improving the PixelCNN with Discretized Logistic Mixture Likelihood and Other Modifications*, 2017

# Outline

## 1. Discrete data vs continuous model

Discretization of continuous distribution

Dequantization of discrete data

## 2. ELBO surgery

## 3. VAE limitations

VAE prior

VAE posterior


# Uniform discretization

Let discrete distribution  $\Pi(\mathbf{y})$  to continuous distribution  $\pi(\mathbf{x})$  in the following way:  $\mathbf{x} = \mathbf{y} + \mathbf{u}$ , where  $\mathbf{u} \sim U[0, 1]$ .

## Theorem

Fitting continuous model  $p(\mathbf{x}|\theta)$  on uniformly dequantized data is equivalent to maximization of a lower bound on log-likelihood for a discrete model:



$$P(\mathbf{y}|\theta) = \int_{U[0,1]} p(\mathbf{y} + \mathbf{u}|\theta) d\mathbf{u}$$



## Proof

$$\begin{aligned} \mathbb{E}_{\pi} \log p(\mathbf{x}|\theta) &= \int \pi(\mathbf{x}) \log p(\mathbf{x}|\theta) d\mathbf{y} = \sum \Pi(\mathbf{y}) \int_{U[0,1]} \log p(\mathbf{y} + \mathbf{u}|\theta) d\mathbf{u} \leq \\ &\leq \sum \Pi(\mathbf{y}) \log \int_{U[0,1]} p(\mathbf{y} + \mathbf{u}|\theta) d\mathbf{u} = \\ &= \sum \Pi(\mathbf{y}) \log P(\mathbf{y}|\theta) = \mathbb{E}_{\Pi} \log P(\mathbf{y}|\theta). \end{aligned}$$

# Variational dequantization



- ▶  $p(\mathbf{x}|\boldsymbol{\theta})$  assign uniform density to unit hypercubes  $\mathbf{y} + U[0, 1]$  (left fig).
- ▶ Smooth dequantization is more natural (right fig).
- ▶ Neural network density models are smooth function approximators.

Introduce variational dequantization noise distribution  $q(\mathbf{u}|\mathbf{y})$ , which tells what kind of noise we have to add to our discrete data. Treat it as an approximate posterior as in VAE model.

# Variational dequantization

## Variational lower bound

$$\begin{aligned}\log P(\mathbf{y}|\boldsymbol{\theta}) &= \left[ \log \int q(\mathbf{u}|\mathbf{y}) \frac{p(\mathbf{y} + \mathbf{u}|\boldsymbol{\theta})}{q(\mathbf{u}|\mathbf{y})} d\mathbf{u} \right] \geq \\ &\geq \int q(\mathbf{u}|\mathbf{y}) \log \frac{p(\mathbf{y} + \mathbf{u}|\boldsymbol{\theta})}{q(\mathbf{u}|\mathbf{y})} d\mathbf{u} = \mathcal{L}(q, \boldsymbol{\theta}).\end{aligned}$$

Uniform dequantization is a special case of variational dequantization ( $q(\mathbf{u}|\mathbf{x}) = U[0, 1]$ ).

## Flow++: flow-based variational dequantization

Let  $\mathbf{u} = g(\epsilon, \mathbf{x}, \boldsymbol{\lambda})$  is a flow model with base distribution  $\epsilon \sim p(\epsilon)$ :

$$q(\mathbf{u}|\mathbf{x}) = p(f(\mathbf{u}, \mathbf{x}, \boldsymbol{\lambda})) \cdot \left| \det \frac{\partial f(\mathbf{u}, \mathbf{x}, \boldsymbol{\lambda})}{\partial \mathbf{u}} \right|.$$

$$\log P(\mathbf{x}|\boldsymbol{\theta}) \geq \mathcal{L}(\boldsymbol{\lambda}, \boldsymbol{\theta}) = \int p(\epsilon) \log \left( \frac{p(\mathbf{x} + g(\epsilon, \mathbf{x}, \boldsymbol{\lambda})|\boldsymbol{\theta})}{p(\epsilon) \cdot |\det \mathbf{J}_g|^{-1}} \right) d\epsilon.$$

# Outline

## 1. Discrete data vs continuous model

Discretization of continuous distribution

Dequantization of discrete data

## 2. ELBO surgery

## 3. VAE limitations

VAE prior

VAE posterior

# ELBO surgery

$$\frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(q, \theta) = \frac{1}{n} \sum_{i=1}^n \left[ \mathbb{E}_{q(\mathbf{z}|\mathbf{x}_i)} \log p(\mathbf{x}_i|\mathbf{z}, \theta) - KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) \right].$$

## Theorem

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) = \textcolor{violet}{KL(q_{\text{agg}}(\mathbf{z})||p(\mathbf{z}))} + \textcolor{teal}{\mathbb{I}_q[\mathbf{x}, \mathbf{z}]}$$

- ▶  $q_{\text{agg}}(\mathbf{z}) = \frac{1}{n} \sum_{i=1}^n q(\mathbf{z}|\mathbf{x}_i)$  – **aggregated** posterior distribution.
- ▶  $\mathbb{I}_q[\mathbf{x}, \mathbf{z}]$  – mutual information between  $\mathbf{x}$  and  $\mathbf{z}$  under empirical data distribution and distribution  $q(\mathbf{z}|\mathbf{x})$ .
- ▶ **First term** pushes  $q_{\text{agg}}(\mathbf{z})$  towards the prior  $p(\mathbf{z})$ .
- ▶ **Second term** reduces the amount of information about  $\mathbf{x}$  stored in  $\mathbf{z}$ .



# ELBO surgery

## Theorem

$$\frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) = KL(q_{\text{agg}}(\mathbf{z})||p(\mathbf{z})) + \mathbb{I}_q[\mathbf{x}, \mathbf{z}].$$

## Proof

$$\begin{aligned} \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z})) &= \frac{1}{n} \sum_{i=1}^n \int q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z}|\mathbf{x}_i)}{p(\mathbf{z})} d\mathbf{z} = \\ &= \frac{1}{n} \sum_{i=1}^n \int q(\mathbf{z}|\mathbf{x}_i) \log \frac{q_{\text{agg}}(\mathbf{z})q(\mathbf{z}|\mathbf{x}_i)}{p(\mathbf{z})q_{\text{agg}}(\mathbf{z})} d\mathbf{z} = \int \frac{1}{n} \sum_{i=1}^n q(\mathbf{z}|\mathbf{x}_i) \log \frac{q_{\text{agg}}(\mathbf{z})}{p(\mathbf{z})} d\mathbf{z} + \\ &+ \frac{1}{n} \sum_{i=1}^n \int q(\mathbf{z}|\mathbf{x}_i) \log \frac{q(\mathbf{z}|\mathbf{x}_i)}{q_{\text{agg}}(\mathbf{z})} d\mathbf{z} = KL(q_{\text{agg}}(\mathbf{z})||p(\mathbf{z})) + \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||q_{\text{agg}}(\mathbf{z})) \end{aligned}$$

Without proof:

$$\mathbb{I}_q[\mathbf{x}, \mathbf{z}] = \frac{1}{n} \sum_{i=1}^n KL(q(\mathbf{z}|\mathbf{x}_i)||q_{\text{agg}}(\mathbf{z})) \in [0, \log n].$$

# ELBO surgery

## ELBO revisiting

$$\begin{aligned}\frac{1}{n} \sum_{i=1}^n \mathcal{L}_i(q, \theta) &= \frac{1}{n} \sum_{i=1}^n [\mathbb{E}_{q(\mathbf{z}|\mathbf{x}_i)} \log p(\mathbf{x}_i|\mathbf{z}, \theta) - KL(q(\mathbf{z}|\mathbf{x}_i)||p(\mathbf{z}))] = \\ &= \underbrace{\frac{1}{n} \sum_{i=1}^n \mathbb{E}_{q(\mathbf{z}|\mathbf{x}_i)} \log p(\mathbf{x}_i|\mathbf{z}, \theta)}_{\text{Reconstruction loss}} - \underbrace{\mathbb{I}_q[\mathbf{x}, \mathbf{z}]}_{\text{MI}} - \underbrace{KL(q_{\text{agg}}(\mathbf{z})||p(\mathbf{z}))}_{\text{Marginal KL}}\end{aligned}$$

Prior distribution  $p(\mathbf{z})$  is only in the last term.

## Optimal VAE prior

$$KL(q_{\text{agg}}(\mathbf{z})||p(\mathbf{z})) = 0 \quad \Leftrightarrow \quad p(\mathbf{z}) = q_{\text{agg}}(\mathbf{z}) = \frac{1}{n} \sum_{i=1}^n q(\mathbf{z}|\mathbf{x}_i).$$

The optimal prior  $p(\mathbf{z})$  is the aggregated posterior  $q_{\text{agg}}(\mathbf{z})$ !

---

Hoffman M. D., Johnson M. J. *ELBO surgery: yet another way to carve up the variational evidence lower bound*, 2016

# Outline

## 1. Discrete data vs continuous model

Discretization of continuous distribution  
Dequantization of discrete data

## 2. ELBO surgery

## 3. VAE limitations

VAE prior

VAE posterior

# Outline

## 1. Discrete data vs continuous model

Discretization of continuous distribution

Dequantization of discrete data

## 2. ELBO surgery

## 3. VAE limitations

VAE prior

VAE posterior

# VAE limitations

- ▶ Poor generative distribution (decoder)

$$p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_{\boldsymbol{\theta}}(\mathbf{z}), \boldsymbol{\sigma}_{\boldsymbol{\theta}}^2(\mathbf{z})) \quad \text{or} \quad = \text{Softmax}(\boldsymbol{\pi}_{\boldsymbol{\theta}}(\mathbf{z})).$$

- ▶ Loose lower bound

$$\log p(\mathbf{x}|\boldsymbol{\theta}) - \mathcal{L}(q, \boldsymbol{\theta}) = (?).$$

- ▶ **Poor prior distribution**

$$p(\mathbf{z}) = \mathcal{N}(0, \mathbf{I}).$$

- ▶ Poor variational posterior distribution (encoder)


$$q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi}) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_{\boldsymbol{\phi}}(\mathbf{x}), \boldsymbol{\sigma}_{\boldsymbol{\phi}}^2(\mathbf{x})).$$

# Optimal VAE prior


How to choose the optimal  $p(\mathbf{z})$ ?

- ▶ Standard Gaussian  $p(\mathbf{z}) = \mathcal{N}(0, I) \Rightarrow$  over-regularization;
- ▶  $p(\mathbf{z}) = q_{\text{agg}}(\mathbf{z}) = \frac{1}{n} \sum_{i=1}^n q(\mathbf{z}|\mathbf{x}_i) \Rightarrow$  overfitting and highly expensive.

Non learnable prior  $p(\mathbf{z})$



Learnable prior  $p(\mathbf{z}|\lambda)$



# Flows-based VAE prior

## Flow model in latent space

$$\log p(\mathbf{z}|\boldsymbol{\lambda}) = \log p(\mathbf{z}^*) + \log \left| \det \left( \frac{d\mathbf{z}^*}{d\mathbf{z}} \right) \right| = \log p(g(\mathbf{z}, \boldsymbol{\lambda})) + \log |\det(\mathbf{J}_g)|$$

$$\mathbf{z} = f(\mathbf{z}^*, \boldsymbol{\lambda}) = g^{-1}(\mathbf{z}^*, \boldsymbol{\lambda})$$

- ▶ RealNVP flow.
- ▶ Autoregressive flow (MAF).

Why it is not a good idea to use IAF for VAE prior?

## ELBO with flow-based VAE prior

$$\begin{aligned} \mathcal{L}(\phi, \theta) &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} [\log p(\mathbf{x}|\mathbf{z}, \theta) + \log p(\mathbf{z}|\boldsymbol{\lambda}) - \log q(\mathbf{z}|\mathbf{x}, \phi)] \\ &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \left[ \log p(\mathbf{x}|\mathbf{z}, \theta) + \underbrace{\left( \log p(g(\mathbf{z}, \boldsymbol{\lambda})) + \log |\det(\mathbf{J}_g)| \right)}_{\text{flow-based prior}} - \log q(\mathbf{z}|\mathbf{x}, \phi) \right] \end{aligned}$$

# Outline

## 1. Discrete data vs continuous model

Discretization of continuous distribution  
Dequantization of discrete data

## 2. ELBO surgery

## 3. VAE limitations

VAE prior

VAE posterior



# VAE limitations

- ▶ Poor generative distribution (decoder)

$$p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_{\boldsymbol{\theta}}(\mathbf{z}), \boldsymbol{\sigma}_{\boldsymbol{\theta}}^2(\mathbf{z})) \quad \text{or} \quad = \text{Softmax}(\boldsymbol{\pi}_{\boldsymbol{\theta}}(\mathbf{z})).$$

- ▶ Loose lower bound

$$\log p(\mathbf{x}|\boldsymbol{\theta}) - \mathcal{L}(q, \boldsymbol{\theta}) = (?).$$

- ▶ Poor prior distribution

$$p(\mathbf{z}) = \mathcal{N}(0, \mathbf{I}).$$

- ▶ **Poor variational posterior distribution (encoder)**

$$q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi}) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_{\boldsymbol{\phi}}(\mathbf{x}), \boldsymbol{\sigma}_{\boldsymbol{\phi}}^2(\mathbf{x})).$$

# Variational posterior

## ELBO

$$\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(q, \boldsymbol{\theta}) + KL(q(\mathbf{z}|\mathbf{x}, \phi) || p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})).$$

- ▶ In E-step of EM-algorithm we wish  $KL(q(\mathbf{z}|\mathbf{x}, \phi) || p(\mathbf{z}|\mathbf{x}, \boldsymbol{\theta})) = 0$ .  
(In this case the lower bound is tight  $\log p(\mathbf{x}|\boldsymbol{\theta}) = \mathcal{L}(q, \boldsymbol{\theta})$ ).
- ▶ Normal variational distribution  $q(\mathbf{z}|\mathbf{x}, \phi) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_\phi(\mathbf{x}), \boldsymbol{\sigma}_\phi^2(\mathbf{x}))$  is poor (e.g. has only one mode).
- ▶ Flows models convert a simple base distribution to a complex one using invertible transformation with simple Jacobian. How to use flows in VAE posterior?

## Flows in VAE posterior

Apply a sequence of transformations to the random variable

$$\mathbf{z} \sim q(\mathbf{z}|\mathbf{x}, \phi) = \mathcal{N}(\mathbf{z}|\mu_{\phi}(\mathbf{x}), \sigma_{\phi}^2(\mathbf{x})).$$

Let  $q(\mathbf{z}|\mathbf{x}, \phi)$  (VAE encoder) be a base distribution for a flow model.

### Flow model in latent space

$$\log q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda) = \log q(\mathbf{z}|\mathbf{x}, \phi) + \log \left| \det \left( \frac{\partial g(\mathbf{z}, \lambda)}{\partial \mathbf{z}} \right) \right|$$

$$\mathbf{z}^* = g(\mathbf{z}, \lambda) = f^{-1}(\mathbf{z}, \lambda)$$

Here  $g(\mathbf{z}, \lambda)$  is a flow model (e.g. stack of planar/coupling/AR layers) parameterized by  $\lambda$ .

Let use  $q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda)$  as a variational distribution. Here  $\phi$  – encoder parameters,  $\lambda$  – flow parameters.

## Flows-based VAE posterior

- ▶ Encoder outputs base distribution  $q(\mathbf{z}|\mathbf{x}, \phi)$ .
- ▶ Flow model  $\mathbf{z}^* = g(\mathbf{z}, \lambda)$  transforms the base distribution  $q(\mathbf{z}|\mathbf{x}, \phi)$  to the distribution  $q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda)$ .
- ▶ Distribution  $q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda)$  is used as a variational distribution for ELBO maximization.

### Flow model in latent space

$$\log q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda) = \log q(\mathbf{z}|\mathbf{x}, \phi) + \log \left| \det \left( \frac{\partial g(\mathbf{z}, \lambda)}{\partial \mathbf{z}} \right) \right|$$

### ELBO with flow-based VAE posterior

$$\begin{aligned} \mathcal{L}(\phi, \theta, \lambda) &= \mathbb{E}_{q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda)} [\log p(\mathbf{x}, \mathbf{z}^*|\theta) - \log q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda)] \\ &= \mathbb{E}_{q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda)} \log p(\mathbf{x}|\mathbf{z}^*, \theta) - KL(q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda) || p(\mathbf{z}^*)). \end{aligned}$$

The second term in ELBO is reverse KL divergence. Planar flows was originally proposed for variational inference in VAE.

# Flows-based VAE posterior

## Flow model in latent space

$$\log q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda) = \log q(\mathbf{z}|\mathbf{x}, \phi) + \log \left| \det \left( \frac{\partial g(\mathbf{z}, \lambda)}{\partial \mathbf{z}} \right) \right|$$

## ELBO objective

$$\begin{aligned} \mathcal{L}(\phi, \theta, \lambda) &= \mathbb{E}_{q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda)} [\log p(\mathbf{x}, \mathbf{z}^*|\theta) - \log q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda)] = \\ &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} [\log p(\mathbf{x}, \mathbf{z}^*|\theta) - \log q(\mathbf{z}^*|\mathbf{x}, \phi, \lambda)] \Big|_{\mathbf{z}^*=g(\mathbf{z}, \lambda)} = \\ &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \left[ \log p(\mathbf{x}, g(\mathbf{z}, \lambda)|\theta) - \log q(\mathbf{z}|\mathbf{x}, \phi) - \log |\det(\mathbf{J}_g)| \right]. \end{aligned}$$

- ▶ Obtain samples  $\mathbf{z}$  from the encoder  $q(\mathbf{z}|\mathbf{x}, \phi)$ .
- ▶ Apply flow model  $\mathbf{z}^* = g(\mathbf{z}, \lambda)$ .
- ▶ Compute likelihood for  $\mathbf{z}^*$  using the decoder, base distribution for  $\mathbf{z}^*$  and the Jacobian.

# Inverse autoregressive flow (IAF)

$$\mathbf{x} = g(\mathbf{z}, \boldsymbol{\theta}) \quad \Rightarrow \quad x_i = \tilde{\sigma}_i(\mathbf{z}_{1:i-1}) \cdot z_i + \tilde{\mu}_i(\mathbf{z}_{1:i-1}).$$

$$\mathbf{z} = f(\mathbf{x}, \boldsymbol{\theta}) \quad \Rightarrow \quad z_i = (x_i - \tilde{\mu}_i(\mathbf{z}_{1:i-1})) \cdot \frac{1}{\tilde{\sigma}_i(\mathbf{z}_{1:i-1})}.$$

## Reverse KL for flow model



$$KL(p||\pi) = \mathbb{E}_{p(\mathbf{z})} \left[ \log p(\mathbf{z}) - \log \left| \det \left( \frac{\partial g(\mathbf{z}, \boldsymbol{\theta})}{\partial \mathbf{z}} \right) \right| - \log \pi(g(\mathbf{z}, \boldsymbol{\theta})) \right]$$

- ▶ We don't need to think about computing the function  $f(\mathbf{x}, \boldsymbol{\theta})$ .
- ▶ Inverse autoregressive flow is a natural choice for using flows in VAE:


$$\mathbf{z} = \boldsymbol{\sigma}(\mathbf{x}) \odot \boldsymbol{\epsilon} + \boldsymbol{\mu}(\mathbf{x}), \quad \boldsymbol{\epsilon} \sim \mathcal{N}(0, 1); \quad \sim q(\mathbf{z}|\mathbf{x}, \phi).$$

$$\mathbf{z}_k = \tilde{\boldsymbol{\sigma}}_k(\mathbf{z}_{k-1}) \odot \mathbf{z}_{k-1} + \tilde{\boldsymbol{\mu}}_k(\mathbf{z}_{k-1}), \quad k \geq 1; \quad \sim q_k(\mathbf{z}_k|\mathbf{x}, \phi, \{\boldsymbol{\lambda}_j\}_{j=1}^k).$$


# Inverse autoregressive flow (IAF)



(a) Prior distribution



(b) Posteriors in standard VAE



(c) Posteriors in VAE with IAF

# Flows-based VAE prior vs posterior

## Theorem

VAE with the flow-based prior for latent code  $\mathbf{z}$  is equivalent to VAE with flow-based posterior for latent code  $\mathbf{z}$ .

## Proof

$$\begin{aligned}\mathcal{L}(\phi, \theta, \lambda) &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \log p(\mathbf{x}|\mathbf{z}, \theta) - \underbrace{KL(q(\mathbf{z}|\mathbf{x}, \phi) || p(\mathbf{z}|\lambda))}_{\text{flow-based prior}} \\ &= \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \log p(\mathbf{x}|\mathbf{z}, \theta) - \underbrace{KL(q(\mathbf{z}|\mathbf{x}, \phi, \lambda) || p(\mathbf{z}))}_{\text{flow-based posterior}}\end{aligned}$$

(Here we use Flow KL duality theorem from Lecture 5)

## Flows in VAE posterior


$$\mathcal{L}(\phi, \theta, \lambda) = \mathbb{E}_{q(\mathbf{z}|\mathbf{x}, \phi)} \left[ \log p(\mathbf{x}, g(\mathbf{z}, \lambda) | \theta) - \log q(\mathbf{z}|\mathbf{x}, \phi) - \log |\det(\mathbf{J}_g)| \right].$$



# Flows-based VAE prior vs posterior

- ▶ IAF posterior decoder path:  $p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta})$ ,  $\mathbf{z} \sim p(\mathbf{z})$ .
- ▶ AF prior decoder path:  $p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta})$ ,  $\mathbf{z} = f(\mathbf{z}^*, \boldsymbol{\lambda})$ ,  $\epsilon \sim p(\mathbf{z}^*)$ .

The AF prior and the IAF posterior have the same computation cost, so using the AF prior makes the model more expressive at no training time cost.



# VAE limitations

- ▶ Poor generative distribution (decoder)

$$p(\mathbf{x}|\mathbf{z}, \boldsymbol{\theta}) = \mathcal{N}(\mathbf{x}|\boldsymbol{\mu}_{\boldsymbol{\theta}}(\mathbf{z}), \boldsymbol{\sigma}_{\boldsymbol{\theta}}^2(\mathbf{z})) \quad \text{or} \quad = \text{Softmax}(\boldsymbol{\pi}_{\boldsymbol{\theta}}(\mathbf{z})).$$

- ▶ Loose lower bound

$$\log p(\mathbf{x}|\boldsymbol{\theta}) - \mathcal{L}(q, \boldsymbol{\theta}) = (?).$$

- ▶ Poor prior distribution

$$p(\mathbf{z}) = \mathcal{N}(0, \mathbf{I}).$$

- ▶ Poor variational posterior distribution (encoder)

$$q(\mathbf{z}|\mathbf{x}, \boldsymbol{\phi}) = \mathcal{N}(\mathbf{z}|\boldsymbol{\mu}_{\boldsymbol{\phi}}(\mathbf{x}), \boldsymbol{\sigma}_{\boldsymbol{\phi}}^2(\mathbf{x})).$$

# Summary

- ▶ Dequantization allows to fit discrete data using continuous model.
- ▶ Uniform dequantization is the simplest form of dequantization. Variational dequantization is a more natural type that was proposed in Flow++ model.
- ▶ The ELBO surgery reveals insights about a prior distribution in VAE. The optimal prior is the aggregated posterior.
- ▶ We could use flow-based prior in VAE (moreover, autoregressive).
- ▶ We could use flows to make variational posterior more expressive. This is equivalent to the flow-based prior.