

ATOMIC BASES AND T-PATH FORMULA FOR CLUSTER ALGEBRAS OF TYPE D

Emily Gunawan – University of Minnesota, Minneapolis (joint work with Gregg Musiker)

We work with cluster algebras with principal coefficients

What is a cluster algebra? (Fomin and Zelevinsky, 2000)

- A (coefficient-free) *cluster algebra* of rank n is a \mathbb{Z} -subalgebra of $\mathbb{Q}(x_1, \dots, x_n)$ generated by elements called *cluster variables*:
 - Start with an *initial seed*: a cluster $\mathbf{x} = \{x_1, x_2, \dots, x_n\}$ and a skew-symmetrizable *exchange matrix* $B = (b_{ij})$.
 - For each $k = 1, \dots, n$, we *mutate* in the k -th direction $(\{x_1, \dots, x_k, \dots, x_n\}, B) \xrightarrow{\mu_k} (\{x_1, \dots, x'_k, \dots, x_n\}, \mu_k(B))$ to obtain a new seed where


$$x'_k = \frac{1}{x_k} \left(\prod_{b_{ik} > 0} x_i^{b_{ik}} + \prod_{b_{ik} < 0} x_i^{-b_{ik}} \right) \text{ and } \mu_k(B)_{ij} = \begin{cases} -b_{ij} & \text{if } k = i \text{ or } k = j \\ b_{ij} + \frac{1}{2} (|b_{ik}|b_{kj} + b_{ik}|b_{kj}|) & \text{otherwise} \end{cases}$$

- Apply all possible sequences of mutations to produce all cluster variables (usually infinitely many).
- Laurent Phenomenon:** each cluster variable can be expressed as a Laurent polynomial in $\{x_1, \dots, x_n\}$.
- Positivity:** this Laurent polynomial has positive coefficients (2014, [Lee and Schiffler], [Gross, Hacking, Keel, and Kontsevich], and special cases by others).

Cluster algebras from orientable surfaces


Definition: ordinary arcs

- An *ordinary arc* γ is a non-contractible curve between marked points such that γ does not cross itself or the boundary, and γ is not homotopic to a boundary edge.
- An *ideal triangulation* is a maximum collection of distinct arcs that pairwise do not cross.



Definition: tagged arcs


- A *tagged arc* is an ordinary arc (which does not cut out a monogon with 1 puncture $\ell \curvearrowright$) decorated (plain or with a \bowtie) at each endpoint.
- A *tagged triangulation* is a maximum collection of distinct tagged arcs that are pairwise “compatible”.



Theorem (Fomin, Shapiro, and Thurston, 2006)

One can define a cluster algebra from an orientable Riemann surface + interior points (called *punctures*) and/or marked points on the boundary:

seed $(\mathbf{x}_T, B_T) \longleftrightarrow$ tagged triangulation $T = \{\tau_1, \dots, \tau_n\}$
 cluster variable $x_\gamma \longleftrightarrow$ tagged arc γ
 cluster mutation \longleftrightarrow “flipping diagonals”



Result 1: T-path formula for cluster variables for punctured surfaces

We generalize (to surfaces with punctures) Schiffler and Thomas’ T -path definition and formula for unpunctured surfaces (2009). Let T° be an ideal triangulation and γ an ordinary arc that crosses T° . Let Δ_k denote the k -th ideal triangle crossed by γ .

Definition: quasi-arc

If τ is an ordinary arc, let an associated *quasi-arc* τ' be a curve (not passing through the puncture p) which agrees with τ outside of a small radius- ϵ disk D_ϵ around p .

Definition: T -path

A (complete) (T°, γ) -path $w = (w_1, \dots, w_{2d+1})$ is a concatenation of quasi-arcs and boundary edges such that:

(T1) Each even step w_{2k} ($k = 1, \dots, d$) is the k -th arc that γ crosses.

(T2) The path w is homotopic to γ , and satisfies the following:

Let p_1, \dots, p_d be the intersection points of γ and T° . Let γ_k be the segment along γ between p_k and p_{k+1} . Then the segment γ_k is homotopic to the segment from p_k following w_{2k} , following w_{2k+1} , following w_{2k+2} until p_{k+1} .

(T3) The step w_{2k+1} traverses a side of the triangle Δ_k , and starts and finishes in the interior of Δ_k or at a boundary marked point.

new

Theorem (T -path formula)

The cluster variable x_γ expressed in the variables corresponding to T° is

$$x_\gamma = \sum_w x(w)$$


over all (T°, γ) -paths $w = (w_1, \dots, w_{2d+1})$, where $x(w) := \left(\prod_{i \text{ odd}} x_{w_i} \right) \left(\prod_{i \text{ even}} x_{w_i}^{-1} \right)$.

In addition, we have:

- a similar formula for cluster algebras with principal and arbitrary coefficient systems
- a T -path formula for a tagged arc with \bowtie on its endpoint/s

Example: once-punctured 4-gon

A tagged triangulation of a once-punctured square \longleftrightarrow A quiver that is mutation equivalent to an orientation of a type D_4 Dynkin diagram.



This example corresponds to

$$B = \begin{pmatrix} 1 & 2 & r & r^{\bowtie} \\ 0 & -1 & 0 & 0 \\ 2 & 1 & 0 & 1 \\ 0 & -1 & 0 & 0 \\ r^{\bowtie} & 0 & -1 & 0 \end{pmatrix}$$

Example: once-punctured 3-gon




Fig. 1: There are $\binom{2n}{n} - \binom{2n-2}{n-1}$ tagged triangulations of a once-punctured n -gon.

Atomic bases

Definition: atomic bases (Sherman and Zelevinsky, 2003)

Let \mathcal{A} be a (coefficient-free) cluster algebra.

- Let the *positive cone* of \mathcal{A} be $\mathcal{A}^+ := \{\text{positive elements}\} = \{\text{elements that are positive Laurent polynomials with respect to every cluster}\}$.
- The subset \mathcal{B} of all indecomposable positive elements (*i.e.*, those that cannot be written as a sum of two positive elements) is called the *atomic basis* if it forms a \mathbb{Z} -basis of \mathcal{A} .

The existence of this atomic basis is not known in general. The cluster algebra with the exchange matrix $\begin{pmatrix} 0 & b \\ -c & 0 \end{pmatrix}$ has no atomic basis if $bc \geq 5$.

Result 2: Combinatorial proof (by T-path) for type D cluster algebras

Definition: cluster monomial

a *cluster monomial* is a product of cluster variables all coming from the same cluster, e.g. a^3be^2 is a cluster monomial if $\{a, b, c, d, e\}$ is a cluster.

A cluster monomial corresponds to a multi-tagged dissection (*i.e.*, a partial tagged triangulation allowing multiple copies of tagged arcs).

Theorem (atomic basis)


For a cluster algebra of type A , D , or E , the basis of cluster monomials is atomic.

- Representation theory proof by [Cerulli Irelli, 2011] and [Cerulli Irelli, Keller, Labardini-Fragoso, and Plamondon, 2012].
- We give a combinatorial proof (using the T -path formula) for type D , inspired by work on types A and \tilde{A} by [Dupont and Thomas, 2011].

Atomic bases for other surfaces

Conjecture (Fomin, Shapiro, and Thurston, 2008, unpublished, based on Fock and Goncharov, 2006)



A candidate for atomic bases: the “*bracelets collection*” consisting of all cluster monomials + a class of elements. A *bracelet* is a closed loop in the interior of the surface which wraps around itself once or multiple times and avoids marked points.



- True for annuli, type \tilde{A} (Dupont and Thomas, 2011).
- The bracelets collection forms a basis for unpunctured surfaces (Musiker, Schiffler, and Williams, 2011).

Current work: type affine D

Type \tilde{D}_{n-1} cluster algebras ($(n-3)$ -gons with 2 punctures), *e.g.* type \tilde{D}_6 cluster algebra comes from a twice-punctured disk with 4 marked points on the boundary.



- We thank the referees, P. Pylyavskyy, V. Reiner, H. Thomas, P. Webb, and NSF Grants DMS-1067183 and DMS-1148634
- Email: egunawan@umn.edu
- Home page: umn.edu/home/egunawan