

PSO - Stands for Particle Swarm Optimization

The movement of the particles is given by

$$\text{Velocity} \quad v_{t+1} = w_t v_t + c_1 r_1 (p_t^b - x_t) + c_2 r_2 (p^g - x_t) \quad \text{--- (1)}$$

$$\text{Position} \quad x_{t+1} = x_t + v_{t+1} \quad \text{--- (2)}$$

t : iteration number ; w_t : inertia weight ✓
 r_1, r_2 : random number between 0 & 1; c_1, c_2 : correction factor ✓
 p_t^b : pbest position at t iteration; p^g : gbest position ✓
 v_t, x_t : velocity and position at t iteration

Maximize $f(X) = x_1^2 - x_1 x_2 + x_2^2 + 2x_1 + 4x_2 + 3$

where $-5 \leq x_1, x_2 \leq 5$

PSO Parameter Setting (used only for illustration)

Randomly chosen ✓

Population Size = 5 ✓

$$c_1 = c_2 = 1.5$$

Max. Iteration = 20

Dimension of the problem = 2 ✓

Inertia weight (w) = 0.9

Goal is to update these values ALWAYS

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \end{bmatrix}$	$\begin{bmatrix} \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \end{bmatrix}$	$\begin{bmatrix} \text{.....} \\ \text{.....} \\ \text{.....} \\ \text{.....} \end{bmatrix}$	$\begin{bmatrix} \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \end{bmatrix}$
$p^g(\text{gbest}) = [\text{.....} \quad \text{.....}]$			

Iteration 1:

Randomly chosen Velocity
between 0 & 1



Goal is to update these values ALWAYS

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \end{bmatrix}$	$\begin{bmatrix} \text{.....} \\ \text{.....} \\ \text{.....} \\ \text{.....} \\ \text{.....} \end{bmatrix}$	$\begin{bmatrix} \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \end{bmatrix}$
$p^g(\text{gbest}) = [\text{.....} \quad \text{.....}]$			

Use $x = L + \text{rand.} * (U - L)$

Lower rand. upper

Iteration 1:

Randomly chosen Velocity
between 0 & 1

Randomly Initialize Position
between -5 & 5

Goal is to update these values ALWAYS

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} \text{.....} \\ \text{.....} \\ \text{.....} \\ \text{.....} \\ \text{.....} \end{bmatrix}$	$\begin{bmatrix} \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \\ \text{.....} & \text{.....} \end{bmatrix}$
$p^g(\text{gbest}) = [\text{.....} \quad \text{.....}]$			

Use $x = L + \text{rand.} * (U - L)$

Iteration 1:

- ✚ Randomly chosen Velocity between 0 & 1
- ✚ Randomly Initialize Position between -5 & 5
- ✚ Calculate fitness values $f(x)$:

Goal is to update these values ALWAYS

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
	x_1 x_2		
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \end{bmatrix}$

$$p^g(\text{gbest}) = [\text{-----} \quad \text{-----}]$$

$$\text{Maximize } f(X) = x_1^2 - x_1x_2 + x_2^2 + 2x_1 + 4x_2 + 3$$

Iteration 1:

- ✚ Randomly chosen Velocity between 0 & 1
- ✚ Randomly Initialize Position between -5 & 5
- ✚ Calculate fitness values $f(x)$:
- ✚ Since there is NO previous iteration exists, so pbest is $p^b = x$

Goal is to update these values ALWAYS

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
	x_1 x_2		
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$

$$p^g(\text{gbest}) = [\text{-----} \quad \text{-----}]$$

$$\text{Maximize } f(X) = x_1^2 - x_1x_2 + x_2^2 + 2x_1 + 4x_2 + 3$$

Iteration 1:

- ✚ Randomly chosen Velocity between 0 & 1
- ✚ Randomly Initialize Position between -5 & 5
- ✚ Calculate fitness values $f(x)$:
- ✚ Since there is NO previous iteration exists, so pbest is $p^b = x$
- ✚ Calculate gbest p^g
Maximum fitness value = 48.6753

Goal is to update these values ALWAYS

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
	x_1 x_2		
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$

$$p^g(\text{gbest}) = [\text{-----} \quad \text{-----}]$$

$$\text{Maximize } f(X) = x_1^2 - x_1x_2 + x_2^2 + 2x_1 + 4x_2 + 3$$

Iteration 1:

✚ Randomly chosen Velocity
between 0 & 1

✚ Randomly Initialize Position
between -5 & 5

✚ Calculate fitness values $f(x)$:

✚ Since there is NO previous iteration
exists, so pbest is $p^b = x$

✚ Calculate gbest p^g

Maximum fitness value = 48.6753

Goal is to update these values ALWAYS

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$

$$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$$

$$\text{Maximize } f(X) = x_1^2 - x_1 x_2 + x_2^2 + 2x_1 + 4x_2 + 3$$

Iteration 2:

Iteration 1:

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$

$$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$$

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots \\ \dots \\ \dots \\ \dots \\ \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$

$$p^g(\text{gbest}) = [\dots \quad \dots]$$

Iteration 2:

Velocity update:

For 1st particle, 1st component

$$v_{11} = wv_{11} + c_1 r_1 (p_{11}^b - x_{11}) + c_2 r_2 (p^g - x_{11})$$

Iteration 1:

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$

$$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$$

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots \\ \dots \\ \dots \\ \dots \\ \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$

$$p^g(\text{gbest}) = [\dots \quad \dots]$$

Iteration 2:

Velocity update:

For 1st particle, 1st component

$$\begin{aligned}
 v_{11} &= wv_{11} + c_1r_1(p_{11}^b - x_{11}) \\
 &\quad + c_2r_2(p^g - x_{11}) \\
 &= 0.9 \times 0.2194 \\
 &\quad + (1.5)(0.5949)(3.1472 - 3.1472) \\
 &\quad + (1.5)(0.0855)(4.1338 - 3.1472) \\
 &= 0.3240
 \end{aligned}$$

Iteration 1:

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$
p^g (gbest) = [4.1338 4.5751]			

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.3240 & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots \\ \dots \\ \dots \\ \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$
p^g (gbest) = [-----]			

Iteration 2:

Velocity update:

For 1st particle, 1st component

$$\begin{aligned}
 v_{11} &= wv_{11} + c_1r_1(p_{11}^b - x_{11}) \\
 &\quad + c_2r_2(p^g - x_{11}) \\
 &= 0.9 \times 0.2194 \\
 &\quad + (1.5)(0.5949)(3.1472 - 3.1472) \\
 &\quad + (1.5)(0.0855)(4.1338 - 3.1472) \\
 &= 0.3240
 \end{aligned}$$

Position update:

$$\begin{aligned}
 x_{11} &= v_{11} + x_{11} \\
 &= 3.4712 \in (-5, 5)
 \end{aligned}$$

$$= 0.3240 + 3.1472$$

Iteration 1:

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$
p^g (gbest) = [4.1338 4.5751]			

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.3240 & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots \\ \dots \\ \dots \\ \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$
p^g (gbest) = [-----]			

Iteration 2:

Velocity update:

For 1st particle, 1st component

$$\begin{aligned}
 v_{11} &= wv_{11} + c_1r_1(p_{11}^b - x_{11}) \\
 &\quad + c_2r_2(p^g - x_{11}) \\
 &= 0.9 \times 0.2194 \\
 &\quad + (1.5)(0.5949)(3.1472 - 3.1472) \\
 &\quad + (1.5)(0.0855)(4.1338 - 3.1472) \\
 &= 0.3240
 \end{aligned}$$

Position update:

$$\begin{aligned}
 x_{11} &= v_{11} + x_{11} \\
 &= 3.4712 \in (-5, 5)
 \end{aligned}$$

Iteration 1:

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$
p^g (gbest) = [4.1338 4.5751]			

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.3240 & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} 3.4712 & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots \\ \dots \\ \dots \\ \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$
p^g (gbest) = [-----]			

Iteration 2:

Velocity update:

1st particle; 2nd component:

$$\begin{aligned}
 v_{12} &= wv_{12} + c_1 r_1 (p_{12}^b - x_{12}) \\
 &\quad + c_2 r_2 (p^g - x_{12}) \\
 &= (0.9)(0.2449) \\
 &\quad + (1.5)(0.5949)(-4.0246 + 4.0246) \\
 &\quad + (1.5)(0.0855)(4.5751 + 4.0246) \\
 &= 1.3233
 \end{aligned}$$

Iteration 1:

v (velocity)	x (position)	f(x)	p _i ^b (pbest)
0.2194 0.2449	x ₁ x ₂		
0.1908 0.2228	3.1472 -4.0246	31.9645	3.1472 -4.0246
0.3828 0.3232	4.0579 -2.2150	32.6168	4.0579 -2.2150
0.3976 0.3547	-3.7301 0.4688	13.2071	-3.7301 0.4688
0.0934 0.3773	4.1338 4.5751	48.6753	4.1338 4.5751
	1.3236 4.6489	41.4537	1.3236 4.6489

$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$

v (velocity)	x (position)	f(x)	p _i ^b (pbest)
0.3240	3.4712		
.....
.....
.....
.....
.....

$p^g(\text{gbest}) = [..... \quad]$

Iteration 2:

Velocity update:

1st particle; 2nd component:

$$\begin{aligned}
 v_{12} &= wv_{12} + c_1 r_1 (p_{12}^b - x_{12}) \\
 &\quad + c_2 r_2 (p^g - x_{12}) \\
 &= (0.9)(0.2449) \\
 &\quad + (1.5)(0.5949)(-4.0246 + 4.0246) \\
 &\quad + (1.5)(0.0855)(4.5751 + 4.0246) \\
 &= 1.3233
 \end{aligned}$$

Position update:

$$\begin{aligned}
 x_{12} &= v_{12} + x_{12} \\
 &= -2.7013 \in (-5, 5)
 \end{aligned}$$

Iteration 1:

v (velocity)	x (position)	f(x)	p _i ^b (pbest)
0.2194 0.2449	x ₁ x ₂		
0.1908 0.2228	3.1472 -4.0246	31.9645	3.1472 -4.0246
0.3828 0.3232	4.0579 -2.2150	32.6168	4.0579 -2.2150
0.3976 0.3547	-3.7301 0.4688	13.2071	-3.7301 0.4688
0.0934 0.3773	4.1338 4.5751	48.6753	4.1338 4.5751
	1.3236 4.6489	41.4537	1.3236 4.6489

$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$

v (velocity)	x (position)	f(x)	p _i ^b (pbest)
0.3240 1.3233	3.4712		
.....
.....
.....
.....
.....

$p^g(\text{gbest}) = [..... \quad]$

Iteration 2:

Velocity update:

1st particle; 2nd component:

$$\begin{aligned}
 v_{12} &= wv_{12} + c_1 r_1 (p_{12}^b - x_{12}) \\
 &\quad + c_2 r_2 (p^g - x_{12}) \\
 &= (0.9)(0.2449) \\
 &\quad + (1.5)(0.5949)(-4.0246 + 4.0246) \\
 &\quad + (1.5)(0.0855)(4.5751 + 4.0246) \\
 &= 1.3233
 \end{aligned}$$

Position update:

$$\begin{aligned}
 x_{12} &= v_{12} + x_{12} \\
 &= -2.7013 \in (-5, 5)
 \end{aligned}$$

Iteration 1:

v (velocity)	x (position)	f(x)	p _i ^b (pbest)
0.2194 0.2449	x ₁ x ₂		
0.1908 0.2228	3.1472 -4.0246	31.9645	3.1472 -4.0246
0.3828 0.3232	4.0579 -2.2150	32.6168	4.0579 -2.2150
0.3976 0.3547	-3.7301 0.4688	13.2071	-3.7301 0.4688
0.0934 0.3773	4.1338 4.5751	48.6753	4.1338 4.5751
	1.3236 4.6489	41.4537	1.3236 4.6489

$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$

v (velocity)	x (position)	f(x)	p _i ^b (pbest)
0.3240 1.3233	3.4712 -2.7013		
.....
.....
.....
.....
.....

$p^g(\text{gbest}) = [..... \quad]$

Iteration 2:

Velocity update:

For 2nd particle; 1st component

$$v_{21} = wv_{21} + c_1r_1(p_{21}^b - x_{21}) + c_2r_2(p^g - x_{21})$$

$$= 0.1815$$

$$x_{21} = v_{21} + x_{21}$$

$$= 4.2394 \in (-5, 5)$$

Iteration 1:

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
	x_1		
	x_2		
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$
$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$			

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots \\ \dots \\ \dots \\ \dots \\ \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$
$p^g(\text{gbest}) = [\dots \quad \dots]$			

Iteration 2:

Velocity update:

For 2nd particle; 1st component

$$v_{21} = wv_{21} + c_1r_1(p_{21}^b - x_{21}) + c_2r_2(p^g - x_{21})$$

$$= 0.1815$$

$$x_{21} = v_{21} + x_{21}$$

$$= 4.2394 \in (-5, 5)$$

Similarly, for 2nd particle; 2nd component,
we can get $v_{22} = 1.0714$;
 $x_{22} = -1.1436$

Iteration 1:

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
	x_1		
	x_2		
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$
$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$			

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$	$\begin{bmatrix} \dots \\ \dots \\ \dots \\ \dots \\ \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$
$p^g(\text{gbest}) = [\dots \quad \dots]$			

Iteration 2:

Similarly, we can found for other pairs the

Velocity & Position updates.

Iteration 1:

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
	x_1		
	x_2		
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$
$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$			

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3790 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} \dots \\ \dots \\ \dots \\ \dots \\ \dots \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$
$p^g(\text{gbest}) = [\dots \quad \dots]$			

Iteration 2:

Similarly, we can find for other pairs the

Velocity & Position updates.

Calculate the fitness values:

$$f_i(X) = x_1^2 - x_1 x_2 + x_2^2 + 2x_1 + 4x_2 + 3$$

Iteration 1:

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$
$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$			

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} 27.8600 \\ 31.0325 \\ 13.7537 \\ 53.7063 \\ 45.5655 \end{bmatrix}$	$\begin{bmatrix} \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \end{bmatrix}$
$p^g(\text{gbest}) = [\text{-----} \quad \text{-----}]$			

Iteration 2:

Similarly, we can find for other pairs the

Velocity & Position updates.

Calculate the fitness values:

Calculate gbest (p^g):

Iteration 1:

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2071 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$
$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$			

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} 27.8600 \\ 31.0325 \\ 13.7537 \\ 53.7063 \\ 45.5655 \end{bmatrix}$	$\begin{bmatrix} \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \end{bmatrix}$
$p^g(\text{gbest}) = [\text{-----} \quad \text{-----}]$			

Iteration 2:

Similarly, we can find for other pairs the

Velocity & Position updates.

Calculate the fitness values:

Calculate gbest (p^g):

Since $53.7063 > 48.6753$ & its position is 4th.

So updated gbest is $[4.4916 \quad 4.8943]$

Iteration 1:

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} x_1 & x_2 \\ 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2971 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$
$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$			

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} 27.8600 \\ 31.0325 \\ 13.7537 \\ 53.7063 \\ 45.5655 \end{bmatrix}$	$\begin{bmatrix} \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \end{bmatrix}$
$p^g(\text{gbest}) = [4.4916 \quad 4.8943]$			

Iteration 2:

Similarly, we can find for other pairs the

Velocity & Position updates.

Calculate the fitness values:

Calculate gbest (p^g):

Since $53.7063 > 48.6753$ & its position is 4th.

So updated gbest is [4.4916 4.8943]

Calculate pbest (p^b):

For 1st particle:

Since $27.8600 < 31.9645$, so pbest position remains same

For 3rd particle:

Since $13.7537 < 13.2971$, so updated pbest position

Iteration 1:

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
	x_1 x_2		
$\begin{bmatrix} 0.2194 & 0.2449 \\ 0.1908 & 0.2228 \\ 0.3828 & 0.3232 \\ 0.3976 & 0.3547 \\ 0.0934 & 0.3773 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$	$\begin{bmatrix} 31.9645 \\ 32.6168 \\ 13.2971 \\ 48.6753 \\ 41.4537 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -3.7301 & 0.4688 \\ 4.1338 & 4.5751 \\ 1.3236 & 4.6489 \end{bmatrix}$

$$p^g(\text{gbest}) = [4.1338 \quad 4.5751]$$

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} 27.8600 \\ 31.0325 \\ 13.7537 \\ 53.7063 \\ 45.5655 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$

$$p^g(\text{gbest}) = [4.4916 \quad 4.8943]$$

Iteration 3:

Update the velocity & position:

$$v_{11} = w v_{11} + c_1 r_1 (p_{11}^b - x_{11}) + c_2 r_2 (p^g - x_{11})$$

And $x_{11} = v_{11} + x_{11}$

Iteration 2: results

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} 27.8600 \\ 31.0325 \\ 13.7537 \\ 53.7063 \\ 45.5655 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$
$p^g(\text{gbest}) = [4.4916 \quad 4.8943]$			

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.1334 & 0.9843 \\ 0.0337 & 0.7826 \\ 2.0987 & 1.1985 \\ 0.3221 & 0.2873 \\ 0.7493 & 0.2862 \end{bmatrix}$	$\begin{bmatrix} 3.6045 & -1.7170 \\ 4.2731 & -0.3611 \\ -0.2784 & 2.4848 \\ 4.8137 & 5.0000 \\ 2.5174 & 5.0000 \end{bmatrix}$	$\begin{bmatrix} \text{-----} \\ \text{-----} \\ \text{-----} \\ \text{-----} \\ \text{-----} \end{bmatrix}$	$\begin{bmatrix} \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \end{bmatrix}$
$p^g(\text{gbest}) = [\text{-----} \quad \text{-----}]$			

Iteration 3:

Update the velocity & position:

Calculate the fitness value:

$$f(X) = x_1^2 - x_1 x_2 + x_2^2 + 2x_1 + 4x_2 + 3$$

Iteration 2: results

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} 27.8600 \\ 31.0325 \\ 13.7537 \\ 53.7063 \\ 45.5655 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$
$p^g(\text{gbest}) = [4.4916 \quad 4.8943]$			

$v(\text{velocity})$	$x(\text{position})$	$f(x)$	$p_i^b(\text{pbest})$
$\begin{bmatrix} 0.1334 & 0.9843 \\ 0.0337 & 0.7826 \\ 2.0987 & 1.1985 \\ 0.3221 & 0.2873 \\ 0.7493 & 0.2862 \end{bmatrix}$	$\begin{bmatrix} 3.6045 & -1.7170 \\ 4.2731 & -0.3611 \\ -0.2784 & 2.4848 \\ 4.8137 & 5.0000 \\ 2.5174 & 5.0000 \end{bmatrix}$	$\begin{bmatrix} 25.4710 \\ 30.0346 \\ 19.3259 \\ 56.7306 \\ 46.7851 \end{bmatrix}$	$\begin{bmatrix} \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \\ \text{-----} & \text{-----} \end{bmatrix}$
$p^g(\text{gbest}) = [\text{-----} \quad \text{-----}]$			

Iteration 3:

Update the velocity & position:

Calculate the fitness value:

Iteration 2: results

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} 27.8600 \\ 31.0325 \\ 13.7537 \\ 53.7063 \\ 45.5655 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$

$$p^g(\text{gbest}) = [4.4916 \quad 4.8943]$$

Update gbest (p^g):

Since $56.7306 > 53.7063$, so updated gbest is
gbest = [4.8137 5.0000]

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.1334 & 0.9843 \\ 0.0337 & 0.7826 \\ 2.0987 & 1.1985 \\ 0.3221 & 0.2873 \\ 0.7493 & 0.2862 \end{bmatrix}$	$\begin{bmatrix} 3.6045 & -1.7170 \\ 4.2731 & -0.3611 \\ -0.2784 & 2.4848 \\ 4.8137 & 5.0000 \\ 2.5174 & 5.0000 \end{bmatrix}$	$\begin{bmatrix} 25.4710 \\ 30.0346 \\ 19.3259 \\ 56.7306 \\ 46.7851 \end{bmatrix}$	$\begin{bmatrix} \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \\ \dots & \dots \end{bmatrix}$

$$p^g(\text{gbest}) = [4.8137 \quad 5.0000]$$

Iteration 3:

Update the velocity & position:

Calculate the fitness value:

Update gbest (p^g):

Since $56.7306 > 53.7063$, so updated gbest is
gbest = [4.8137 5.0000]

Update pbest (p^b):

Iteration 2: results

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} 27.8600 \\ 31.0325 \\ 13.7537 \\ 53.7063 \\ 45.5655 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$

$$p^g(\text{gbest}) = [4.4916 \quad 4.8943]$$

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.1334 & 0.9843 \\ 0.0337 & 0.7826 \\ 2.0987 & 1.1985 \\ 0.3221 & 0.2873 \\ 0.7493 & 0.2862 \end{bmatrix}$	$\begin{bmatrix} 3.6045 & -1.7170 \\ 4.2731 & -0.3611 \\ -0.2784 & 2.4848 \\ 4.8137 & 5.0000 \\ 2.5174 & 5.0000 \end{bmatrix}$	$\begin{bmatrix} 25.4710 \\ 30.0346 \\ 19.3259 \\ 56.7306 \\ 46.7851 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -0.2784 & 2.4848 \\ 4.8137 & 5.0000 \\ 2.5174 & 5.0000 \end{bmatrix}$

$$p^g(\text{gbest}) = [4.8137 \quad 5.0000]$$

Iteration 4:

Update the velocity & position:

Calculate the fitness value:

Update gbest (p^g):

Since 58.000 > 56.7306

gbest = [5.0000 5.0000]

Update pbest (p^b):

Iteration 3: results

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
0.1334 0.9843	3.6045 -1.7170	25.4710	3.4712 -2.7013
0.0337 0.7826	4.2731 -0.3611	30.0346	4.2394 -1.1436
2.0987 1.1985	-0.2784 2.4848	19.3259	-0.2784 2.4848
0.3221 0.2873	4.8137 5.0000	56.7306	4.8137 5.0000
0.7493 0.2862	2.5174 5.0000	46.7851	2.5174 5.0000

p^g (gbest) = [4.8137 5.0000]

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
0.1561 0.8690	3.7606 -0.8480	25.1798	3.6045 -1.7170
0.0696 0.6935	4.3427 0.3325	30.5409	4.3427 0.3325
2.5419 1.4012	2.2635 3.8860	34.4996	2.2635 3.8860
0.2899 0.2586	5.0000 5.0000	58.0000	5.0000 5.0000
0.9689 0.2576	3.4863 5.0000	49.6953	3.4863 5.0000

p^g (gbest) = [5.0000 5.0000]

Iteration 5:

Update the velocity & position:

Calculate the fitness value:

Update gbest (p^g):

Update pbest (p^b):

Since there is no change in the fitness value (58.000) and hence optimum will reach.

$$x_1 = 5, x_2 = 5$$

$$f(x_1, x_2) = 58$$

Iteration 4: results

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
0.1561 0.8690	3.7606 -0.8480	25.1798	3.6045 -1.7170
0.0696 0.6935	4.3427 0.3325	30.5409	4.3427 0.3325
2.5419 1.4012	2.2635 3.8860	34.4996	2.2635 3.8860
0.2899 0.2586	5.0000 5.0000	58.0000	5.0000 5.0000
0.9689 0.2576	3.4863 5.0000	49.6953	3.4863 5.0000

p^g (gbest) = [5.0000 5.0000]

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
0.1601 0.7567	3.9208 -0.0914	26.2152	3.9208 -0.0914
0.1469 1.2228	4.4896 1.5553	33.7933	4.4896 1.5553
2.6386 1.4040	4.9021 5.0000	57.3244	4.9021 5.0000
0.2609 0.2327	5.0000 5.0000	58.0000	5.0000 5.0000
1.0661 0.2319	4.5524 5.0000	55.0672	4.5524 5.0000

p^g (gbest) = [5.0000 5.0000]

x_1 x_2

Iteration 3:

Update the velocity & position:

Calculate the fitness value:

Update gbest (p^g):

Since $56.7306 > 53.7063$, so updated gbest is
 $G_{best} = [4.8137 \ 5.0000]$

Update pbest (p^b):

Iteration 2: results

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.3240 & 1.3233 \\ 0.1815 & 1.0714 \\ 1.3531 & 0.8175 \\ 0.3578 & 0.3192 \\ 0.4445 & 0.3301 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$	$\begin{bmatrix} 27.8600 \\ 31.0325 \\ 13.7537 \\ 53.7063 \\ 45.5655 \end{bmatrix}$	$\begin{bmatrix} 3.1472 & -4.0246 \\ 4.0579 & -2.2150 \\ 2.3770 & 1.2863 \\ 4.4916 & 4.8943 \\ 1.7681 & 4.9790 \end{bmatrix}$

$$p^g(gbest) = [4.4916 \ 4.8943]$$

v (velocity)	x (position)	$f(x)$	p_i^b (pbest)
$\begin{bmatrix} 0.1334 & 0.9843 \\ 0.0337 & 0.7826 \\ 2.0987 & 1.1985 \\ 0.3221 & 0.2873 \\ 0.7493 & 0.2862 \end{bmatrix}$	$\begin{bmatrix} 3.6045 & -1.7170 \\ 4.2731 & -0.3611 \\ -0.2784 & 2.4848 \\ 4.8137 & 5.0000 \\ 2.5174 & 5.0000 \end{bmatrix}$	$\begin{bmatrix} 25.4710 \\ 30.0346 \\ 19.3259 \\ 56.7306 \\ 46.7851 \end{bmatrix}$	$\begin{bmatrix} 3.4712 & -2.7013 \\ 4.2394 & -1.1436 \\ -0.2784 & 2.4848 \\ 4.8137 & 5.0000 \\ 2.5174 & 5.0000 \end{bmatrix}$

$$p^g(gbest) = [4.8137 \ 5.0000]$$

$$|f(x) - f(x_{pre})| < \epsilon$$

Summary:

Update the velocity & position:

$$v_{t+1} = w_t v_t + c_1 r_1 (p_t^b - x_t) + c_2 r_2 (p^g - x_t)$$

$$x_{t+1} = x_t + v_{t+1}$$

Calculate the fitness value:

Max / min

Calculate the objective function values $f(x)$

Update gbest (p^g):

Compare the fitness value

Compare the individual particle fitness value & select the best one.

Update pbest (p^b):