

Observation of a Charmed Baryon Decaying to $D^0 p$ at a Mass Near $2.94 \text{ GeV}/c^2$

B. Aubert,¹ R. Barate,¹ M. Bona,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹
V. Tisserand,¹ A. Zghiche,¹ E. Grauges,² A. Palano,³ M. Pappagallo,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴
Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ M. Battaglia,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶
R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ J. A. Kadyk,⁶ L. T. Kerth,⁶
Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶
N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷
S. E. Morgan,⁷ A. T. Watson,⁷ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸
T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ W. N. Cottingham,⁹ D. Walker,⁹ T. Cuhadar-Donszelmann,¹⁰
B. G. Fulson,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹
M. Saleem,¹¹ L. Teodorescu,¹¹ V. E. Blinov,¹² A. D. Bukin,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² A. P. Onuchin,¹²
S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² K. Yu. Todyshev,¹² D. S. Best,¹³ M. Bondioli,¹³
M. Bruinsma,¹³ M. Chao,¹³ S. Curry,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ P. Lund,¹³ M. Mandelkern,¹³
R. K. Mommensen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ S. Abachi,¹⁴ C. Buchanan,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵
O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ L. Zhang,¹⁵ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ H. P. Paar,¹⁶ S. Rahatlou,¹⁶
V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ D. Kovalskyi,¹⁷
J. D. Richman,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸
G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸
J. Albert,¹⁹ E. Chen,¹⁹ A. Dvoretzskii,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹
A. Samuel,¹⁹ R. Andreassen,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. C. Bloom,²¹
S. Chen,²¹ W. T. Ford,²¹ J. F. Hirschauer,²¹ A. Kreisel,²¹ U. Nauenberg,²¹ A. Olivas,²¹ W. O. Ruddick,²¹
J. G. Smith,²¹ K. A. Ulmer,²¹ S. R. Wagner,²¹ J. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² A. Soffer,²² W. H. Toki,²²
R. J. Wilson,²² F. Winklmeier,²² Q. Zeng,²² D. D. Altenburg,²³ E. Feltresi,²³ A. Hauke,²³ H. Jasper,²³ B. Spaan,²³
T. Brandt,²⁴ V. Klose,²⁴ H. M. Lacker,²⁴ W. F. Mader,²⁴ R. Nogowski,²⁴ A. Petzold,²⁴ J. Schubert,²⁴
K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ A. Volk,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,²⁵ *
E. Latour,²⁵ Ch. Thiebaux,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ W. Gradl,²⁶ F. Muheim,²⁶ S. Playfer,²⁶
A. I. Robertson,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷
M. Negrini,²⁷ A. Petrella,²⁷ L. Piemontese,²⁷ E. Prencipe,²⁷ F. Anulli,²⁸ R. Baldini-Ferrolì,²⁸ A. Calcaterra,²⁸
R. de Sangro,²⁸ G. Finocchiaro,²⁸ S. Pacetti,²⁸ P. Patteri,²⁸ I. M. Peruzzi,²⁸ † M. Piccolo,²⁸ M. Rama,²⁸
A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ M. Lo Vetere,²⁹ M. M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹
C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰
J. Wu,³⁰ R. S. Dubitzky,³¹ J. Marks,³¹ S. Schenk,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³²
U. Egede,³² R. L. Flack,³² J. R. Gaillard,³² J. A. Nash,³² M. B. Nikolich,³² W. Panduro Vazquez,³² X. Chai,³³
M. J. Charles,³³ U. Mallik,³³ N. T. Meyer,³³ V. Ziegler,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ L. Dong,³⁴ V. Eyges,³⁴
W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ A. V. Gritsan,³⁵ M. Fritsch,³⁶ G. Schott,³⁶
N. Arnaud,³⁷ M. Davier,³⁷ G. Grosdidier,³⁷ A. Höcker,³⁷ F. Le Diberder,³⁷ V. Lepeltier,³⁷ A. M. Lutz,³⁷
A. Oyanguren,³⁷ S. Pruvot,³⁷ S. Rodier,³⁷ P. Roudeau,³⁷ M. H. Schune,³⁷ A. Stocchi,³⁷ W. F. Wang,³⁷
G. Wormser,³⁷ C. H. Cheng,³⁸ D. J. Lange,³⁸ D. M. Wright,³⁸ C. A. Chavez,³⁹ I. J. Forster,³⁹ J. R. Fry,³⁹
E. Gabathuler,³⁹ R. Gamet,³⁹ K. A. George,³⁹ D. E. Hutchcroft,³⁹ D. J. Payne,³⁹ K. C. Schofield,³⁹
C. Touramanis,³⁹ A. J. Bevan,⁴⁰ F. Di Lodovico,⁴⁰ W. Menges,⁴⁰ R. Sacco,⁴⁰ C. L. Brown,⁴¹ G. Cowan,⁴¹
H. U. Fläeher,⁴¹ D. A. Hopkins,⁴¹ P. S. Jackson,⁴¹ T. R. McMahon,⁴¹ S. Ricciardi,⁴¹ F. Salvatore,⁴¹ D. N. Brown,⁴²
C. L. Davis,⁴² J. Allison,⁴³ N. R. Barlow,⁴³ R. J. Barlow,⁴³ Y. M. Chia,⁴³ C. L. Edgar,⁴³ M. P. Kelly,⁴³
G. D. Lafferty,⁴³ M. T. Naisbit,⁴³ J. C. Williams,⁴³ J. I. Yi,⁴³ C. Chen,⁴⁴ W. D. Hulsbergen,⁴⁴ A. Jawahery,⁴⁴
C. K. Lae,⁴⁴ D. A. Roberts,⁴⁴ G. Simi,⁴⁴ G. Blaylock,⁴⁵ C. Dallapiccola,⁴⁵ S. S. Hertzbach,⁴⁵ X. Li,⁴⁵ T. B. Moore,⁴⁵
S. Saremi,⁴⁵ H. Staengle,⁴⁵ S. Y. Willocq,⁴⁵ R. Cowan,⁴⁶ K. Koenke,⁴⁶ G. Sciolla,⁴⁶ S. J. Sekula,⁴⁶ M. Spitznagel,⁴⁶
F. Taylor,⁴⁶ R. K. Yamamoto,⁴⁶ H. Kim,⁴⁷ P. M. Patel,⁴⁷ C. T. Potter,⁴⁷ S. H. Robertson,⁴⁷ A. Lazzaro,⁴⁸
V. Lombardo,⁴⁸ F. Palombo,⁴⁸ J. M. Bauer,⁴⁹ L. Cremaldi,⁴⁹ V. Eschenburg,⁴⁹ R. Godang,⁴⁹ R. Kroeger,⁴⁹
J. Reidy,⁴⁹ D. A. Sanders,⁴⁹ D. J. Summers,⁴⁹ H. W. Zhao,⁴⁹ S. Brunet,⁵⁰ D. Côté,⁵⁰ M. Simard,⁵⁰ P. Taras,⁵⁰
F. B. Viaud,⁵⁰ H. Nicholson,⁵¹ N. Cavallo,⁵² ‡ G. De Nardo,⁵² D. del Re,⁵² F. Fabozzi,⁵² ‡ C. Gatto,⁵² L. Lista,⁵²
D. Monorchio,⁵² P. Paolucci,⁵² D. Piccolo,⁵² C. Sciacca,⁵² M. Baak,⁵³ H. Bulten,⁵³ G. Raven,⁵³ H. L. Snoek,⁵³
C. P. Jessop,⁵⁴ J. M. LoSecco,⁵⁴ T. Allmendinger,⁵⁵ G. Benelli,⁵⁵ K. K. Gan,⁵⁵ K. Honscheid,⁵⁵ D. Hufnagel,⁵⁵

P. D. Jackson,⁵⁵ H. Kagan,⁵⁵ R. Kass,⁵⁵ T. Pulliam,⁵⁵ A. M. Rahimi,⁵⁵ R. Ter-Antonyan,⁵⁵ Q. K. Wong,⁵⁵
 N. L. Blount,⁵⁶ J. Brau,⁵⁶ R. Frey,⁵⁶ O. Igonkina,⁵⁶ M. Lu,⁵⁶ R. Rahmat,⁵⁶ N. B. Sinev,⁵⁶ D. Strom,⁵⁶
 J. Strube,⁵⁶ E. Torrence,⁵⁶ F. Galeazzi,⁵⁷ A. Gaz,⁵⁷ M. Margoni,⁵⁷ M. Morandin,⁵⁷ A. Pompili,⁵⁷ M. Posocco,⁵⁷
 M. Rotondo,⁵⁷ F. Simonetto,⁵⁷ R. Stroili,⁵⁷ C. Voci,⁵⁷ M. Benayoun,⁵⁸ J. Chauveau,⁵⁸ P. David,⁵⁸ L. Del Buono,⁵⁸
 Ch. de la Vaissière,⁵⁸ O. Hamon,⁵⁸ B. L. Hartfiel,⁵⁸ M. J. J. John,⁵⁸ Ph. Leruste,⁵⁸ J. Malcès,⁵⁸ J. Ocariz,⁵⁸
 L. Roos,⁵⁸ G. Therin,⁵⁸ P. K. Behera,⁵⁹ L. Gladney,⁵⁹ J. Panetta,⁵⁹ M. Biasini,⁶⁰ R. Covarelli,⁶⁰ M. Pioppi,⁶⁰
 C. Angelini,⁶¹ G. Batignani,⁶¹ S. Bettarini,⁶¹ F. Bucci,⁶¹ G. Calderini,⁶¹ M. Carpinelli,⁶¹ R. Cenci,⁶¹ F. Forti,⁶¹
 M. A. Giorgi,⁶¹ A. Lusiani,⁶¹ G. Marchiori,⁶¹ M. A. Mazur,⁶¹ M. Morganti,⁶¹ N. Neri,⁶¹ E. Paoloni,⁶¹ G. Rizzo,⁶¹
 J. Walsh,⁶¹ M. Haire,⁶² D. Judd,⁶² D. E. Wagoner,⁶² J. Biesiada,⁶³ N. Danielson,⁶³ P. Elmer,⁶³ Y. P. Lau,⁶³ C. Lu,⁶³
 J. Olsen,⁶³ A. J. S. Smith,⁶³ A. V. Telnov,⁶³ F. Bellini,⁶⁴ G. Cavoto,⁶⁴ A. D'Orazio,⁶⁴ E. Di Marco,⁶⁴ R. Faccini,⁶⁴
 F. Ferrarotto,⁶⁴ F. Ferroni,⁶⁴ M. Gaspero,⁶⁴ L. Li Gioi,⁶⁴ M. A. Mazzoni,⁶⁴ S. Morganti,⁶⁴ G. Piredda,⁶⁴ F. Polci,⁶⁴
 F. Safai Tehrani,⁶⁴ C. Voena,⁶⁴ M. Ebert,⁶⁵ H. Schröder,⁶⁵ R. Waldi,⁶⁵ T. Adye,⁶⁶ N. De Groot,⁶⁶ B. Franek,⁶⁶
 E. O. Olaiya,⁶⁶ F. F. Wilson,⁶⁶ S. Emery,⁶⁷ A. Gaidot,⁶⁷ S. F. Ganzhur,⁶⁷ G. Hamel de Monchenault,⁶⁷
 W. Kozanecki,⁶⁷ M. Legendre,⁶⁷ B. Mayer,⁶⁷ G. Vasseur,⁶⁷ Ch. Yèche,⁶⁷ M. Zito,⁶⁷ W. Park,⁶⁸ M. V. Purohit,⁶⁸
 A. W. Weidemann,⁶⁸ J. R. Wilson,⁶⁸ M. T. Allen,⁶⁹ D. Aston,⁶⁹ R. Bartoldus,⁶⁹ P. Bechtel,⁶⁹ N. Berger,⁶⁹
 A. M. Boyarski,⁶⁹ R. Claus,⁶⁹ J. P. Coleman,⁶⁹ M. R. Convery,⁶⁹ M. Cristinziani,⁶⁹ J. C. Dingfelder,⁶⁹ D. Dong,⁶⁹
 J. Dorfan,⁶⁹ G. P. Dubois-Felsmann,⁶⁹ D. Dujmic,⁶⁹ W. Dunwoodie,⁶⁹ R. C. Field,⁶⁹ T. Glanzman,⁶⁹ S. J. Gowdy,⁶⁹
 M. T. Graham,⁶⁹ V. Halyo,⁶⁹ C. Hast,⁶⁹ T. Hryn'ova,⁶⁹ W. R. Innes,⁶⁹ M. H. Kelsey,⁶⁹ P. Kim,⁶⁹ M. L. Kocian,⁶⁹
 D. W. G. S. Leith,⁶⁹ S. Li,⁶⁹ J. Libby,⁶⁹ S. Luitz,⁶⁹ V. Luth,⁶⁹ H. L. Lynch,⁶⁹ D. B. MacFarlane,⁶⁹ H. Marsiske,⁶⁹
 R. Messner,⁶⁹ D. R. Muller,⁶⁹ C. P. O'Grady,⁶⁹ V. E. Ozcan,⁶⁹ A. Perazzo,⁶⁹ M. Perl,⁶⁹ B. N. Ratcliff,⁶⁹
 A. Roodman,⁶⁹ A. A. Salnikov,⁶⁹ R. H. Schindler,⁶⁹ J. Schwiening,⁶⁹ A. Snyder,⁶⁹ J. Stelzer,⁶⁹ D. Su,⁶⁹
 M. K. Sullivan,⁶⁹ K. Suzuki,⁶⁹ S. K. Swain,⁶⁹ J. M. Thompson,⁶⁹ J. Va'vra,⁶⁹ N. van Bakel,⁶⁹ M. Weaver,⁶⁹
 A. J. R. Weinstein,⁶⁹ W. J. Wisniewski,⁶⁹ M. Wittgen,⁶⁹ D. H. Wright,⁶⁹ A. K. Yarritu,⁶⁹ K. Yi,⁶⁹ C. C. Young,⁶⁹
 P. R. Burchat,⁷⁰ A. J. Edwards,⁷⁰ S. A. Majewski,⁷⁰ B. A. Petersen,⁷⁰ C. Roat,⁷⁰ L. Wilden,⁷⁰ S. Ahmed,⁷¹
 M. S. Alam,⁷¹ R. Bula,⁷¹ J. A. Ernst,⁷¹ V. Jain,⁷¹ B. Pan,⁷¹ M. A. Saeed,⁷¹ F. R. Wappler,⁷¹ S. B. Zain,⁷¹
 W. Bugg,⁷² M. Krishnamurthy,⁷² S. M. Spanier,⁷² R. Eckmann,⁷³ J. L. Ritchie,⁷³ A. Satpathy,⁷³ C. J. Schilling,⁷³
 R. F. Schwitters,⁷³ J. M. Izen,⁷⁴ I. Kitayama,⁷⁴ X. C. Lou,⁷⁴ S. Ye,⁷⁴ F. Bianchi,⁷⁵ F. Gallo,⁷⁵ D. Gamba,⁷⁵
 M. Bomben,⁷⁶ L. Bosisio,⁷⁶ C. Cartaro,⁷⁶ F. Cossutti,⁷⁶ G. Della Ricca,⁷⁶ S. Dittongo,⁷⁶ S. Grancagnolo,⁷⁶
 L. Lanceri,⁷⁶ L. Vitale,⁷⁶ V. Azzolini,⁷⁷ F. Martinez-Vidal,⁷⁷ Sw. Banerjee,⁷⁸ B. Bhuyan,⁷⁸ C. M. Brown,⁷⁸
 D. Fortin,⁷⁸ K. Hamano,⁷⁸ R. Kowalewski,⁷⁸ I. M. Nugent,⁷⁸ J. M. Roney,⁷⁸ R. J. Sobie,⁷⁸ J. J. Back,⁷⁹
 P. F. Harrison,⁷⁹ T. E. Latham,⁷⁹ G. B. Mohanty,⁷⁹ H. R. Band,⁸⁰ X. Chen,⁸⁰ B. Cheng,⁸⁰ S. Dasu,⁸⁰ M. Datta,⁸⁰
 A. M. Eichenbaum,⁸⁰ K. T. Flood,⁸⁰ J. J. Hollar,⁸⁰ J. R. Johnson,⁸⁰ P. E. Kutter,⁸⁰ H. Li,⁸⁰ R. Liu,⁸⁰ B. Mellado,⁸⁰
 A. Mihalyi,⁸⁰ A. K. Mohapatra,⁸⁰ Y. Pan,⁸⁰ M. Pierini,⁸⁰ R. Prepost,⁸⁰ P. Tan,⁸⁰ S. L. Wu,⁸⁰ Z. Yu,⁸⁰ and H. Neal⁸¹

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²Universitat de Barcelona, Facultat de Física Dept. ECM, E-08028 Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Institute of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

¹⁴University of California at Los Angeles, Los Angeles, California 90024, USA

¹⁵University of California at Riverside, Riverside, California 92521, USA

¹⁶University of California at San Diego, La Jolla, California 92093, USA

¹⁷University of California at Santa Barbara, Santa Barbara, California 93106, USA

¹⁸University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA

¹⁹California Institute of Technology, Pasadena, California 91125, USA

²⁰University of Cincinnati, Cincinnati, Ohio 45221, USA

²¹University of Colorado, Boulder, Colorado 80309, USA

- ²²Colorado State University, Fort Collins, Colorado 80523, USA
- ²³Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany
- ²⁴Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany
- ²⁵Ecole Polytechnique, LLR, F-91128 Palaiseau, France
- ²⁶University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom
- ²⁷Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy
- ²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy
- ²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy
- ³⁰Harvard University, Cambridge, Massachusetts 02138, USA
- ³¹Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany
- ³²Imperial College London, London, SW7 2AZ, United Kingdom
- ³³University of Iowa, Iowa City, Iowa 52242, USA
- ³⁴Iowa State University, Ames, Iowa 50011-3160, USA
- ³⁵Johns Hopkins University, Baltimore, Maryland 21218, USA
- ³⁶Universität Karlsruhe, Institut für Experimentelle Kernphysik, D-76021 Karlsruhe, Germany
- ³⁷Laboratoire de l'Accélérateur Linéaire, IN2P3-CNRS et Université Paris-Sud 11, Centre Scientifique d'Orsay, B.P. 34, F-91898 ORSAY Cedex, France
- ³⁸Lawrence Livermore National Laboratory, Livermore, California 94550, USA
- ³⁹University of Liverpool, Liverpool L69 7ZE, United Kingdom
- ⁴⁰Queen Mary, University of London, E1 4NS, United Kingdom
- ⁴¹University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom
- ⁴²University of Louisville, Louisville, Kentucky 40292, USA
- ⁴³University of Manchester, Manchester M13 9PL, United Kingdom
- ⁴⁴University of Maryland, College Park, Maryland 20742, USA
- ⁴⁵University of Massachusetts, Amherst, Massachusetts 01003, USA
- ⁴⁶Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA
- ⁴⁷McGill University, Montréal, Québec, Canada H3A 2T8
- ⁴⁸Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
- ⁴⁹University of Mississippi, University, Mississippi 38677, USA
- ⁵⁰Université de Montréal, Physique des Particules, Montréal, Québec, Canada H3C 3J7
- ⁵¹Mount Holyoke College, South Hadley, Massachusetts 01075, USA
- ⁵²Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy
- ⁵³NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands
- ⁵⁴University of Notre Dame, Notre Dame, Indiana 46556, USA
- ⁵⁵Ohio State University, Columbus, Ohio 43210, USA
- ⁵⁶University of Oregon, Eugene, Oregon 97403, USA
- ⁵⁷Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
- ⁵⁸Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France
- ⁵⁹University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA
- ⁶⁰Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy
- ⁶¹Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy
- ⁶²Prairie View A&M University, Prairie View, Texas 77446, USA
- ⁶³Princeton University, Princeton, New Jersey 08544, USA
- ⁶⁴Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
- ⁶⁵Universität Rostock, D-18051 Rostock, Germany
- ⁶⁶Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom
- ⁶⁷DSM/Daphnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France
- ⁶⁸University of South Carolina, Columbia, South Carolina 29208, USA
- ⁶⁹Stanford Linear Accelerator Center, Stanford, California 94309, USA
- ⁷⁰Stanford University, Stanford, California 94305-4060, USA
- ⁷¹State University of New York, Albany, New York 12222, USA
- ⁷²University of Tennessee, Knoxville, Tennessee 37996, USA
- ⁷³University of Texas at Austin, Austin, Texas 78712, USA
- ⁷⁴University of Texas at Dallas, Richardson, Texas 75083, USA
- ⁷⁵Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
- ⁷⁶Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
- ⁷⁷IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain
- ⁷⁸University of Victoria, Victoria, British Columbia, Canada V8W 3P6
- ⁷⁹Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom
- ⁸⁰University of Wisconsin, Madison, Wisconsin 53706, USA
- ⁸¹Yale University, New Haven, Connecticut 06511, USA

(Dated: July 5, 2018)

A search for charmed baryons decaying to $D^0 p$ reveals two states: the $\Lambda_c(2880)^+$ baryon and a previously unobserved state at a mass of $[2939.8 \pm 1.3 \text{ (stat.)} \pm 1.0 \text{ (syst.)}] \text{ MeV}/c^2$ and with an intrinsic width of $[17.5 \pm 5.2 \text{ (stat.)} \pm 5.9 \text{ (syst.)}] \text{ MeV}$. Consistent and significant signals are observed for the $K^-\pi^+$ and $K^-\pi^+\pi^-\pi^+$ decay modes of the D^0 in 287 fb^{-1} annihilation data recorded by the *BABAR* detector at a center-of-mass energy of 10.58 GeV . There is no evidence in the D^+p spectrum of doubly-charged partners. The mass and intrinsic width of the $\Lambda_c(2880)^+$ baryon and relative yield of the two baryons are also measured.

PACS numbers: 14.20.Lq, 13.85.Ni

Charmed baryons are expected to exhibit a rich spectrum of states. Only a few of these states have been confirmed [1]. The heaviest singly-charmed baryon previously observed is the $\Lambda_c(2880)^+$ decaying to $\Lambda_c\pi^+\pi^-$ [2]. The $\Lambda_c(2880)^+$ baryon is notable not only due to its narrow width ($< 8 \text{ MeV}$) but also because it is one of only two singly-charmed baryons, along with the $\Xi_c(2815)$ [3], found above the Dp mass threshold.

Presented in this Letter is the observation of a new charmed baryon decaying to $D^0 p$ [4] with a mass of approximately $2.94 \text{ GeV}/c^2$ and an intrinsic width of approximately 20 MeV . This baryon, tentatively labeled the $\Lambda_c(2940)^+$, is observed in 287 fb^{-1} of e^+e^- annihilation data collected near $\sqrt{s} = 10.58 \text{ GeV}$ by the *BABAR* detector [5] at the PEP-II asymmetric-energy e^+e^- storage rings. Along with this new baryon, the decay $\Lambda_c(2880)^+ \rightarrow D^0 p$ is also observed. The masses, intrinsic widths of both baryons and their relative production rate are measured.

The goal of this analysis is to study the inclusive $D^0 p$ mass spectrum. Two samples of D^0 mesons are identified using the $K^-\pi^+$ and $K^-\pi^+\pi^-\pi^+$ final states. Each sample is produced by combining charged tracks of the appropriate composition in a geometric fit to a common vertex. The χ^2 probability of this fit is required to exceed 2%. Charged particle species (K^+, π^+, p) are separated using a likelihood algorithm that combines data from a ring-imaging Cherenkov detector with the measured energy loss in the tracking systems [5]. Each proton candidate is combined with each D^0 candidate using a geometric vertex fit that assumes a common production point within the nominal beam envelope. The χ^2 probability of this fit is required to be better than 2%.

Requirements are imposed on three additional quantities to improve the signal purity of the $D^0 p$ samples: Δm , the difference between the reconstructed D^0 mass and the accepted value of $m_{D^0} = 1864.6 \text{ MeV}/c^2$ [1]; p^* , the center-of-mass momentum of the $D^0 p$ system; and $\cos\vartheta$, where ϑ is angle of the proton with respect to the e^+e^- system in the $D^0 p$ center-of-mass frame. For isotropic production (expected for the $\Lambda_c(2940)^+$), the $\cos\vartheta$ distribution will be flat whereas background tends to peak at ± 1 . Studies of Monte Carlo (MC) simulated data samples are used to determine the specific requirements on these quantities that maximize the expected significance of signals introduced in the mass re-




FIG. 1: The solid points are the $D^0 p$ invariant mass distribution of the final sample. Also shown are (gray) the contribution from false D^0 candidates estimated from D^0 mass sidebands and (open points) the mass distribution from wrong-sign $\bar{D}^0 p$ candidates. The solid curve is the fit described in the text. The dashed curve is the portion of that fit attributed to combinatorial background.

gion near $2940 \text{ MeV}/c^2$. The resulting best criteria are $|\Delta m| < 14 \text{ MeV}/c^2$, $p^* > 2.6 \text{ GeV}/c$, and $\cos\vartheta < 0.8$ for the $D^0 \rightarrow K^-\pi^+$ sample and $|\Delta m| < 9 \text{ MeV}/c^2$, $p^* > 2.8 \text{ GeV}/c$, and $\cos\vartheta < 0.8$ for the $D^0 \rightarrow K^-\pi^+\pi^-\pi^+$ sample. The Δm requirements correspond to approximately two standard deviations in D^0 mass resolution. The p^* requirement removes all sources of $D^0 p$ combinations from B meson decay.

A MC simulation of a baryon of mass $2.94 \text{ GeV}/c^2$ decaying to $D^0 p$ predicts selection efficiencies between 30% and 38% for the $D^0 \rightarrow K^-\pi^+$ final state depending on p^* and between 12% and 14% for the $D^0 \rightarrow K^-\pi^+\pi^-\pi^+$ final state. A proton purity of approximately 83% in the final $D^0 p$ sample is estimated from studies of a comparable MC sample.

To calculate a $D^0 p$ invariant mass, each D^0 candidate is assigned an energy that is consistent with a D^0 mass of m_{D^0} . The resulting combined $D^0 p$ invariant mass spec-

trum is shown in Fig. 1. Two peaks are apparent. The clear signal at $2.88 \text{ GeV}/c^2$ is likely due to the decay of the $\Lambda_c(2880)^+$ baryon. The signal at $2.94 \text{ GeV}/c^2$ is the evidence for the new $\Lambda_c(2940)^+$ baryon. No similar structures are observed in the wrong-sign $\bar{D}^0 p$ candidate combinations. Candidates selected from D^0 mass sidebands are used to estimate the contribution from non- D^0 sources (see Fig. 1). This sideband sample shows no structure.

An unbinned likelihood fit is used to model the $D^0 p$ spectrum from the kinematic limit up to $3.05 \text{ GeV}/c^2$. This fit includes $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ states, each modeled by a relativistic Breit-Wigner lineshape $\sigma(m)$ convolved with a Gaussian resolution function. The Breit-Wigner line shape $\sigma(m)$ is:

$$\sigma(m) \propto \frac{q(m)}{(m^2 - m_0^2)^2 + m_0^2 \Gamma^2}, \quad (1)$$

where Γ is the intrinsic width and is constant (i.e. not mass dependent), m_0 is the mass pole, and q is the three-momentum magnitude of the D^0 or proton in the $D^0 p$ rest frame for a given mass m . The detector resolution is obtained from MC simulation which predicts $1.8 \text{ MeV}/c^2$ and $1.3 \text{ MeV}/c^2$ for the $D^0 \rightarrow K^-\pi^+$ and $D^0 \rightarrow K^-\pi^+\pi^-\pi^+$ samples, respectively.

The product of a fourth-order polynomial and two-body phase space [1] is used to model the combinatorial background. A fit based on this background shape and the $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ signals is shown in Fig. 1 and results in a $\Lambda_c(2940)^+$ mass of $2939.8 \pm 1.3 \text{ MeV}/c^2$, a width of $17.5 \pm 5.2 \text{ MeV}$, and a raw yield of 2280 ± 310 decays (statistical errors only). The $\Lambda_c(2880)^+$ properties obtained are a mass of $2881.9 \pm 0.1 \text{ MeV}/c^2$ and a width of $5.8 \pm 1.5 \text{ MeV}$, consistent with the CLEO results [2], and a raw yield of 2800 ± 190 decays (statistical errors only). If the $\Lambda_c(2940)^+$ signal is removed from the fit, the log likelihood changes by 38.2, which is equivalent (in one degree of freedom) to a signal significance of 8.7 standard deviations. If the $D^0 \rightarrow K^-\pi^+$ and $D^0 \rightarrow K^-\pi^+\pi^-\pi^+$ samples are fit separately, the resulting masses, widths, and relative yields of the $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ baryons are consistent within statistical errors. After accounting for selection efficiency and D^0 branching fractions, the absolute yields for the two D^0 decays modes are consistent for both the $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ baryons.

The above likelihood fit models the mass spectrum near $2.84 \text{ GeV}/c^2$ as a smooth distribution (Fig. 2(a)). There is, however, a non-distinct structure near a mass of $2.84 \text{ GeV}/c^2$ whose origin is not understood, and so this model may not be accurate. Various modifications of the fit are employed as systematic checks. At one extreme, if the likelihood fit is limited to masses above $2.8525 \text{ GeV}/c^2$ (Fig. 2(b)), the result is a substantial decrease (29%) in the $\Lambda_c(2940)^+$ yield, a $0.5 \text{ MeV}/c^2$ shift




FIG. 2: Three examples of how the structure near a $D^0 p$ mass of $2.84 \text{ GeV}/c^2$ can be modeled. Shown are the results of fits that (a) assume a smooth distribution (as used for the central result) (b) exclude data below a mass of $2.8525 \text{ GeV}/c^2$, and (c) add an extra resonance contribution.

in mass, and a smaller width (12.5 MeV). The changes in the fitted $\Lambda_c(2940)^+$ properties are much smaller if a third signal line shape (of variable mass and width) is added to the fit (Fig. 2(c)). None of these alternate fits lead to a reduction in the statistical significance of the $\Lambda_c(2940)^+$ signal below 7.2 standard deviations.

Because the $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ are near the $D^0 p$ threshold, the systematic uncertainty in mass from possible detector biases is relatively small. This uncertainty is calculated by considering appropriate variations in the assumed B field strength and detector material using a procedure developed for measuring the Λ_c mass [6]. This procedure is also used to calculate small ($< 0.1 \text{ MeV}/c^2$) corrections to the reconstructed $D^0 p$ mass. An additional uncertainty of $0.5 \text{ MeV}/c^2$ arises from the current knowledge of m_{D^0} . The results for the $\Lambda_c(2940)^+$ baryon are:

$$m = [2939.8 \pm 1.3 \text{ (stat.)} \pm 1.0 \text{ (syst.)}] \text{ MeV}/c^2$$

$$\Gamma = [17.5 \pm 5.2 \text{ (stat.)} \pm 5.9 \text{ (syst.)}] \text{ MeV}.$$

For the $\Lambda_c(2880)^+$ baryon the results are:

$$m = [2881.9 \pm 0.1 \text{ (stat.)} \pm 0.5 \text{ (syst.)}] \text{ MeV}/c^2$$

$$\Gamma = [5.8 \pm 1.5 \text{ (stat.)} \pm 1.1 \text{ (syst.)}] \text{ MeV}.$$

From the baryon yields obtained from the likelihood fits, the following ratio of production cross sections and decay

branching ratios is calculated:

$$\frac{\sigma(\Lambda_c(2940)^+)\mathcal{B}r(\Lambda_c(2940)^+ \rightarrow D^0 p)}{\sigma(\Lambda_c(2880)^+)\mathcal{B}r(\Lambda_c(2880)^+ \rightarrow D^0 p)} = 0.81 \pm 0.13 \text{ (stat.)} \pm 0.35 \text{ (syst.)},$$

where the systematic uncertainty is dominated by uncertainties in the background shape.

Various tests are applied to the data to confirm the $\Lambda_c(2940)^+$ signal. Since the signal is observed in two different D^0 decay modes, it appears to be associated with real D^0 decays. The lack of any structure in the D^0 side-band samples and the relative size of these samples support this conclusion. Since the sample of protons is 83% pure, it is unlikely that the $\Lambda_c(2940)^+$ signal could arise from proton mis-identification. As further confirmation, when the K^+ or π^+ mass is assigned to the protons, the resulting $D^0 K^+$ and $D^0 \pi^+$ invariant mass distributions show no evidence of structure.

Even if the observed signal is attributed to a combination of D^0 and protons, it is still possible to produce a false signal from the reflection of heavier states. One example of such a possible reflection is a hypothetical baryon of mass near $3.10 \text{ GeV}/c^2$ decaying to either $D^*(2010)^+ p$ or $D^*(2007)^0 p$. Such a baryon, if sufficiently narrow, would produce a $D^0 p$ mass spectrum (after ignoring the π^+ or π^0 from D^* decay) of approximately the correct mass and width. Such a baryon would also be clearly visible in the $D^*(2010)^+ p$ or $D^*(2007)^0 p$ mass distributions. An explicit search in those mass distributions shows no signal, and thus this hypothesis is strongly disfavored.

Another possible reflection is from a baryon of mass $3.13 \text{ GeV}/c^2$ decaying to $D^0 \Sigma^+$. The kinematics of such a decay could produce peaks at both $2.85 \text{ GeV}/c^2$ and $2.94 \text{ GeV}/c^2$ if the Σ^+ had the appropriate spin alignment. The Σ^+ , however, is a long-lived particle, and MC studies indicate that for this decay the proton vertex χ^2 probability distribution would peak at zero. An investigation of the χ^2 probability of the $\Lambda_c(2940)^+$ signal seen in the data indicates a flat distribution. Thus, a reflection from $D^0 \Sigma^+$ decay is also strongly disfavored.

The simplest interpretation of the $\Lambda_c(2940)^+$ signal is that it arises from a charmed baryon of quark content cdu . Under this scenario the decay to $D^0 p$ involves simple $u\bar{u}$ gluon splitting. The remaining question is whether the $\Lambda_c(2940)^+$ belongs to an isotriplet. The most direct way to address this question is to explicitly search for a neutral or doubly-charged partner of nearly the same mass and width, analogous to the Σ_c^0 and Σ_c^{++} . The BABAR detector cannot isolate the most obvious neutral decay mode ($D^0 n$). It is possible, however, to search for a doubly-charged baryon decaying to $D^+ p$.

To select a sample of D^+ candidates, the same methods used for the D^0 samples are applied to the decay $D^+ \rightarrow K^- \pi^+ \pi^+$. The selection requirements for the




FIG. 3: The invariant mass distribution of selected $D^+ p$ candidates. The curve is the result of the fit described in the text. The curves below are the lineshapes of the $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ baryons obtained from the $D^0 p$ data, drawn approximately to scale after correcting for selection efficiency and D^0 and D^+ branching fractions.

$D^+ p$ sample are $|\Delta m| < 12 \text{ MeV}/c^2$, $p^* > 2.7 \text{ GeV}/c$, and $\cos \vartheta < 0.8$. The efficiency for this selection is approximately 23%.

The resulting $D^+ p$ distribution is shown in Fig. 3. No signals corresponding to either the $\Lambda_c(2880)^+$ or $\Lambda_c(2940)^+$ baryon are apparent. A likelihood fit which assumes a doubly-charged partner of the $\Lambda_c(2940)^+$ of identical mass and width results in a yield of -40 ± 120 candidates (statistical error only).

Based on previous observations, such as the CLEO measurement of the Σ_c^0 and Σ_c^{++} [7], one would expect similar production rates for the $\Lambda_c(2940)^+$ and a hypothetical doubly-charged partner. Under the additional assumption that the branching fraction of the doubly-charged baryon to Dp is the same, the expected doubly-charged signal yield would be approximately 2200 decays once the D^0 and D^+ branching fractions and selection efficiencies are accounted for (see Fig. 3). It thus seems unlikely that a doubly-charged partner exists, unless its production is largely suppressed or it decays in an unexpected fashion.

The $\Lambda_c(2940)^+$ baryon is interesting for several reasons. Relativistic quark model calculations [8] predict three excited Λ_c baryons of different spin-parity quantum numbers near a mass of $2.94 \text{ GeV}/c^2$. The DN decay mode, although not unexpected [9, 10], is a final state that has received relatively little theoretical investigation. If this baryon had a significant branching fraction to $\Lambda_c \pi^+ \pi^-$ it probably would have been observed with the $\Lambda_c(2880)^+$ by CLEO [2]. It is not clear, however, why this particular decay mode, which is favored by phase space, is suppressed. One observation which is notable, even if it might be a simple coincidence, is that at

a mass of $2939.8 \text{ MeV}/c^2$, the $\Lambda_c(2940)^+$ is just $6 \text{ MeV}/c^2$ below the $D^{*0}p$ threshold. It is also interesting that the $\Lambda_c(2940)^+$ is approximately one pion mass heavier than the $\Sigma_c(2800)^+$, a charmed baryon recently discovered by BELLE [11] decaying to $\Lambda_c\pi^0$.

The $\Lambda_c(2880)^+$ mass and width results presented here are consistent with but more precise than the CLEO measurement of $m = 2880.9 \pm 2.3 \text{ MeV}/c^2$ and $\Gamma < 8 \text{ MeV}$ (at 90% CL). The existence of the decay $\Lambda_c(2880)^+ \rightarrow D^0p$ rules out various interpretations of this baryon [10].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mexico), Marie Curie EIF (European Union), the A. P. Sloan Foundation, the Research Corporation, and the Alexander von Humboldt Foundation.

[†] Also with Università di Perugia, Dipartimento di Fisica, Perugia, Italy

[‡] Also with Università della Basilicata, Potenza, Italy

- [1] S. Eidelman et al. (Particle Data Group), Phys. Lett. **B592**, 1 (2004).
- [2] M. Artuso et al. (CLEO), Phys. Rev. Lett. **86**, 4479 (2001).
- [3] J. P. Alexander et al. (CLEO), Phys. Rev. Lett. **83**, 3390 (1999).
- [4] Inclusion of charge conjugate states is implied throughout this paper.
- [5] B. Aubert et al. (*BABAR*), Nucl. Instrum. Meth. **A479**, 1 (2002).
- [6] B. Aubert et al. (*BABAR*), Phys. Rev. **D72**, 052006 (2005).
- [7] M. Artuso et al. (CLEO), Phys. Rev. **D65**, 071101 (2002).
- [8] S. Migura, D. Merten, B. Metsch, and H.-R. Petry (2006), submitted to Eur. Phys. J. A, hep-ph/0602153.
- [9] D. Pirjol and T.-M. Yan, Phys. Rev. **D56**, 5483 (1997).
- [10] A. E. Blechman, A. F. Falk, D. Pirjol, and J. M. Yelton, Phys. Rev. **D67**, 074033 (2003).
- [11] R. Mizuk et al. (Belle), Phys. Rev. Lett. **94**, 122002 (2005).

* Also at Laboratoire de Physique Corpusculaire, Clermont-Ferrand, France