

day 34 prep

working on over-relaxation and gauss-seidel methods

first over-relaxation

```
In [1]: M = 100 # grid size
        V = 1.0 # voltage

        target = 2**-.16

        phi = np.zeros([M+1, M+1], float)
        phi[0, :] = V

        phiprime = np.empty([M+1, M+1], float)


        delta = 1.0

        while delta > target:
            for i in range(M+1):
                for j in range(M+1):
                    if i == 0 or i == M or j == 0 or j == M:
                        phiprime[i, j] = phi[i, j]
                    else:
                        phiprime[i, j] = (phi[i+1, j] + phi[i-1, j] + phi[i, j+1] + phi[i, j-1])
            delta = np.max(np.abs(phi - phiprime))
            phi, phiprime = phiprime, phi

        fig0, ax0 = plt.subplots()
        ax0.imshow(phi)
```

```
Out[1]: <matplotlib.image.AxesImage at 0x701dd7753fb0>
```

Figure



```
In [2]: phi = np.zeros([M+1, M+1], float)
phi[0,:] = V


dphi = np.empty([M+1,M+1],float)

delta = 1.0
while delta>target:
    for i in range(M+1):
        for j in range(M+1):
            if i==0 or i==M or j==0 or j==M: # so this is around the boundary
                #phiprime[i,j] = phi[i,j]
                phi[i,j] = phi[i,j]
                dphi[i,j] = 0
            else:
                dphi[i,j] = (phi[i+1,j]+phi[i-1,j]+phi[i,j+1]+phi[i,j-1])/4-
                phi[i,j]
                phi[i,j] = phi[i,j]+dphi[i,j]
        delta = np.max(np.abs(dphi))
    # phi, phiprime = phiprime, phi

fig1, ax1 = plt.subplots()
ax1.imshow(phi)
```

```
Out[2]: <matplotlib.image.AxesImage at 0x701dd6432f60>
```

Figure



```
In [3]: phi = np.zeros([M+1, M+1], float)
        phi[0,:] = V
        # phi[M,:] = V


        dphi = np.empty([M+1,M+1],float)
        omega = 0.9

        delta = 1.0
        while delta>target:
            for i in range(M+1):
                for j in range(M+1):
                    if i==0 or i==M or j==0 or j==M: # so this is around the boundary
                        phi[i,j] = phi[i,j]
                        dphi[i,j] = 0
                    else:
                        dphi[i,j] = (1+omega)*((phi[i+1,j]+phi[i-1,j]+phi[i,j+1]+phi[i,j-1]) - 4*phi[i,j])
                        phi[i,j] = phi[i,j]+dphi[i,j]
                delta = np.max(np.abs(dphi))

        fig2, ax2 = plt.subplots()
        ax2.imshow(phi)
```

Out[3]: <matplotlib.image.AxesImage at 0x701dd64b9190>

Figure



so for homework I did not make a mistake but I need to clarify:

what i told the students to do was to make those spots be like boundaries in that they could be set to be +1 and -1 volts. if i do that problem that is slightly different from the problem as it is stated and intended to be worked out in the book. So first I'll work my problem, and then i'll work the problem in the book and then i'll look at some variations to speed up my code after that

```
In [16]: M = 100 # grid size
V = 1

target = 2** -16

phi = np.zeros([M+1, M+1], float)


phi[20:40, 60:80] = V

phi[60:80, 20:40] = -V

fig3, ax3 = plt.subplots()
image = ax3.imshow(phi.T, origin='lower')
```

```
plt.colorbar(image)
fig3.tight_layout()
```

Figure




```
In [7]: dphi = np.empty([M+1,M+1],float)
        omega = 0.9

        delta = 1.0
        while delta>target:
            for i in range(M+1):
                for j in range(M+1):
                    if i==0 or i==M or j==0 or j==M or (20<=i<40 and 60<=j<80) or (60<=i<80 and 20<=j<40):
                        phi[i,j] = phi[i,j]
                        dphi[i,j] = 0
                    else:
                        dphi[i,j] = (1+omega)*((phi[i+1,j]+phi[i-1,j]+phi[i,j+1]+phi[i,j-1]) - 4*phi[i,j])
                        phi[i,j] = phi[i,j]+dphi[i,j]
            delta = np.max(np.abs(dphi))

        fig4, ax4 = plt.subplots()
        ax4.imshow(phi, origin='lower')
        ax4.set_xlabel('x')
        ax4.set_ylabel('y')
```

Out[7]: Text(0, 0.5, 'y')

Figure



Ok but I'm not sure that is what the question is asking and I think I mislead the students.

I think the question is coming from painting in the charge density, not painting in the voltage, and that difference is important.

$$\nabla^2 \phi = -\frac{\rho}{\epsilon_0}$$

which means we are designating the boundaries of some charge density... no voltage.

Which means that to first order approximation:

$$\nabla^2 \phi = \frac{\partial^2 \phi}{\partial x^2} + \frac{\partial^2 \phi}{\partial y^2} = \frac{\phi(x+a, y) + \phi(x-a, y) + \phi(x, y+a) + \phi(x, y-a) - 4\phi(x, y)}{a^2}$$

So, if that is equal to zero, then we have the formula that we used up above:

$$\frac{\phi(x+a, y) + \phi(x-a, y) + \phi(x, y+a) + \phi(x, y-a) - 4\phi(x, y)}{a^2} = 0$$

$$\phi(x+a, y) + \phi(x-a, y) + \phi(x, y+a) + \phi(x, y-a) - 4\phi(x, y) = 0$$

$$\phi(x+a, y) + \phi(x-a, y) + \phi(x, y+a) + \phi(x, y-a) = 4\phi(x, y)$$

$$\frac{\phi(x+a, y) + \phi(x-a, y) + \phi(x, y+a) + \phi(x, y-a)}{4} = \phi(x, y)$$

And that is just fine. But when this is Poisson's equation then things change a bit

$$\frac{\phi(x+a, y) + \phi(x-a, y) + \phi(x, y+a) + \phi(x, y-a) - 4\phi(x, y)}{a^2} = \frac{\rho}{\epsilon_0}$$

So rearranging this means that a^2 sticks around with the ρ/ϵ .

$$\phi(x, y) = \frac{1}{4}(\phi(x+a, y) + \phi(x-a, y) + \phi(x, y+a) + \phi(x, y-a) + a^2 \frac{\rho}{\epsilon_0})$$

Now, don't forget that ρ is a spacial dependent quantity. So we should really define at the beginning a matrix for ρ and then we can use it in what comes after.

```
In [11]: M = 100 # grid size
V = 1

target = 2**-16

phi = np.zeros([M+1, M+1], float)

rho_0 = 1

rho = np.zeros([M+1, M+1], float)

rho[20:40, 60:80] = rho_0


rho[60:80, 20:40] = -rho_0

dphi = np.empty([M+1, M+1], float)
omega = 0.9

delta = 1.0
while delta > target:
    for i in range(M+1):
        for j in range(M+1):
            # this time I don't need that weird boundary so this is nice!
            if i==0 or i==M or j==0 or j==M:
                # continue
                phi[i, j] = phi[i, j]
                dphi[i, j] = 0
            else:
                dphi[i, j] = (phi[i+1, j] + phi[i-1, j] + phi[i, j+1] + phi[i, j-1] + rho[i, j] - 4*phi[i, j]) / a**2
                phi[i, j] = phi[i, j] + (1+omega)*dphi[i, j]
        delta = np.max(np.abs(dphi))

fig4, ax4 = plt.subplots()
image = ax4.imshow(phi, origin='lower')
ax4.set_xlabel('x')
ax4.set_ylabel('y')
```

```
plt.colorbar(image)
fig4.tight_layout()
```



Now I want to subtly mess with things to optimize them all the way. like do i even need that dphi matrix at all?

```
In [23]: M = 100 # grid size
V = 1

phi = np.zeros([M+1, M+1], float)

phi[0,:] = V

rho_0 = 1

rho = np.zeros([M+1, M+1], float)

rho[20:40, 60:80] = rho_0

rho[60:80, 20:40] = -rho_0

dphi = np.empty([M+1,M+1],float)
omega = 0.9

delta = 1.0
while delta>target:
    for i in range(M):
```



```


for j in range(M):
    # this time I don't need that weird boundary so this is nice!
    if i!=0 or i!=M or j!=0 or j!=M:
        dphi[i,j] = (phi[i+1,j]+phi[i-1,j]+phi[i,j+1]+phi[i,j-1]+rho
        phi[i,j] = phi[i,j]+(1+omega)*dphi[i,j]
delta = np.max(np.abs(dphi))

fig4, ax4 = plt.subplots()
ax4.imshow(phi, origin='lower')
ax4.set_xlabel('x')
ax4.set_ylabel('y')

```

Out[23]: Text(0, 0.5, 'y')

Figure



In []: