

day 27 prep

we are now ready to handle 2nd order differential equations. the method will be similar to coupled first order equations because that is exactly how we will treat any second order equation: as a set of coupled first order equations like below.

Let's take an example second order equation and look at the technique. We will take a mass on a spring under damped conditions:

$$ma = -cv - kx$$

The first thing we want to do is put this in terms of all of the derivative:

$$m \frac{d^2x}{dt^2} = -c \frac{dx}{dt} - kx$$

Then we want to isolate the second derivative.

$$\frac{d^2x}{dt^2} = -\frac{c}{m} \frac{dx}{dt} - \frac{k}{m}x$$

Then we want to break this up into two first order differential equations. And this looks a little bit backwards, but we need to realize that we just replaced v with $\frac{dx}{dt}$ so what if we went backwards that in terms of the substitution of variables.

$$\frac{dv}{dt} = -\frac{c}{m}v - \frac{k}{m}x$$

and then

$$\frac{dx}{dt} = v$$

```
In [1]: %matplotlib ipynpl

import numpy as np
import matplotlib.pyplot as plt
```

```
In [20]: # Runge-Kutta 4th order

def f(r,t):
    x = r[0]
    v = r[1]
    c = 5
    m = 1
    k = 100
    fx = v
    fv = -c/m*v - k/m*x
```

```

    return(np.array([fx,fv],float))

# define boundary conditions

t0 = 0.0 # starting point
tf = 2.0 # ending point
N = 1000 # number of points between a and b
dt = (tf-t0)/N

r = np.array([100,0],float) # initial condition

tpoints = np.arange(t0, tf, dt)
xpoints = []
vpoints = []


for t in tpoints:
    xpoints.append(r[0])
    vpoints.append(r[1])
    k1 = dt*f(r,t)
    k2 = dt*f(r+0.5*k1,t+0.5*dt)
    k3 = dt*f(r+0.5*k2,t+0.5*dt)
    k4 = dt*f(r+k3, t+dt)
    r = r + (k1+2*k2+2*k3+k4)/6

fig0, ax0 = plt.subplots()
ax0.plot(tpoints, xpoints)
#ax0.plot(tpoints, vpoints)

```

Out[20]: [

Figure



In []: