

day 13 | 250926 F

Let's integrate data.

```
In [1]: import numpy as np
import pandas as pd
import matplotlib.pyplot as plt
```

```
In [2]: df = pd.read_csv('../data/velocities.txt', sep='\t', names=['t', 'v'])
df
```

```
Out[2]:
```

	t	v
0	0	0.000000
1	1	0.069478
2	2	0.137694
3	3	0.204332
4	4	0.269083
...	...	...
96	96	0.223073
97	97	0.254244
98	98	0.283753
99	99	0.311479
100	100	0.337308

101 rows × 2 columns

```
In [52]: def trap(x_data, y_data):
    if type(x_data) == pd.core.series.Series:
        x_data = np.asarray(x_data)
        y_data = np.asarray(y_data)
    dx = (x_data[-1] - x_data[0]) / len(x_data)
    s = dx / 2 * (y_data[0] + y_data[-1])
    s = s + np.sum(y_data[1:-2] * dx)
    return(s)

def simp(func, a, b, steps):
    if steps % 2 != 0:
        steps = steps + 1
        print('i added one step')
    dx = (b - a) / steps
    x = np.linspace(a, b, steps + 1)
```

```

y = func(x)
s = dx/3*(func(a)+func(b))
s = s + dx/3*(4*np.sum(y[1:steps:2]) + 2*np.sum(y[2:steps-1:2]))
return(s)

```

```

In [53]: x = np.linspace(0,2, 10000)
y = x**4 - 2*x + 1

```

```

In [54]: trap(x, y)

```

```

Out[54]: 4.396961306605322

```

```

In [55]: trap(df['t'], df['v'])

```

```

Out[55]: 7.8291534653465344

```

```

In [56]: integrals = [0]

```

```

for i in range(1,101):
    w = trap(df['t'][0:i], df['v'][0:i])
    integrals.append(w)

```

```


In [57]: fig, ax = plt.subplots()
ax.plot(df['t'], df['v'])
ax.plot(df['t'], integrals)

```

```

Out[57]: [<matplotlib.lines.Line2D at 0x76b1d48bc740>]

```



```
In [58]: integrals[-1]
```

```
Out[58]: 7.534669724999998
```

```
In [ ]:
```

```
In [63]: df['t'].values[-1]
```

```
Out[63]: 100
```

```
In [29]: df['t'].index
```

```
Out[29]: RangeIndex(start=0, stop=101, step=1)
```

```
In [33]: range(1,10)
```

```
Out[33]: range(1, 10)
```

```
In [ ]:
```