

Chapter 3 - Momentum + Angular Momentum

Recall: $\vec{p} = m\vec{v}$ $\sum \vec{F}_n = \dot{\vec{p}}$

$$\vec{P} = \vec{p}_1 + \vec{p}_2 + \vec{p}_3 \dots = \sum \vec{p}_n$$

$$\dot{\vec{P}} = \vec{F}_{\text{ext}}$$

If $\vec{F}_{\text{ext}} = 0$

$$\dot{\vec{P}} = 0$$

$\rightarrow \vec{P}_i = \vec{P}_f$ Conservation of Momentum

If two objects,

$$\vec{p}_{1i} + \vec{p}_{2i} = \vec{p}_{1f} + \vec{p}_{2f}$$

$$m_1 \vec{v}_{1i} + m_2 \vec{v}_{2i} = m_1 \vec{v}_{1f} + m_2 \vec{v}_{2f}$$

← really 2 equations

If collision is perfectly inelastic
↳ stick together

then,

$$m_1 \vec{v}_{1i} + m_2 \vec{v}_{2i} = m_1 \vec{v}_f + m_2 \vec{v}_f$$
$$= (m_1 + m_2) \vec{v}_f$$

mass add
up!


$$\vec{v}_f = \frac{m_1 \vec{v}_{1i} + m_2 \vec{v}_{2i}}{m_1 + m_2}$$

✓ ← two equations!

Rockets + Explosions

At the beginning

$$P = 0$$



Round 1

$$P_i = 0 = P_f = p_{ME} + p_{ball}$$

$$0 = \underbrace{(75 \text{ kg} + 99(0.5 \text{ kg}))}_ {124.5 \text{ kg}} v + (0.5 \text{ kg})(-15 \text{ m/s})$$

$$0 = 124.5 v - 7.5 \frac{\text{kg m}}{\text{s}}$$


$$v = \frac{7.5 \text{ kg m/s}}{124.5 \text{ kg}} = 0.0602 \text{ m/s}$$

Row 12

$$V' = V_{ice} - V_{s'}$$
$$-15 \text{ m/s} = V_{ice} - 0.0602 \text{ m/s}$$

$$-15 + 0.06 = V_{ice} = -14.94 \text{ m/s}$$

0 =



ice

$$P_i = P_f$$

$$(124.5)(0.0602 \text{ m/s}) + 0.5(-15) = 124 \cdot v + 0.5(-15) + 0.5(-14.94)$$

= 0

$$-0.5(-15 - 14.94) = 124 \cdot v$$

$$v = \frac{-0.5(-15 - 14.94)}{124} = \frac{-0.5(-15 - 15 + 0.0602)}{124}$$

$$= 0.1207 \text{ m/s}$$

but we need a pattern to do this numerically - this way is not giving that away

$$(m + \Delta m)(v + \Delta v) = mv + \Delta mv + m\Delta v + \Delta m\Delta v$$

$$\Delta P_{ME} = \Delta P_{ball}$$

$$124.5(0.0602) - 124v = 0.5(-14.94)$$

$m\Delta v \quad mv \quad \Delta mv$

$$0.5(15) + 0.5 \cdot 0.0602$$

$$\frac{0.5(14.94)}{124}$$

$$0.0602410$$

$$\frac{0.5(15) - 0.5(15)}{124}$$

$$\frac{0.5(15)}{124.5}$$

$$0.0602406$$

$$\frac{0.5(15-x) - x}{124} = ?$$

IGNORE

THESE

NOTES,
REMINGTON
AT WORK

$$v = \frac{-0.5(-15 - 14.94)}{124} = \frac{0.5(15 + 14.94)}{124}$$
$$= \frac{0.5(15)}{124} + \frac{0.5(14.94)}{124}$$

$0.06048 \quad 0.0602$

$$\frac{0.5(14.94)}{124} \stackrel{?}{=} \frac{0.5(15)}{124.5}$$

$0.0602 = 0.0602$

Round 1 Redux - my frame

$$P_i = 0 = P_f = p_{ME} + p_{ball}$$

$$0 = \underbrace{(75 \text{ kg} + 99(0.5 \text{ kg}))}_ {124.5 \text{ kg}} v + (0.5 \text{ kg})(-15 \text{ m/s})$$

$$0 = 124.5 v - 7.5 \frac{\text{kg m}}{\text{s}}$$

$$v = \frac{7.5 \text{ kg m/s}}{124.5 \text{ kg}} = 0.0602 \text{ m/s}$$

Round 2 Redux - my frame

P_i'

$$0 = 124.0 v + 0.5(-15)$$

$$v = \frac{0.5(15)}{124} = 0.06048 \text{ m/s}$$

$$\frac{0.5(15 - 0.602)}{124} =$$

$$\frac{0.5(15) - 0.5(15)}{124} = \frac{0.5(15)}{124.5}$$

$$\frac{1}{124} - \frac{0.5}{124.5} = \frac{1}{124.5}$$

Round 3 Redux - my frame

P_i''

$$D = 123.5 \text{ V} + 0.5(-15)$$

$$V = \frac{0.5(15)}{123.5} = 0.0607 \text{ m/s}$$

0.18

Generalize

$$dV = \frac{dP_{ex}}{m} \quad \leftarrow \text{Variable}$$

$$\int_{V_i}^{V_f} dV = \int_{m_0 + m_f}^{m_0} \frac{V_{ex} dm}{m} = V_{ex} \int_{m_0 + m_f}^{m_0} \frac{dm}{m}$$

$$V \Big|_{v_i}^{v_f} = V_{ex} \ln(m) \Big|_{m_0+m_f}^{m_0}$$

$$v_f - v_i = V_{ex} \left[\ln(m_0) - \ln(m_0 + m_f) \right]$$

$$\underline{v_f = V_{ex} \ln\left(\frac{m_0}{m_0 + m_f}\right) + v_i}$$

$$\hookrightarrow \underline{\underline{V(t) = V_{ex} \ln\left(\frac{m(t)}{m_0 + m_f}\right) + v_i}}$$

test w berechnung

$$\begin{aligned} v_f &= -15 \ln\left(\frac{75\text{kg}}{125\text{kg}}\right) \\ &= \underline{\underline{+7.66 \text{ m/s}}} \end{aligned}$$

$$\rightarrow V(t) = V_{0x} \ln \left(\frac{m(t)}{m_0 + m_f} \right) + V_0$$

$$0 = +15 \ln \left(\frac{m_0}{m_0 + 0.5n} \right) + 10$$

How many
baseballs to stop
from 10 m/s

$$-10 = +15 \ln \left(\frac{m_0}{m_0 + 0.5n} \right)$$

$$\frac{-10}{+15} = \ln \left(\frac{m_0}{m_0 + 0.5n} \right)$$

$$e^{\frac{-10}{15}} = \frac{m_0}{m_0 + 0.5n}$$


$$e^{\frac{-10}{15}} (m_0 + 0.5n) = m_0$$

$$m_0 + 0.5n = m_0 e^{\frac{10}{15}}$$

$$n = \frac{m_0 (e^{\frac{10}{15}} - 1)}{0.5}$$

$$m_0 = 75 \text{ kg}$$

$$n = 142.16 \leftarrow \# \text{ of baseballs}$$



Center of Mass

$$\vec{R} = \frac{m_1 \vec{r}_1 + m_2 \vec{r}_2 + \dots}{m_1 + m_2 + \dots}$$

total, M

$$\vec{R} = \frac{1}{M} \sum_{\alpha=1}^N m_{\alpha} \vec{r}_{\alpha}$$

center of mass is a position

$$X = \frac{1}{M} \sum_{\alpha=1}^N m_{\alpha} X_{\alpha}$$

x-component (x position)

$$Y = \frac{1}{M} \sum_{\alpha=1}^N m_{\alpha} Y_{\alpha}$$

$$Z = \dots$$

$$\dot{\vec{R}} = \frac{1}{M} \sum_{\alpha=1}^N m_{\alpha} \dot{\vec{r}}_{\alpha}$$

$$\vec{P} = M \dot{\vec{R}}$$

for continuous medium

$$m_{\alpha} \rightarrow dm$$

$$\vec{R} = \frac{1}{M} \int \vec{r} dm$$

volume

$$\vec{R} = \frac{1}{M} \int \vec{r} \rho dV$$

$$\rho = \frac{M}{V}$$

$$\vec{P} = \cancel{M} \left(\frac{1}{\cancel{M}} \sum_{\alpha} m_{\alpha} \dot{\vec{r}}_{\alpha} \right)$$

$$= \sum_{\alpha} m_{\alpha} \dot{\vec{r}}_{\alpha}$$

$$\vec{P} = \sum_{\alpha} \vec{p}_{\alpha} \quad \leftarrow \text{already knew from Ch. 1.}$$

$$\dot{\vec{P}} = M \ddot{\vec{R}} = \vec{F}_{\text{ext}} \quad \leftarrow \text{can treat a collection as a single particle at the center of mass}$$