


## IX. Fluid-particle interaction (FPI)



### IX.1 Fluid-to-particle transfer rates

#### IX.1.1 Momentum transfer – forces and moments on a particle

#### IX.1.2 Heat transfer

#### IX.1.3 Particle tracking numerics


Chap 9

By E. Amani

## Heat transfer

- DNS formulation:

$$\begin{aligned}
 & \overbrace{m_p \frac{d\bar{T}_p}{dt}}^{q_{in}} \\
 &= \underbrace{\int_{A_p} -\mathbf{q}_{\text{cond}}'' \cdot \mathbf{n} dA}_{q_{\text{conv}}} + \dot{m}_p (h_{v,\text{surf}} - \bar{h}_p + w' w' / 2) + q_R
 \end{aligned}$$



Chap 9

By E. Amani

## Heat transfer

- Point-particle formulation:

- Assuming **uniform temperature inside droplet**:

$$q_{\text{in}} = m_p \frac{d\bar{T}_p}{dt} = m_p \frac{di_p}{dt} = m_p c_p \frac{dT_p}{dt}$$

- Convection modeling:

$$\begin{aligned}
 q_{\text{conv}} &= h_m \underbrace{A_p}_{\pi d_p^2} (T_s - T_{\text{surf}}) = \pi d_p \lambda_m \text{Nu}_m (T_s - T_{\text{surf}}) \\
 \text{Nu}_m &= \frac{h_m d_p}{\lambda_m}
 \end{aligned}$$

- Therefore,

$$m_p c_p \frac{dT_p}{dt} = \pi d_p \lambda_m \text{Nu}_m \underbrace{(T_s - T_{\text{surf}})}_{\substack{T_p \\ \tau_M}} + \dot{m}_p \left( h_{v,\text{surf}} - \bar{h}_p + \underbrace{w' w' / 2}_{\substack{\text{neglected} \\ \text{By E. Amani}}} \right) + q_R$$

Chap 9

By E. Amani

## Heat transfer

- Finally:

$$\frac{dT_p}{dt} = \frac{T_s - T_p}{\tau_T} - \frac{1}{\tau_M} \frac{L_{v,\text{surf}}}{c_p} + \frac{q_R}{m_p c_p}; \quad \tau_T = \frac{\rho_p c_p d_p^2}{6 \lambda_m \text{Nu}_m}$$

Heat transfer time  
scale


- Different models, different correlations for  $\text{Nu}_m$ , e.g. [13, 14]
- See “FPISummary.pdf” for a short summary.

Chap 9

By E. Amani

## Heat transfer

- Application: internal combustion engines




▲ CI engine simulation- fuel atomization, **evaporation**, mixing, and combustion

Chap 9

By E. Amani


# Other topics

## ● Mass transfer




# Other topics

## ● Two-way coupling



## Other topics

### ● Dispersion



Chap 9

By E. Amani

## Numerics of particle tracking

The general form of ODE Lagrangian equations is:

$$\frac{d\Phi_p}{dt} = \frac{\Phi_s - \Phi_p}{\tau_\phi} + a \quad (74.9)$$

By proper selection of  $\Phi$ ,  $\tau_\phi$ ,  $a$ , the Lagrangian governing equations of a particle are obtained.

Several approaches have been proposed for the numerical solution of this equation

+ Implicit Euler:  $\mathcal{O}(\Delta t)$

$$\frac{\Phi_p^{n+1} - \Phi_p^n}{\Delta t} = \frac{\Phi_s^{n+1} - \Phi_p^{n+1}}{\tau_\phi^{n+1}} + a^{n+1} \rightarrow$$

$$\Phi_p^{n+1} = \left[ \Phi_p^n + \Delta t \left( a^{n+1} + \frac{\Phi_s^{n+1}}{\tau_\phi^{n+1}} \right) \right] / \left( 1 + \Delta t / \tau_\phi^{n+1} \right) \quad (75.9)$$

Chap 9

By E. Amani

## Numerics of particle tracking

+ Trapezoidal at  $n+\frac{1}{2}$   $O(\Delta t^2)$

$$\frac{\phi_p^{n+1} - \phi_p^n}{\Delta t} = \frac{\phi_s^{n+\frac{1}{2}} - \phi_p^{n+\frac{1}{2}}}{r_{\phi}^{n+\frac{1}{2}}} + a^{n+\frac{1}{2}} \rightarrow \frac{1}{2}(\phi_p^n + \phi_p^{n+1})$$

$$\phi_p^{n+1} = \left[ \phi_p^n \left(1 - \frac{\Delta t}{2 r_{\phi}^{n+\frac{1}{2}}}\right) + \frac{\Delta t}{r_{\phi}^{n+\frac{1}{2}}} \phi_s^{n+\frac{1}{2}} + \Delta t a^{n+\frac{1}{2}} \right] / \left(1 + \frac{\Delta t}{2 r_{\phi}^{n+\frac{1}{2}}}\right) \quad (76.9)$$

Chap 9

By E. Amani

## Numerics of particle tracking

Note 1:  $\phi_s^{n+1}$  is estimated by one of the following formula:

a)  $\phi_s^{n+1} \simeq \phi_s^n \quad (77.9)$

b)  $\phi_s^{n+1} \simeq \phi_s^n + (q\Delta t) \vec{U}_p \cdot (\vec{\nabla} \phi_s)^n \quad (78.9)$  needs  $\vec{\nabla} \phi_s$

c)  $\phi_s^{n+1} \simeq \phi_s(\vec{x}_p^{n+1})$ ,  $\vec{x}_p^{n+1} \simeq \vec{x}_p^n + q\Delta t \vec{U}_p^{(n)} \quad (79.9)$  inaccurate near the boundaries needs extrapolation outside domain  
 $\vec{x}_p^{n+1}$  approximation of particle location at  $n+1$

For  $a$  and  $r_{\phi}^{n+1}$ , the approximation like Eq. (77.9) is usually used.

Note 2: The position equation in these two methods is advanced by

$$\left[ \frac{d\vec{x}_p}{dt} = \vec{U}_p \right]^{n+\frac{1}{2}} \rightarrow \frac{\vec{x}_p^{n+1} - \vec{x}_p^n}{\Delta t} = \vec{U}_p^{n+\frac{1}{2}} = \frac{1}{2}(\vec{U}_p^n + \vec{U}_p^{n+1}) \rightarrow$$

$$\vec{x}_p^{n+1} = \vec{x}_p^n + \frac{\Delta t}{2}(\vec{U}_p^n + \vec{U}_p^{n+1}) \quad (80.9)$$

Chap 9

By E. Amani



## Numerics of particle tracking

+ Analytic method (or  $\Delta t$ )

Assuming  $\tau_p$ ,  $\phi_s$ , and  $a$  to be constant at  $t = \oplus$  where  $\oplus$  can be  $n, n+\frac{1}{2}$ , or  $n+1$ , Eq. (74.9) can be integrated analytically between  $t_n$  to  $t_{n+1}$  to yield (exercise)

$$\phi_p^{n+1} - \phi_p^n = [\phi_s^\oplus - \phi_p^n + a^\oplus \tau_p^\oplus] [1 - \exp(-\frac{\Delta t}{\tau_p^\oplus})] \quad (81.9)$$

then (exercise), inserting into  $d\vec{x}_p/dt = \vec{U}_p$  results in

$$\vec{x}_p^{n+1} = \vec{x}_p^n + \Delta t (\phi_s^\oplus + a^\oplus \tau_p^\oplus) - \tau_p^\oplus (\phi_p^{n+1} - \phi_p^n) \quad (82.9)$$

the values at  $\oplus$  can be estimated by Eqs. (77.9) - (79.9).


Chap 9

By E. Amani

## Project#2: ELSS of multiphase flows

### Part I

- Lagrangian particle tracking
- Python coding
- Validation against a simple case
- Particle in 2D vortex
- Error analysis of discretization schemes




Chap 9

By E. Amani

## Hands-on practice

### ● HW#9:

- Some useful analytical solutions for particle tracking
- Test of FLUENT Lagrangian particle tracking



**Chap 9**

**By E. Amani**

## References

1. Schiller, L. and A. Naumann, A drag coefficient correlation. Vdi Zeitung, 77: 318–320, 1935. Cité page. 38.
2. Sommerfeld, M., Theoretical and experimental modelling of particulate flows. Lecture series, 2000. 6: p. 3-7.
3. Auton, T., J. Hunt, and M. Prud'Homme, The force exerted on a body in inviscid unsteady non-uniform rotational flow. Journal of Fluid Mechanics, 1988. 197: p. 241-257.
4. Reeks, M. and S. McKee, The dispersive effects of Basset history forces on particle motion in a turbulent flow. The Physics of fluids, 1984. 27(7): p. 1573-1582.
5. Michaelides, E.E. and A. Roig, A reinterpretation of the Odar and Hamilton data on the unsteady equation of motion of particles. AIChE Journal, 2011. 57(11): p. 2997-3002.
6. Mei, R. and J. Klausner, Shear lift force on spherical bubbles. International journal of heat and fluid flow, 1994. 15(1): p. 62-65.

**Chap 9**

**By E. Amani**



## References (continued)

7. Rubinow, S. and J.B. Keller, The transverse force on a spinning sphere moving in a viscous fluid. *Journal of Fluid Mechanics*, 1961. 11(3): p. 447-459.
8. Oesterle, B. and T.B. Dinh, Experiments on the lift of a spinning sphere in a range of intermediate Reynolds numbers. *Experiments in Fluids*, 1998. 25(1): p. 16-22.
9. Li, A. and G. Ahmadi, Dispersion and deposition of spherical particles from point sources in a turbulent channel flow. *Aerosol science and technology*, 1992. 16(4): p. 209-226.
10. Ounis, H., G. Ahmadi, and J.B. McLaughlin, Brownian particle deposition in a directly simulated turbulent channel flow. *Physics of Fluids A: Fluid Dynamics*, 1993. 5(6): p. 1427-1432.
11. Brock, J.R., On the theory of thermal forces acting on aerosol particles. *Journal of Colloid Science*, 1962. 17(8): p. 768-780.

**Chap 9**

**By E. Amani**

## References (continued)

12. Tomiyama, A., Struggle with computational bubble dynamics. *Multiphase Science and Technology*, 1998. 10(4): p. 369-405.
13. Amani, E. and M. Nobari, A calibrated evaporation model for the numerical study of evaporation delay in liquid fuel sprays. *International journal of heat and mass transfer*, 2013. 56(1): p. 45-58.
14. Amani, E. and M. Nobari, Systematic tuning of dispersion models for simulation of evaporating sprays. *International Journal of Multiphase Flow*, 2013. 48: p. 11-31.
15. Behbahani, M.A., Edrisi, M., Rashidi, F. and Amani, E., Tuning a multi-fluid model for gas lift simulations in wells, *Chemical Engineering Research and Design*, 2012. 90(4): p. 471-486.
16. Tofighian, H., Amani, E. and Saffar-Avval, M., A large eddy simulation study of cyclones: The effect of sub-models on efficiency and erosion prediction, *Powder Technology*, 2020. 360, p. 1237-1252.
17. Amani, E., Mehrabian, M. and Movahed, S., A discrete phase hybrid continuum-atomistic model for electrokinetics in nanofluidics, *Physics of Fluids*, 2018. 30(7), p.072003.

**Chap 9**

**By E. Amani**

## References (continued)

18. Turns, S., An introduction to combustion, 2011, Third edition, McGraw-hill.

**Chap 9**

**By E. Amani**

