

Calculus A II One-to-One Tutoring

Chang, Yung-Hsuan

April 21, 2024

Example 3.1.

Compute all the first partial derivatives of $f(x, y) = x^3 + x^2y^2 - 2y^2$.

Recall 3.2 (Chain Rule).

If g is differentiable at x and f is differentiable at $g(x)$, then $(f \circ g)(x)$ is differentiable at x and

$$\frac{d}{dx} ((f \circ g)(x)) = \frac{df}{dx}(g(x)) \cdot \frac{dg}{dx}(x).$$

In Leibniz notation, if $y = f(u)$ and $u = g(x)$ are both differentiable, then

$$\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}.$$

Theorem 3.3 (Chain Rule).

Let $z = f(x, y)$, $x = x(t)$, $y = y(t)$. Then

$$\frac{dz}{dt} = \frac{\partial z}{\partial x} \cdot \frac{dx}{dt} + \frac{\partial z}{\partial y} \cdot \frac{dy}{dt}.$$

Example 3.4.

Let $z = x^2y + 3xy^4$ with $(x, y) = (\sin(2t), \cos(t))$. Compute $\frac{dz}{dt}$.

Example 3.5.

Let $u = x^4y + y^2z^3$ with $(x, y, z) = (rse^t, rs^2e^{-t}, r^2s \sin(t))$. Compute $\frac{\partial u}{\partial s}$.

Example 3.6.

Compute all first partial derivatives for $f(x, y) = e^{\sqrt{x^2+y}}$.

Example 3.7.

Let $z = x^2 + y^2$ with $(x, y) = (t - \cos t, 1 - \sin t)$. Compute $\frac{dz}{dt}$.

Example 3.8.

Compute all first partial derivatives for $f(x, y) = \arcsin\left(\frac{x^2 - y^2}{x^2 + y^2}\right)$ with $xy > 0$.

Example 3.9.

Let $z = e^{xy}$ with $(x, y) = \left(\ln \sqrt{u^2 + v^2}, \arctan \frac{v}{u} \right)$. Compute $\frac{\partial z}{\partial u}$.

Definition 3.10 (Directional Derivative).

Let f be a real-valued function of two variables. The directional derivative of f at (x_0, y_0) for the direction $\hat{\mathbf{u}} = (u_1, u_2)$ (a unit vector) is defined when the following limit exists:

$$D_{\hat{\mathbf{u}}}f(x_0, y_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + hu_1, y_0 + hu_2) - f(x_0, y_0)}{h}.$$

Remark 3.11.

In the previous definition, if $\hat{\mathbf{u}} = (1, 0)$, then

$$D_{\hat{\mathbf{u}}}f(x_0, y_0) = \frac{\partial f}{\partial x}(x_0, y_0).$$

Example 3.12.

Determine the directional derivative of

$$f(x, y) = \begin{cases} \frac{xy}{x^2 + y^2}, & \text{if } (x, y) \neq (0, 0); \\ 0, & \text{if } (x, y) = (0, 0), \end{cases}$$

for the direction $(1, 1)$ at $(0, 0)$.

Example 3.13.

Determine the directional derivative of $f(x, y) = xy^2 \arctan(z)$ for the direction $(1, 1, 1)$ at $(2, 1, 1)$.

Definition 3.14 (Limit).

The limit

$$\lim_{(x,y) \rightarrow (x_0,y_0)} f(x,y) = L$$

if for all $\varepsilon > 0$ (the greek letter ε stands for “error”), there exists a $\delta > 0$ (the greek letter δ stands for “difference”) such that $\sqrt{(x - x_0)^2 + (y - y_0)^2} < \delta$ implies $|f(x, y) - f(x_0, y_0)| < \varepsilon$.

Remark 3.15.

If there exist two path $C_1 : y = f_1(x)$ and $C_2 : y = f_2(x)$ such that

$$\lim_{\substack{(x,y) \rightarrow (x_0,y_0) \\ (x,y) \in C_1}} f(x,y) \neq \lim_{\substack{(x,y) \rightarrow (x_0,y_0) \\ (x,y) \in C_2}} f(x,y),$$

then the limit $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x,y)$ does not exist.

Example 3.16.

Determine the existence of $\lim_{(x,y) \rightarrow (0,0)} f(x,y)$ for each of the following functions:

1. $f(x,y) = \frac{x^2 - y^2}{x^2 + y^2}$;
2. $f(x,y) = \frac{xy}{x^2 + y^2}$;
3. $f(x,y) = \frac{x^2 y}{x^2 + y^2}$; and
4. $f(x,y) = \frac{x^{\frac{2}{3}} \sin y}{x^2 + y^2}$.

Example 3.17.

Determine $\lim_{(x,y) \rightarrow (0,0)} \frac{\sin(x^2 + y^2)}{x^2 + y^2}$.

Definition 3.18 (Continuity).

We say a function f is continuous at (x_0, y_0) if

$$\lim_{(x,y) \rightarrow (x_0, y_0)} f(x,y) = f(x_0, y_0).$$

We say a function is continuous if it is continuous at every point in its domain.

Example 3.19.

Determine the continuity of the function $f(x, y) = \frac{x^2 y}{x^2 + y^2}$ at $(0, 0)$.