

Homework 2 of Computational Mathematics

Chang, Yung-Hsuan

111652004

Department of Applied Mathematics

March 22, 2024

Problem 1. Use a fixed-point iteration method to determine a solution accurate to within 10^{-2} for $x^3 - x - 1 = 0$ on $[1, 2]$. Use $p_0 = 1$.

Solution. We want to find the solution to $f(x) = x$ with $f(x) = \sqrt{1 + \frac{1}{x}}$. Then, by calculator,

$$p_1 = \sqrt{2} = 1.414,$$

$$p_2 = 1.307,$$

$$p_3 = 1.329,$$

$$p_4 = 1.324,$$

which is accurate to within 10^{-2} for $x^3 - x - 1 = 0$ on $[1, 2]$.

□

Problem 2. Use Theorem 2.1 to find a bound for the number of iterations needed to achieve an approximation with accuracy 10^{-3} to the solution of $x^3 + x - 4 = 0$ lying in the interval $[1, 4]$. Find an approximation to the root with this degree of accuracy.

Solution. Let $f(x) = x^3 + x - 4$. Since $f \in C[1, 4]$ and $f(1) \cdot f(4) = (-2) \cdot 64 < 0$. The Bisection method generates a sequence $\{p_n\}_{n=1}^{\infty}$ approaches to a zero p of f with

$$|p_n - p| \leq \frac{4 - 1}{2^n}, \quad \text{when } n \geq 1.$$

Then,

$$\frac{3}{2^n} \leq 10^{-3} \implies n \geq \log_2(3000) > 11.$$

```

111652004_CM_HW2_P2.py > ...
1  end_point_1 = 1
2  end_point_2 = 4
3  tolerance = 0.001
4  maximum_number_of_iterations = 50
5  p_0 = 2.5
6  i = 1
7
8  def f(x):
9      return x**3 + x - 4
10
11  FA = f(end_point_1)
12
13  while i <= maximum_number_of_iterations:
14      p = end_point_1 + (end_point_2 - end_point_1) / 2
15      FP = f(p)
16      if (p == 0 or (end_point_2 - end_point_1) / 2 < tolerance):
17          print(f"p = {p} with {i} iterations.")
18          exit(0)
19      i = i + 1
20      if FA * FP > 0:
21          end_point_1 = p
22          FA = FP
23      else:
24          end_point_2 = p
25
26  print(f"Method failed after {maximum_number_of_iterations}.")

```

PROBLEMS OUTPUT DEBUG CONSOLE TERMINAL PORTS GITLENS COMMENTS

PS E:\Eiken\Visual Studio Code Git Sync\CM_HW2 & C:/Users/yungh/AppData/Local/Microsoft/WindowsApps/python3.11.exe "e
p = 1.378662109375 with 12 iterations.

Using the Bisection method, by Python, $p = 1.37866$.

□

Problem 3. The following four methods are preposed to compute $21^{1/3}$. Rank them in order, based on their apparent speed of convergence, assuming $p_0 = 1$.

a. $p_n = \frac{20p_{n-1} + \frac{21}{p_{n-1}^2}}{21}$

b. $p_n = p_{n-1} - \frac{p_{n-1}^3 - 21}{3p_{n-1}^2}$

c. $p_n = p_{n-1} - \frac{p_{n-1}^4 - 21p_{n-1}}{p_{n-1}^2 - 21}$

d. $p_n = \sqrt{\frac{21}{p_{n-1}}}$

Solution.

Problem 4. Use Theorem 2.3 to show that $g(x) = 2^{-x}$ has a unique fixed point on $\left[\frac{1}{3}, 1\right]$. Use fixed-point iteration to find an approximation to the fixed point accurate to within 10^{-4} . Use corollary 2.5 to estimate the number of iterations required to achieve 10^{-4} accuracy, and compare this theoretical estimate to the number actually needed.

Solution. We know that $g \in C\left[\frac{1}{3}, 1\right]$ and $g(x) \in [0.5, 0.9637] \subseteq \left[\frac{1}{3}, 1\right]$. Then g has at least a fixed point in $[a, b]$. Moreover, $g'(x)$ exists on $\left(\frac{1}{3}, 1\right)$. Choose $k = 0.7$. Then


$$\begin{aligned} \left| \frac{d}{dx} 2^{-x} \right| &= \ln 2 \cdot 2^{-x} \\ &< \ln 2 \cdot 2^{-0} \\ &= \ln 2 \\ &< k \end{aligned}$$

for all $x \in (0, \infty)$. Hence, $g'(x)$ exists on $\left(\frac{1}{3}, 1\right)$ and a positive $0 < k < 1$ exists with $|g'(x)| \leq k$ for all $x \in \left(\frac{1}{3}, 1\right)$. Then there exists exactly one fixed point in $\left[\frac{1}{3}, 1\right]$. It is known the assumption of Theorem 2.4 holds, i.e., $g'(x)$ exists on $\left(\frac{1}{3}, 1\right)$ and a positive $0 < k < 1$ exists with $|g'(x)| \leq k$ for all $x \in \left(\frac{1}{3}, 1\right)$. By Corollary 2.5,

$$|p_n - p| \leq 0.7^n \max\{0.6 - \frac{1}{3}, 1 - 0.6\}.$$

Then,

$$0.7^n \max\{0.6 - \frac{1}{3}, 1 - 0.6\} \leq 10^{-4} \implies n > 23.$$



The image shows a Visual Studio Code editor window with two tabs: '111652004_CM_HW2.tex U' and '111652004_CM_HW2_P4.py U'. The Python file is open, displaying a script for finding the root of a function $g(x) = 0.5x$ using the bisection method. The script starts with $p_0 = 0.6$, a tolerance of 0.0001 , and a maximum of 50 iterations. It defines $g(x)$ and enters a while loop that updates p until the tolerance is met or iterations are exhausted. The terminal output shows the script ran successfully, printing the result after 9 iterations.

```
HW2 > 111652004_CM_HW2_P4.py > ...
1  p_0 = 0.6
2  tolerance = 0.0001
3  maximum_number_of_iterations = 50
4  i = 1
5
6  def g(x):
7      return 0.5**x
8
9  while i <= maximum_number_of_iterations:
10     p = g(p_0)
11     if abs(p - p_0) < tolerance:
12         print(f"p = {p} with {i} iterations.")
13         exit(0)
14     i = i + 1
15     p_0 = p
16
17 print(f"Method failed after {maximum_number_of_iterations}.")
```

PROBLEMS OUTPUT DEBUG CONSOLE TERMINAL PORTS GITLENS COMMENTS

PS E:\Eiken\Visual Studio Code Git Sync\CM_HW> & C:/Users/yungh/AppData/Local/Microsoft/WindowsApps/python3.11.exe "e
p = 0.6412138835623649 with 9 iterations.

By Python, the number of iteration actually needed is 9, which is smaller due to my overestimation for error. □

Problem 5. Let A be a given positive constant and $g(x) = 2x - Ax^2$.

- a. Show that if fixed-point iteration converges to a nonzero limit, then the limit is $p = \frac{1}{A}$, so the inverse of a number can be found using only multiplications and subtractions.
- b. Find an interval about $\frac{1}{A}$ for which fixed-point iteration converges, provided p_0 is in that interval.

Solution.

Problem 6. Show that if A is any positive number, then the sequence defined by

$$x_n = \frac{1}{2}x_{n-1} + \frac{A}{2x_{n-1}}, \quad \text{for } n \geq 1,$$

converges to \sqrt{A} whenever $x_0 > 0$.

Solution.

Problem 7. Let $f(x) = -x^3 - \cos x$. With $p_0 = -1$ and $p_1 = 0$, find p_3 .

- a. Use the Secant method.
- b. Use the method of False Position.

Solution.

Problem 8. Problems involving the amount of money required to pay off a mortgage over a fixed period of time involve the formula

$$A = \frac{P}{i} (1 - (1 + i)^{-n}),$$

known as an *ordinary annuity equation*. In this equation, A is the amount of the mortgage, P is the amount of each payment, and i is the interest rate per period for the n payment periods. Suppose that a 30-year home mortgage in the amount of \$135,000 is needed and that the borrower can afford house payments of at most \$1000 per month. What is the maximal interest rate the borrower can afford to pay?

Solution.

Problem 9.

- a. Show that for any positive integer k , the sequence defined by $p_n = \frac{1}{n^k}$ converges linearly to $p = 0$.
- b. Show that the sequence $p_n = 10^{-2^n}$ converges quadratically to 0.

Solution.

Problem 10.

- a. The following sequences are linearly convergent. Generate the first five terms of the sequence $\{\hat{p}_n\}$ using Aitken's Δ^2 method.

$$p_0 = 0.5, \quad p_n = \cos(p_{n-1}), \quad n \geq 1$$

- b. Use Steffensen's method to find, to an accuracy of 10^{-4} , the root of $x^3 - x - 1 = 0$ that lies in $[1, 2]$.

Solution.

Problem 11. Given a polynomial $P(x) = x^3 - 5x^2 + 8x - 6$, do the following:

- a. Evaluate $P(2)$, $P'(2)$, $P(4)$, and $P'(4)$ by Horner's method.
- b. Find the root of $P(x)$ with error less than 0.00001 between $[2, 4]$ by using the Newton method with initial point $x_0 = 2$ and $x_0 = 4$. Determine which initial point may lead to the root.
- c. Deflate $P(x)$ into a quadratic polynomial by using the results in (b) and find the complex roots of $P(x)$.
- d. Perform one step of Muller's Method starting from initial $(0, P(0))$, $(1, P(1))$ and $(2, P(2))$.
- e. Implement a MATLAB code of Muller's Method to find the complex root within error less than 0.00001 and compare with the answer you find in (c).

Solution.