

# Homework 4 of Computational Mathematics

Chang, Yung-Hsuan

111652004

Department of Applied Mathematics

April 29, 2024

**Problem 1.** Use the following data and the knowledge that the first five derivatives of  $f$  are bounded on  $[1, 5]$  by 2, 3, 6, 12, and 23, respectively, to approximate  $f'(3)$  as accurately as possible. Find a bound for the error.

$x$	1	2	3	4	5
$f(x)$	2.4142	2.6734	2.8974	3.0976	3.2804

**Solution.** By the five-point midpoint formula,

$$\begin{aligned}
 f'(3) &= \frac{1}{12 \cdot 1} (f(1) - 8 \cdot f(2) + 8 \cdot f(4) - f(5)) + \frac{h^4}{30} \cdot f^{(5)}(\xi) \\
 &= \frac{1}{12} (2.4142 - 8 \cdot 2.6734 + 8 \cdot 3.0976 - 3.2804) + \frac{(0.1)^4}{30} \cdot f^{(5)}(\xi) \\
 &= 0.8773 + \frac{(0.1)^4}{30} \cdot f^{(5)}(\xi),
 \end{aligned}$$

where  $\xi \in (1, 5)$ . The bound for the error is

$$\begin{aligned}
 \left| \frac{(0.1)^4}{30} \cdot f^{(5)}(\xi) \right| &\leq \frac{(0.1)^4}{30} \cdot 23 \\
 &= 7.667 \cdot 10^{-5}.
 \end{aligned}$$

□

**Problem 2.** Let  $f(x) = 3xe^x - \cos x$ . Use the following data and equation (4.9) to approximate  $f''(1.3)$  with  $h = 0.1$  and with  $h = 0.01$ .

$x$	1.20	1.29	1.30	1.31	1.40
$f(x)$	11.59006	13.78176	14.04276	14.30741	16.86187

Compare your results to  $f''(1.3)$ .

**Solution.** We first deal with the case  $h = 0.1$ . By the second derivative midpoint formula,

$$\begin{aligned}
 f''(1.3) &= \frac{1}{(0.1)^2} (f(1.20) - 2 \cdot f(1.30) + f(1.40)) - \frac{(0.1)^2}{12} \cdot f^{(4)}(\xi_1) \\
 &= \frac{1}{0.01} \cdot (11.59006 - 2 \cdot 14.04276 + 16.86187) - \frac{0.01}{12} \cdot f^{(4)}(\xi_1) \\
 &= 36.641 - \frac{0.01}{12} \cdot f^{(4)}(\xi_1),
 \end{aligned}$$

where  $\xi_1 \in (1.20, 1.40)$ . We now deal with the case  $h = 0.01$ . Again by the second derivative midpoint formula,

$$\begin{aligned}
 f''(1.3) &= \frac{1}{(0.01)^2} (f(1.29) - 2 \cdot f(1.30) + f(1.31)) - \frac{(0.01)^2}{12} \cdot f^{(4)}(\xi_2) \\
 &= \frac{1}{0.0001} \cdot (13.78176 - 2 \cdot 14.04276 + 14.30741) - \frac{0.0001}{12} \cdot f^{(4)}(\xi_2) \\
 &= 36.5 - \frac{0.0001}{12} \cdot f^{(4)}(\xi_2),
 \end{aligned}$$

where  $\xi_2 \in (1.29, 1.31)$ . The actual value of  $f''(1.3)$  can be calculated as follows:

$$\begin{aligned}
 f'(x) &= 3e^x + 3xe^x + \sin x \\
 \implies f''(x) &= 6e^x + 3xe^x + \cos x \\
 \implies f''(1.3) &= 6e^{1.3} + 3 \cdot 1.3 \cdot e^{1.3} + \cos(1.3) \\
 &= 36.59354.
 \end{aligned}$$

The case with  $h = 0.01$  is closed to the true value, and the other is farther to the true value. □

**Problem 3.** Derive an  $\mathcal{O}(h^4)$  five-point formula to approximate  $f'(x_0)$  that uses  $f(x_0 - h)$ ,  $f(x_0)$ ,  $f(x_0 + h)$ ,  $f(x_0 + 2h)$ , and  $f(x_0 + 3h)$ . [Hint: Consider the expression  $A \cdot f(x_0 - h) + B \cdot f(x_0 + h) + C \cdot f(x_0 + 2h) + D \cdot f(x_0 + 3h)$ . Expand in fourth Taylor polynomials, and choose  $A$ ,  $B$ ,  $C$ , and  $D$ , appropriately.]

**Solution.**

**Problem 4.** The following data give approximations to the integral

$$M = \int_0^\pi \sin x \, dx.$$

$$N_1(h) = 1.570796, \quad N_1\left(\frac{h}{2}\right) = 1.896119, \quad N_1\left(\frac{h}{4}\right) = 1.974232, \quad N_1\left(\frac{h}{8}\right) = 1.993570.$$

Assuming  $M = N_1(h) + K_1h^2 + K_2h^4 + K_3h^6 + K_4h^8 + \mathcal{O}(h^{10})$ , construct an extrapolation table to determine  $N_4(h)$ .

**Solution.** By the formula for the  $\mathcal{O}(h^{2j})$  approximation

$$N_j(h) = N_{j-1}\left(\frac{h}{2}\right) + \frac{N_{j-1}\left(\frac{h}{2}\right) - N_{j-1}(h)}{4^{j-1} - 1},$$

we can have the following table:

$\mathcal{O}(h^2)$	$\mathcal{O}(h^4)$	$\mathcal{O}(h^6)$	$\mathcal{O}(h^8)$
1.570796			
1.896119	2.004560		
1.974232	2.000270	1.999984	
1.993570	2.000016	1.999837	1.999834

□

**Problem 5.** The forward-difference formula can be expressed as

$$f'(x_0) = \frac{1}{h} (f(x_0 + h) - f(x_0)) - \frac{h}{2} f''(x_0) - \frac{h^2}{6} f'''(x_0) + \mathcal{O}(h^3).$$

Use extrapolation to derive an  $\mathcal{O}(h^3)$  formula for  $f'(x_0)$ .

**Solution.**

**Problem 6.** Find the degree of precision of the quadrature formula

$$\int_{-1}^1 f(x) \, dx = f\left(-\frac{\sqrt{3}}{3}\right) + f\left(\frac{\sqrt{3}}{3}\right).$$

**Solution.** For  $f(x) = P_0(x) = 1$ , it is clear that

$$\int_{-1}^1 1 \, dx = f\left(-\frac{\sqrt{3}}{3}\right) + f\left(\frac{\sqrt{3}}{3}\right) = 2.$$

For  $f(x) = P_1(x) = x$ , it is also clear that

$$\int_{-1}^1 x \, dx = f\left(-\frac{\sqrt{3}}{3}\right) + f\left(\frac{\sqrt{3}}{3}\right) = 0.$$

For  $f(x) = P_2(x) = x^2$ ,

$$\int_{-1}^1 x^2 \, dx = \frac{2}{3}$$

and

$$f\left(-\frac{\sqrt{3}}{3}\right) + f\left(\frac{\sqrt{3}}{3}\right) = \frac{1}{3} + \frac{1}{3} = \frac{2}{3}.$$

For  $f(x) = P_3(x) = x^3$ , it is clear that

$$\int_{-1}^1 x^3 \, dx = f\left(-\frac{\sqrt{3}}{3}\right) + f\left(\frac{\sqrt{3}}{3}\right) = 0.$$

For  $f(x) = P_4(x) = x^4$ ,

$$\int_{-1}^1 x^4 \, dx = \frac{2}{5}$$

and

$$f\left(-\frac{\sqrt{3}}{3}\right) + f\left(\frac{\sqrt{3}}{3}\right) = \frac{1}{9} + \frac{1}{9} = \frac{2}{9}.$$

Hence, the degree of precision of this quadrature formula is 3.

□

**Problem 7.** Find the constants  $x_0$ ,  $x_1$ , and  $c_1$  so that the quadrature formula

$$\int_0^1 f(x) \, dx = \frac{1}{2}f(x_0) + c_1f(x_1)$$

has the highest possible degree of precision.

**Solution.** Set  $f(x) = P_0(x) = 1$ . Then

$$1 = \frac{1}{2} + c_1.$$

This implies that  $c_1$  must be  $\frac{1}{2}$ . Set  $f(x) = P_1(x) = x$ . Then

$$\frac{1}{2} = \frac{1}{2}x_0 + \frac{1}{2}x_1.$$

Set  $f(x) = P_2(x) = x^2$ . Then

$$\frac{1}{3} = \frac{1}{2}x_0^2 + \frac{1}{2}x_1^2.$$

Since there are two unknowns and two equations, two unknowns may be solved:

$$\frac{1}{3} = \frac{1}{2}(1 - x_1)^2 + \frac{1}{2}x_1^2$$

$$2 = 3(1 - x_1)^2 + 3x_1^2$$

$$x_1 = \frac{3 \pm \sqrt{3}}{6}.$$

Choose  $x_0 < x_1$ . Then  $(x_0, x_1, c_1) = \left(\frac{3 - \sqrt{3}}{6}, \frac{3 + \sqrt{3}}{6}, \frac{1}{2}\right)$ . □



**Problem 8.** Determine the values of  $n$  and  $h$  required to approximate

$$\int_0^2 e^{2x} \sin 3x \, dx$$

to within  $10^{-4}$ . Use

- a. composite trapezoidal rule;
- b. composite Simpson's rule; and
- c. composite midpoint rule.

**Solution.**

**Problem 9.** Determine to within  $10^{-6}$  the length of the graph of the ellipse with  $4x^2 + 9y^2 = 36$ .

**Solution.**

**Problem 10.** Show that the error  $E(f)$  for composite Simpson's rule can be approximated by

$$-\frac{h^4}{180} (f'''(b) - f'''(a)) .$$

[Hint:  $\sum_{j=1}^{n/2} f^{(4)}(\xi_j)(2h)$  is a Riemann sum for  $\int_a^b f^{(4)}(x) \, dx$ .]

**Solution.**

**Problem 11.** Use the following data to approximate  $\int_1^5 f(x) \, dx$  as accurately as possible.

$x$	1	2	3	4	5
$f(x)$	2.4142	2.6734	2.8974	3.0976	3.2804

**Solution.**

**Problem 12.** Show that the approximation obtained from  $R_{k,2}$  is the same as that given by the composite Simpson's rule described in Theorem 4.4 with  $h = h_k$ .

**Solution.**

**Problem 13.** Use composite Simpson's rule with  $n = 4, 6, 8, \dots$ , until successive approximations to the following integrals agree to within  $10^{-6}$ . Determine the number of nodes required. Use the adaptive quadrature algorithm to approximate the integral to within  $10^{-6}$ , and count the number of nodes. Did the adaptive quadrature produce any improvement?

a.  $\int_0^\pi x \sin(x^2) \, dx$ ; and

b.  $\int_0^\pi x^2 \sin x \, dx$ .

**Solution.**

**Problem 14.** Approximate the following integral using Gaussian quadrature with  $n = 2, 3, 4$ . Compare your answers with the exact values of the integral.

$$\int_1^{1.6} \frac{2x}{x^2 - 4} dx.$$

**Solution.** We first transform the interval  $[1, 1.6]$  to  $[-1, 1]$  by using  $t = \frac{2x - 2.6}{0.6}$ . Hence the integral becomes

$$\int_{-1}^1 \frac{1.2t + 5.2}{\left(\frac{0.6t + 2.6}{2}\right)^2 - 4} \cdot \frac{0.6}{2} dt = \int_{-1}^1 \frac{12t + 52}{3t^2 + 26t - 77} dt.$$

We now deal with the case  $n = 2$ . Using Table 4.12, the approximation is

$$1 \cdot \frac{12 \cdot 0.5773502692 + 52}{(0.5773502692)^2 + 26 \cdot 0.5773502692 - 77} + 1 \cdot \frac{12 \cdot (-0.5773502692) + 52}{(-0.5773502692)^2 + 26 \cdot (-0.5773502692) - 77} \\ = -1.4474.$$

The exact value of the integral is

$$\int_1^{1.6} \frac{2x}{x^2 - 4} dx = \int_{-3}^{-1.44} \frac{1}{u} du \\ = 0.364643 - 1.098612 \\ = -0.733969.$$

**Problem 15.** Determine constants  $a$ ,  $b$ ,  $c$ , and  $d$  that will produce a quadrature formula

$$\int_{-1}^1 f(x) \, dx = a \cdot f(-1) + b \cdot f(1) + c \cdot f'(-1) + d \cdot f'(1).$$

that has degree of precision 3.

**Solution.** Set  $f(x) = P_0(x) = 1$ . Then

$$2 = a + b.$$

Set  $f(x) = P_1(x) = x$ . Then

$$0 = -a + b + c + d.$$

Set  $f(x) = P_2(x) = x^2$ . Then

$$\frac{2}{3} = a + b - 2c + 2d.$$

Set  $f(x) = P_3(x) = x^3$ . Then

$$0 = -a + b + 3c + 3d.$$

Hence,  $(a, b, c, d) = (1, 1, 1/3, -1/3)$ .

□



**Problem 16.** The improper integral

$$\int_0^{\infty} f(x) \, dx$$

cannot be converted into an integral with finite limits using the substitution  $t = \frac{1}{x}$  because the limit at zero becomes infinite. The problem is resolved by first writing

$$\int_0^{\infty} f(x) \, dx = \int_0^1 f(x) \, dx + \int_1^{\infty} f(x) \, dx.$$

Apply this technique to approximate the following improper integrals to within  $10^{-6}$ .

a.  $\int_0^{\infty} \frac{1}{1+x^4} \, dx$ ; and

b.  $\int_0^{\infty} \frac{1}{(1+x^2)^3} \, dx$ .

**Solution.**