

Université Paris Cité – LIPADE

Algorithmic Complexity

Introduction

Jean-Guy Mailly (jean-guy.mailly@u-paris.fr)

2022



- ▶ Jean-Guy Mailly : jean-guy.mailly@u-paris.fr
Bureau 814 I
- ▶ 15h de cours : lundi, 10h15–11h45, Curie A
- ▶ 15h de TD : lundi, 12h45–14h15, Cuneo D (RSA/DCI)
vendredi, 10h30–12h00, Vieussens D (IAD/VMI)
- ▶ Modalités de contrôle de connaissances :
 - ▶ Un contrôle continu vers la mi-semestre
 - ▶ Un examen en fin de semestre
 - ▶ Note finale : $\max(EX, \frac{EX+CC}{2})$
- ▶ Moodle: Cours IFFAX020 Complexité Algorithmique
<https://moodle.u-paris.fr/course/view.php?id=5486>



Algorithmic Complexity

Measure the hardness of a problem w.r.t. the efficiency of algorithms to solve it

- ▶ Time
- ▶ Space

Goal of the Course

- ▶ Bases of complexity theory
- ▶ Main complexity classes (time and space)
- ▶ Complexity of usual problems
- ▶ Being able to determine the complexity of a problem



When solving a problem, two kinds of resources are used

- ▶ time
 - ▶ number of seconds
 - ▶ number of steps for the computation
- ▶ space
 - ▶ number of bytes used to execute the program
 - ▶ number of variables used to represent and solve the problem

Complexity Theory

- ▶ Classification of problems w.r.t. the resources required to solve them
 - ▶ The more we need time and/or space, the harder it is
- ▶ Comparison of problems (depending on the class they belong)
- ▶ Solving problems by translating them into other problems (with the same complexity)



Mappings and Asymptotic Bounds

Problems and Languages

Graph Theory

Non-Directed Graphs

Directed Graphs

Logic



Integer mappings

We use mappings $f : \mathbb{N} \rightarrow \mathbb{N}$ to represent the time (or space) used to solve a problem

- ▶ Intuition: If the problem entry has size n , $f(n)$ steps are required to compute the result
- ▶ If needed, we use the closest integer value (e.g. $\log(n)$ means $\lceil \log(n) \rceil$)



Integer mappings

We use mappings $f : \mathbb{N} \rightarrow \mathbb{N}$ to represent the time (or space) used to solve a problem

- ▶ Intuition: If the problem entry has size n , $f(n)$ steps are required to compute the result
- ▶ If needed, we use the closest integer value (e.g. $\log(n)$ means $\lceil \log(n) \rceil$)

Asymptotic bounds – \mathcal{O} notation

Given a mapping f , $\mathcal{O}(f(n))$ is the set of mappings g s.t. $\exists n_0, c$,


$$\forall n \geq n_0, g(n) \leq c \times f(n)$$

- ▶ Intuition: When n is large enough, g is smaller than f modulo some constant c

Example




- We suppose that we have an algorithm which solves a graphs problem in $g(n) = 4 \times n^2 + 3 \times n + 2$ steps, when n is the size of the graph (i.e. number of vertices)



Example




- ▶ We suppose that we have an algorithm which solves a graphs problem in $g(n) = 4 \times n^2 + 3 \times n + 2$ steps, when n is the size of the graph (i.e. number of vertices)
- ▶ When n becomes large enough, $g(n) \leq 5 \times f(n) = 5 \times n^2$, so $g(n) \in \mathcal{O}(n^2)$



Example



- ▶ We suppose that we have an algorithm which solves a graphs problem in $g(n) = 4 \times n^2 + 3 \times n + 2$ steps, when n is the size of the graph (i.e. number of vertices)
- ▶ When n becomes large enough, $g(n) \leq 5 \times f(n) = 5 \times n^2$, so $g(n) \in \mathcal{O}(n^2)$





From Slow to Fast Increase

C is an arbitrary constant, and \log is any logarithmic function

| Family of Functions | Name |
|----------------------------|-----------------|
| $\mathcal{O}(1)$ | Constant |
| $\mathcal{O}(\log(n))$ | Logarithmic |
| $\mathcal{O}((\log(n))^c)$ | Polylogarithmic |
| $\mathcal{O}(n)$ | Linear |
| $\mathcal{O}(n \log(n))$ | Linearithmic |
| $\mathcal{O}(n^2)$ | Quadratic |
| $\mathcal{O}(n^C)$ | Polynomial |
| $\mathcal{O}(C^n)$ | Exponential |
| $\mathcal{O}(n!)$ | Factorial |


On the Sum of Functions

If $f(n) = g(n) + h(n)$, $\mathcal{O}(f(n)) = \max(\mathcal{O}(g(n)), \mathcal{O}(h(n)))$

Explanation of the \mathcal{O} “Hierarchy”




► $\mathcal{O}(1) = \{g \mid \exists n_0, c, \forall n \geq n_0, g(n) \leq c \times 1\}$



Explanation of the \mathcal{O} “Hierarchy”




- ▶ $\mathcal{O}(1) = \{g \mid \exists n_0, c, \forall n \geq n_0, g(n) \leq c \times 1\}$
- ▶ $\mathcal{O}(\log(n)) = \{g \mid \exists n_0, c, \forall n \geq n_0, g(n) \leq c \times \log(n)\}$



Explanation of the \mathcal{O} “Hierarchy”




- ▶ $\mathcal{O}(1) = \{g \mid \exists n_0, c, \forall n \geq n_0, g(n) \leq c \times 1\}$
- ▶ $\mathcal{O}(\log(n)) = \{g \mid \exists n_0, c, \forall n \geq n_0, g(n) \leq c \times \log(n)\}$
- ▶ ...



Explanation of the \mathcal{O} “Hierarchy”



- ▶ $\mathcal{O}(1) = \{g \mid \exists n_0, c, \forall n \geq n_0, g(n) \leq c \times 1\}$
- ▶ $\mathcal{O}(\log(n)) = \{g \mid \exists n_0, c, \forall n \geq n_0, g(n) \leq c \times \log(n)\}$
- ▶ ...
- ▶ $\mathcal{O}(n!) = \{g \mid \exists n_0, c, \forall n \geq n_0, g(n) \leq c \times n!\}$





Mappings and Asymptotic Bounds

Problems and Languages

Graph Theory

Non-Directed Graphs

Directed Graphs

Logic



Decision Problem, Function Problem

A *decision problem* over a set of input data E is a mapping from any element in E to a value in $\{false, true\} (\simeq \{0, 1\} \simeq \{NO, YES\})$.

A *function problem* over a set of input data E is a mapping from any element in E to a single outcome.

Solving an Equation

Decision Problem Does the equation $f(x) = y$ have a solution?

Function Problem Give a solution of the equation $f(x) = y$



Enumeration Problem, Optimization Problem

Given a function problem \mathcal{P} ,

The *enumeration problem* $\text{ENUM-}\mathcal{P}$ over E is a mapping from any element $e_i \in E$ to the set of all outcome of \mathcal{P} over e_i .

The *optimization problem* $\text{OPT-}\mathcal{P}$ over E is a mapping from any element in E to a single outcome which minimizes a given criterion.

Solving an Equation

Enumeration Problem Give all the solutions of the equation $f(x) = y$

Optimization Problem Give a minimal solution of the equation
$$f(x) = y$$



Definition

- ▶ Set of symbols Σ called *vocabulary* or *alphabet*.
- ▶ A *word* w is a sequence of symbols $w_1 w_2 \dots w_k$, with $w_i \in \Sigma$ for all i .
- ▶ $\Sigma^* = \{w_1 w_2 \dots w_k \mid w_i \in \Sigma, k \in \mathbb{N}\}$
- ▶ a *language* \mathcal{L} is any subset of Σ^* , the *complement* of \mathcal{L} is $\bar{\mathcal{L}} = \Sigma^* \setminus \mathcal{L}$

Language \simeq Decision Problem

- ▶ For any language \mathcal{L} , $\mathcal{P}(\mathcal{L})$ is the decision problem:
Given $x \in \Sigma^*$, does x belong to \mathcal{L} ?
- ▶ For any decision problem \mathcal{P} , $\mathcal{L}(\mathcal{P})$ is the language:
 $\{x \in \text{instances of } \mathcal{P} \mid x \text{ is a positive instance of } \mathcal{P}\}$



Mappings and Asymptotic Bounds

Problems and Languages

Graph Theory

- Non-Directed Graphs

- Directed Graphs

Logic



Definition


A non-directed graph is a pair $G = \langle N, E \rangle$ where N is the set of **nodes** and $E \subseteq \text{pairs}(N)$ is the set of **edges**, with $\text{pairs}(N) = \{\{x_i, x_j\} \mid x_i, x_j \in N\}$



Definition

A non-directed graph is a pair $G = \langle N, E \rangle$ where N is the set of **nodes** and $E \subseteq \text{pairs}(N)$ is the set of **edges**, with $\text{pairs}(N) = \{\{x_i, x_j\} \mid x_i, x_j \in N\}$

$G = \langle N, E \rangle$, with $N = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$ and $E = \{\{x_1, x_2\}, \{x_2, x_3\}, \{x_3, x_4\}, \{x_4, x_5\}, \{x_5, x_6\}, \{x_5, x_7\}, \{x_6, x_7\}\}$






Definition

A **subgraph** in a non-directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq \text{pairs}(N') \cap E$. If $E' = \text{pairs}(N') \cap E$, we say that G' is **the subgraph of G induced by N'**

Definition

A **subgraph** in a non-directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq \text{pairs}(N') \cap E$. If $E' = \text{pairs}(N') \cap E$, we say that G' is **the subgraph of G induced by N'**


$G' = \langle \{x_1, x_2, x_3\}, \{\{x_1, x_2\}\} \rangle$ is a subgraph of G



Definition

A **subgraph** in a non-directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq \text{pairs}(N') \cap E$. If $E' = \text{pairs}(N') \cap E$, we say that G' is **the subgraph of G induced by N'**

$G' = \langle \{x_1, x_2, x_3\}, \{\{x_1, x_2\}\} \rangle$ is a subgraph of G




Definition

A **subgraph** in a non-directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq \text{pairs}(N') \cap E$. If $E' = \text{pairs}(N') \cap E$, we say that G' is **the subgraph of G induced by N'**

$G' = \langle \{x_1, x_2, x_3\}, \{\{x_1, x_2\}\} \rangle$ is a subgraph of G

$G' = \langle \{x_1, x_2, x_3\}, \{\{x_1, x_2\}, \{x_2, x_3\}\} \rangle$ is the subgraph of G induced by $\{x_1, x_2, x_3\}$




Definition

A **subgraph** in a non-directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq \text{pairs}(N') \cap E$. If $E' = \text{pairs}(N') \cap E$, we say that G' is **the subgraph of G induced by N'**

$G' = \langle \{x_1, x_2, x_3\}, \{\{x_1, x_2\}\} \rangle$ is a subgraph of G

$G' = \langle \{x_1, x_2, x_3\}, \{\{x_1, x_2\}, \{x_2, x_3\}\} \rangle$ is the subgraph of G induced by $\{x_1, x_2, x_3\}$





Definition


A chain in a non-directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, \{n_i, n_{i+1}\} \in E$.



Definition

A chain in a non-directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, \{n_i, n_{i+1}\} \in E$.

(x_3, x_4, x_5, x_6) and (x_3, x_4, x_5, x_7) are chains of G






Definition

A chain in a non-directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, \{n_i, n_{i+1}\} \in E$.

(x_3, x_4, x_5, x_6) and (x_3, x_4, x_5, x_7) are chains of G






Definition

A chain in a non-directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, \{n_i, n_{i+1}\} \in E$.

(x_3, x_4, x_5, x_6) and (x_3, x_4, x_5, x_7) are chains of G






Definition

A chain in a non-directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, \{n_i, n_{i+1}\} \in E$.

(x_3, x_4, x_5, x_6) and (x_3, x_4, x_5, x_7) are chains of G

(x_1, x_6) is not a chain of G




Definition

A chain in a non-directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, \{n_i, n_{i+1}\} \in E$.

(x_3, x_4, x_5, x_6) and (x_3, x_4, x_5, x_7) are chains of G

(x_1, x_6) is not a chain of G






Definition

A clique in a non-directed graph $G = \langle N, E \rangle$ is subgraph $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq \text{pairs}(N') \cap E$, such that $\forall x_i, x_j \in N', x_i \neq x_j, \{x_i, x_j\} \in E'$

Definition

A clique in a non-directed graph $G = \langle N, E \rangle$ is subgraph $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq \text{pairs}(N') \cap E$, such that $\forall x_i, x_j \in N', x_i \neq x_j, \{x_i, x_j\} \in E'$


$\{\{x_5, x_6, x_7\}, \{\{x_5, x_6\}, \{x_6, x_7\}, \{x_7, x_5\}\}\}$ is a clique in G



Definition

A clique in a non-directed graph $G = \langle N, E \rangle$ is subgraph $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq \text{pairs}(N') \cap E$, such that $\forall x_i, x_j \in N', x_i \neq x_j, \{x_i, x_j\} \in E'$

$\{\{x_5, x_6, x_7\}, \{\{x_5, x_6\}, \{x_6, x_7\}, \{x_7, x_5\}\}\}$ is a clique in G






Definition

A cycle in a non-directed graph $G = \langle N, E \rangle$ is a chain (n_1, \dots, n_k) such that $n_1 = n_k$. The length of the cycle is $k - 1$

Definition

A cycle in a non-directed graph $G = \langle N, E \rangle$ is a chain (n_1, \dots, n_k) such that $n_1 = n_k$. The length of the cycle is $k - 1$

(x_5, x_6, x_7, x_5) is a cycle of length 3 in G






Definition

A cycle in a non-directed graph $G = \langle N, E \rangle$ is a chain (n_1, \dots, n_k) such that $n_1 = n_k$. The length of the cycle is $k - 1$

(x_5, x_6, x_7, x_5) is a cycle of length 3 in G






Definition

A directed graph is a pair $G = \langle N, E \rangle$ where N is the set of **nodes** and $E \subseteq N \times N$ is the set of **edges**

Definition

A directed graph is a pair $G = \langle N, E \rangle$ where N is the set of **nodes** and $E \subseteq N \times N$ is the set of **edges**

$G = \langle N, E \rangle$, with $N = \{x_1, x_2, x_3, x_4, x_5, x_6, x_7\}$ and $E = \{(x_1, x_2), (x_2, x_3), (x_3, x_4), (x_4, x_3), (x_4, x_5), (x_5, x_6), (x_6, x_7), (x_7, x_5)\}$






Definition

A **subgraph** in a directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$. If $E' = (N' \times N') \cap E$, we say that G' is **the subgraph of G induced by N'**

Definition

A **subgraph** in a directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$. If $E' = (N' \times N') \cap E$, we say that G' is **the subgraph of G induced by N'**

$G' = \langle \{x_1, x_2, x_3\}, \{(x_1, x_2)\} \rangle$ is a subgraph of G






Definition

A **subgraph** in a directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$. If $E' = (N' \times N') \cap E$, we say that G' is **the subgraph of G induced by N'**

$G' = \langle \{x_1, x_2, x_3\}, \{(x_1, x_2)\} \rangle$ is a subgraph of G




Definition

A **subgraph** in a directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$. If $E' = (N' \times N') \cap E$, we say that G' is **the subgraph of G induced by N'**

$G' = \langle \{x_1, x_2, x_3\}, \{(x_1, x_2)\} \rangle$ is a subgraph of G

$G' = \langle \{x_1, x_2, x_3\}, \{(x_1, x_2), (x_2, x_3)\} \rangle$ is the subgraph of G induced by $\{x_1, x_2, x_3\}$




Definition

A **subgraph** in a directed graph $G = \langle N, E \rangle$ is a pair $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$. If $E' = (N' \times N') \cap E$, we say that G' is **the subgraph of G induced by N'**

$G' = \langle \{x_1, x_2, x_3\}, \{(x_1, x_2)\} \rangle$ is a subgraph of G

$G' = \langle \{x_1, x_2, x_3\}, \{(x_1, x_2), (x_2, x_3)\} \rangle$ is the subgraph of G induced by $\{x_1, x_2, x_3\}$





Definition


A path in a directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, (n_i, n_{i+1}) \in E$.



Definition

A path in a directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, (n_i, n_{i+1}) \in E$.

(x_3, x_4, x_5, x_6) is a path of G , but (x_3, x_4, x_5, x_7) is not






Definition

A path in a directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, (n_i, n_{i+1}) \in E$.

(x_3, x_4, x_5, x_6) is a path of G , but (x_3, x_4, x_5, x_7) is not






Definition

A path in a directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, (n_i, n_{i+1}) \in E$.


(x_3, x_4, x_5, x_6) is a path of G , but (x_3, x_4, x_5, x_7) is not



Definition

A path in a directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, (n_i, n_{i+1}) \in E$.

(x_3, x_4, x_5, x_6) is a path of G , but (x_3, x_4, x_5, x_7) is not
 (x_1, x_6) is not a path of G




Definition

A path in a directed graph $G = \langle N, E \rangle$ is a vector of nodes (n_1, \dots, n_k) such that $\forall 0 \leq i < k, (n_i, n_{i+1}) \in E$.

(x_3, x_4, x_5, x_6) is a path of G , but (x_3, x_4, x_5, x_7) is not

(x_1, x_6) is not a path of G






Definition

A clique in a directed graph $G = \langle N, E \rangle$ is subgraph $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$, such that $\forall x_i, x_j \in N', x_i \neq x_j, (x_i, x_j) \in E'$

Definition

A clique in a directed graph $G = \langle N, E \rangle$ is subgraph $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$, such that $\forall x_i, x_j \in N', x_i \neq x_j, (x_i, x_j) \in E'$


$\langle \{x_5, x_6, x_7\}, \{(x_5, x_6), (x_6, x_7), (x_7, x_5)\} \rangle$ is not a clique in G



Definition

A clique in a directed graph $G = \langle N, E \rangle$ is subgraph $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$, such that $\forall x_i, x_j \in N', x_i \neq x_j, (x_i, x_j) \in E'$


$\langle \{x_5, x_6, x_7\}, \{(x_5, x_6), (x_6, x_7), (x_7, x_5)\} \rangle$ is not a clique in G



Definition

A clique in a directed graph $G = \langle N, E \rangle$ is subgraph $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$, such that $\forall x_i, x_j \in N', x_i \neq x_j, (x_i, x_j) \in E'$

$\langle \{x_5, x_6, x_7\}, \{(x_5, x_6), (x_6, x_7), (x_7, x_5)\} \rangle$ is not a clique in G
 $\langle \{x_3, x_4\}, \{(x_3, x_4), (x_4, x_3)\} \rangle$ is a clique in G




Definition

A clique in a directed graph $G = \langle N, E \rangle$ is subgraph $G' = \langle N', E' \rangle$, with $N' \subseteq N$ and $E' \subseteq (N' \times N') \cap E$, such that $\forall x_i, x_j \in N', x_i \neq x_j, (x_i, x_j) \in E'$

$\langle \{x_5, x_6, x_7\}, \{(x_5, x_6), (x_6, x_7), (x_7, x_5)\} \rangle$ is not a clique in G

$\langle \{x_3, x_4\}, \{(x_3, x_4), (x_4, x_3)\} \rangle$ is a clique in G





Definition


A cycle in a directed graph $G = \langle N, E \rangle$ is a path (n_1, \dots, n_k) such that $n_1 = n_k$. The length of the cycle is $k - 1$

Definition

A cycle in a directed graph $G = \langle N, E \rangle$ is a path (n_1, \dots, n_k) such that $n_1 = n_k$. The length of the cycle is $k - 1$

(x_3, x_4, x_3) is a cycle of length 2 in G


(x_5, x_6, x_7, x_5) is a cycle of length 3 in G



Definition

A cycle in a directed graph $G = \langle N, E \rangle$ is a path (n_1, \dots, n_k) such that $n_1 = n_k$. The length of the cycle is $k - 1$

(x_3, x_4, x_3) is a cycle of length 2 in G
 (x_5, x_6, x_7, x_5) is a cycle of length 3 in G




Definition

A cycle in a directed graph $G = \langle N, E \rangle$ is a path (n_1, \dots, n_k) such that $n_1 = n_k$. The length of the cycle is $k - 1$

(x_3, x_4, x_3) is a cycle of length 2 in G

(x_5, x_6, x_7, x_5) is a cycle of length 3 in G




Definition

A cycle in a directed graph $G = \langle N, E \rangle$ is a path (n_1, \dots, n_k) such that $n_1 = n_k$. The length of the cycle is $k - 1$

(x_3, x_4, x_3) is a cycle of length 2 in G

(x_5, x_6, x_7, x_5) is a cycle of length 3 in G

(x_2, x_3, x_4, x_3) is not a cycle of G




Definition

A cycle in a directed graph $G = \langle N, E \rangle$ is a path (n_1, \dots, n_k) such that $n_1 = n_k$. The length of the cycle is $k - 1$

(x_3, x_4, x_3) is a cycle of length 2 in G

(x_5, x_6, x_7, x_5) is a cycle of length 3 in G

(x_2, x_3, x_4, x_3) is not a cycle of G





Mappings and Asymptotic Bounds

Problems and Languages

Graph Theory

Non-Directed Graphs

Directed Graphs

Logic



Definition

Propositional logic is a mathematical formalism which studies **propositions**, i.e. logical objects which can be **true** or **false**. A propositional formula is the combinations of propositions with **connectives**



Definition

Propositional logic is a mathematical formalism which studies **propositions**, i.e. logical objects which can be **true** or **false**. A propositional formula is the combinations of propositions with **connectives**

Examples of propositions:

- ▶ it's raining
- ▶ it's sunny
- ▶ I need an umbrella
- ▶ I need sunscreen



Definition

Propositional logic is a mathematical formalism which studies **propositions**, i.e. logical objects which can be **true** or **false**. A propositional formula is the combinations of propositions with **connectives**

Examples of propositions:

- ▶ it's raining
- ▶ it's sunny
- ▶ I need an umbrella
- ▶ I need sunscreen

Propositions are usually represented by *Boolean variables*, i.e. mathematical objects which can be assigned a value from a binary set $\mathbb{B} = \{0, 1\}$ (sometimes written $\{\text{False}, \text{True}\}$ or $\{\perp, \top\}$)

Syntax of Propositional Logic



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables

atom $\forall x_i \in V$, x_i is a (well-formed) formula

negation if φ is a formula, then $\neg\varphi$ is a formula

conjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \wedge \varphi_2$ is a formula

Syntax of Propositional Logic



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables

atom $\forall x_i \in V$, x_i is a (well-formed) formula

negation if φ is a formula, then $\neg\varphi$ is a formula

conjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \wedge \varphi_2$ is a formula

disjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \vee \varphi_2$ is a formula

Syntax of Propositional Logic



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables

atom $\forall x_i \in V$, x_i is a (well-formed) formula

negation if φ is a formula, then $\neg\varphi$ is a formula

conjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \wedge \varphi_2$ is a formula

disjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \vee \varphi_2$ is a formula

implication if φ_1 and φ_2 are formulas, then $\varphi_1 \Rightarrow \varphi_2$ is a formula

Syntax of Propositional Logic



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables

atom $\forall x_i \in V$, x_i is a (well-formed) formula

negation if φ is a formula, then $\neg\varphi$ is a formula

conjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \wedge \varphi_2$ is a formula

disjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \vee \varphi_2$ is a formula

implication if φ_1 and φ_2 are formulas, then $\varphi_1 \Rightarrow \varphi_2$ is a formula

equivalence if φ_1 and φ_2 are formulas, then $\varphi_1 \Leftrightarrow \varphi_2$ is a formula

Syntax of Propositional Logic



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables

atom $\forall x_i \in V$, x_i is a (well-formed) formula

negation if φ is a formula, then $\neg\varphi$ is a formula

conjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \wedge \varphi_2$ is a formula

disjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \vee \varphi_2$ is a formula

implication if φ_1 and φ_2 are formulas, then $\varphi_1 \Rightarrow \varphi_2$ is a formula

equivalence if φ_1 and φ_2 are formulas, then $\varphi_1 \Leftrightarrow \varphi_2$ is a formula

We use parentheses to simplify the reading and clarify priorities

Syntax of Propositional Logic



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables

atom $\forall x_i \in V$, x_i is a (well-formed) formula

negation if φ is a formula, then $\neg\varphi$ is a formula

conjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \wedge \varphi_2$ is a formula

disjunction if φ_1 and φ_2 are formulas, then $\varphi_1 \vee \varphi_2$ is a formula

implication if φ_1 and φ_2 are formulas, then $\varphi_1 \Rightarrow \varphi_2$ is a formula

equivalence if φ_1 and φ_2 are formulas, then $\varphi_1 \Leftrightarrow \varphi_2$ is a formula

We use parentheses to simplify the reading and clarify priorities

Formulas

- ▶ $x_1, \neg x_1, \neg\neg x_1$
- ▶ $x_1 \wedge x_2, x_1 \vee x_2$
- ▶ $x_1 \wedge (x_3 \vee (x_4 \Rightarrow x_1))$
- ▶ $\varphi \wedge \psi$

Not Formulas

- ▶ $x_1 \neg, \neg x_1 \neg$
- ▶ $x_1 \wedge, \vee x_1$
- ▶ $x_1 \Rightarrow (\vee x_2)$
- ▶ $\wedge \psi$



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables. An interpretation ω is a mapping from V to \mathbb{B} . Interpretations can be extended to formulas:

atom if $\varphi = x_i$, then $\omega(\varphi) = \omega(x_i)$

negation if $\varphi = \neg\varphi_1$, then $\omega(\varphi) = 1 - \omega(\varphi_1)$

conjunction if $\varphi = \varphi_1 \wedge \varphi_2$, then $\omega(\varphi) = \min(\omega(\varphi_1), \omega(\varphi_2))$



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables. An interpretation ω is a mapping from V to \mathbb{B} . Interpretations can be extended to formulas:

atom if $\varphi = x_i$, then $\omega(\varphi) = \omega(x_i)$

negation if $\varphi = \neg\varphi_1$, then $\omega(\varphi) = 1 - \omega(\varphi_1)$

conjunction if $\varphi = \varphi_1 \wedge \varphi_2$, then $\omega(\varphi) = \min(\omega(\varphi_1), \omega(\varphi_2))$

disjunction if $\varphi = \varphi_1 \vee \varphi_2$, then $\omega(\varphi) = \max(\omega(\varphi_1), \omega(\varphi_2))$



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables. An interpretation ω is a mapping from V to \mathbb{B} . Interpretations can be extended to formulas:

atom if $\varphi = x_i$, then $\omega(\varphi) = \omega(x_i)$

negation if $\varphi = \neg\varphi_1$, then $\omega(\varphi) = 1 - \omega(\varphi_1)$

conjunction if $\varphi = \varphi_1 \wedge \varphi_2$, then $\omega(\varphi) = \min(\omega(\varphi_1), \omega(\varphi_2))$

disjunction if $\varphi = \varphi_1 \vee \varphi_2$, then $\omega(\varphi) = \max(\omega(\varphi_1), \omega(\varphi_2))$

implication if $\varphi = \varphi_1 \Rightarrow \varphi_2$, then $\omega(\varphi) = \omega(\neg\varphi_1 \vee \varphi_2)$



Let $V = \{x_1, \dots, x_n\}$ be a set of Boolean variables. An interpretation ω is a mapping from V to \mathbb{B} . Interpretations can be extended to formulas:

atom if $\varphi = x_i$, then $\omega(\varphi) = \omega(x_i)$

negation if $\varphi = \neg\varphi_1$, then $\omega(\varphi) = 1 - \omega(\varphi_1)$

conjunction if $\varphi = \varphi_1 \wedge \varphi_2$, then $\omega(\varphi) = \min(\omega(\varphi_1), \omega(\varphi_2))$

disjunction if $\varphi = \varphi_1 \vee \varphi_2$, then $\omega(\varphi) = \max(\omega(\varphi_1), \omega(\varphi_2))$

implication if $\varphi = \varphi_1 \Rightarrow \varphi_2$, then $\omega(\varphi) = \omega(\neg\varphi_1 \vee \varphi_2)$

equivalence if $\varphi = \varphi_1 \Leftrightarrow \varphi_2$, then $\omega(\varphi) = \omega((\varphi_1 \Rightarrow \varphi_2) \wedge (\varphi_2 \Rightarrow \varphi_1))$

Semantics of Propositional Logic

Example of Interpretation



$$V = \{x_1, x_2, x_3, x_4\}, \omega(x_1) = \omega(x_4) = 0, \omega(x_2) = \omega(x_3) = 1$$

$$\blacktriangleright \omega(\neg x_1) = 1 - \omega(x_1) = 1, \omega(\neg x_2) = 1 - \omega(x_2) = 0$$

Semantics of Propositional Logic

Example of Interpretation



$$V = \{x_1, x_2, x_3, x_4\}, \omega(x_1) = \omega(x_4) = 0, \omega(x_2) = \omega(x_3) = 1$$

$$\blacktriangleright \omega(\neg x_1) = 1 - \omega(x_1) = 1, \omega(\neg x_2) = 1 - \omega(x_2) = 0$$

Intuitively:

- ▶ the negation takes the opposite of the original value

Semantics of Propositional Logic

Example of Interpretation



$$V = \{x_1, x_2, x_3, x_4\}, \omega(x_1) = \omega(x_4) = 0, \omega(x_2) = \omega(x_3) = 1$$

- ▶ $\omega(\neg x_1) = 1 - \omega(x_1) = 1, \omega(\neg x_2) = 1 - \omega(x_2) = 0$
- ▶ $\omega(x_1 \wedge x_2) = \min(\omega(x_1), \omega(x_2)) = \min(0, 1) = 0$

Intuitively:

- ▶ the negation takes the opposite of the original value

Semantics of Propositional Logic

Example of Interpretation



$$V = \{x_1, x_2, x_3, x_4\}, \omega(x_1) = \omega(x_4) = 0, \omega(x_2) = \omega(x_3) = 1$$

- ▶ $\omega(\neg x_1) = 1 - \omega(x_1) = 1, \omega(\neg x_2) = 1 - \omega(x_2) = 0$
- ▶ $\omega(x_1 \wedge x_2) = \min(\omega(x_1), \omega(x_2)) = \min(0, 1) = 0$
- ▶ $\omega(x_2 \wedge x_3) = \min(\omega(x_2), \omega(x_3)) = \min(1, 1) = 1$

Intuitively:

- ▶ the negation takes the opposite of the original value

Semantics of Propositional Logic

Example of Interpretation



$$V = \{x_1, x_2, x_3, x_4\}, \omega(x_1) = \omega(x_4) = 0, \omega(x_2) = \omega(x_3) = 1$$

- ▶ $\omega(\neg x_1) = 1 - \omega(x_1) = 1, \omega(\neg x_2) = 1 - \omega(x_2) = 0$
- ▶ $\omega(x_1 \wedge x_2) = \min(\omega(x_1), \omega(x_2)) = \min(0, 1) = 0$
- ▶ $\omega(x_2 \wedge x_3) = \min(\omega(x_2), \omega(x_3)) = \min(1, 1) = 1$

Intuitively:

- ▶ the negation takes the opposite of the original value
- ▶ the conjunction is true iff both values are true

Semantics of Propositional Logic

Example of Interpretation



$$V = \{x_1, x_2, x_3, x_4\}, \omega(x_1) = \omega(x_4) = 0, \omega(x_2) = \omega(x_3) = 1$$

- ▶ $\omega(\neg x_1) = 1 - \omega(x_1) = 1, \omega(\neg x_2) = 1 - \omega(x_2) = 0$
- ▶ $\omega(x_1 \wedge x_2) = \min(\omega(x_1), \omega(x_2)) = \min(0, 1) = 0$
- ▶ $\omega(x_2 \wedge x_3) = \min(\omega(x_2), \omega(x_3)) = \min(1, 1) = 1$
- ▶ $\omega(x_1 \vee x_2) = \max(\omega(x_1), \omega(x_2)) = \max(0, 1) = 1$

Intuitively:

- ▶ the negation takes the opposite of the original value
- ▶ the conjunction is true iff both values are true

Semantics of Propositional Logic

Example of Interpretation



$$V = \{x_1, x_2, x_3, x_4\}, \omega(x_1) = \omega(x_4) = 0, \omega(x_2) = \omega(x_3) = 1$$

- ▶ $\omega(\neg x_1) = 1 - \omega(x_1) = 1, \omega(\neg x_2) = 1 - \omega(x_2) = 0$
- ▶ $\omega(x_1 \wedge x_2) = \min(\omega(x_1), \omega(x_2)) = \min(0, 1) = 0$
- ▶ $\omega(x_2 \wedge x_3) = \min(\omega(x_2), \omega(x_3)) = \min(1, 1) = 1$
- ▶ $\omega(x_1 \vee x_2) = \max(\omega(x_1), \omega(x_2)) = \max(0, 1) = 1$
- ▶ $\omega(x_1 \vee x_4) = \max(\omega(x_1), \omega(x_4)) = \max(0, 0) = 0$

Intuitively:

- ▶ the negation takes the opposite of the original value
- ▶ the conjunction is true iff both values are true

Semantics of Propositional Logic

Example of Interpretation



$$V = \{x_1, x_2, x_3, x_4\}, \omega(x_1) = \omega(x_4) = 0, \omega(x_2) = \omega(x_3) = 1$$

- ▶ $\omega(\neg x_1) = 1 - \omega(x_1) = 1, \omega(\neg x_2) = 1 - \omega(x_2) = 0$
- ▶ $\omega(x_1 \wedge x_2) = \min(\omega(x_1), \omega(x_2)) = \min(0, 1) = 0$
- ▶ $\omega(x_2 \wedge x_3) = \min(\omega(x_2), \omega(x_3)) = \min(1, 1) = 1$
- ▶ $\omega(x_1 \vee x_2) = \max(\omega(x_1), \omega(x_2)) = \max(0, 1) = 1$
- ▶ $\omega(x_1 \vee x_4) = \max(\omega(x_1), \omega(x_4)) = \max(0, 0) = 0$

Intuitively:

- ▶ the negation takes the opposite of the original value
- ▶ the conjunction is true iff both values are true
- ▶ the disjunction is true iff at least one of the values is true



Sets of Atoms

- ▶ An interpretation can be written as the set of atoms which are assigned 1



Sets of Atoms

- ▶ An interpretation can be written as the set of atoms which are assigned 1
- ▶ Example: $V = \{x_1, x_2, x_3\}$, $\omega(x_1) = 0, \omega(x_2) = \omega(x_3) = 1$ is equivalent to $\omega = \{x_2, x_3\}$



Sets of Atoms

- ▶ An interpretation can be written as the set of atoms which are assigned 1
- ▶ Example: $V = \{x_1, x_2, x_3\}$, $\omega(x_1) = 0, \omega(x_2) = \omega(x_3) = 1$ is equivalent to $\omega = \{x_2, x_3\}$

Vectors of Bits

- ▶ An interpretation can be written as a vector of 0 and 1; the order has to be fixed (usually lexicographical order)



Sets of Atoms

- ▶ An interpretation can be written as the set of atoms which are assigned 1
- ▶ Example: $V = \{x_1, x_2, x_3\}$, $\omega(x_1) = 0, \omega(x_2) = \omega(x_3) = 1$ is equivalent to $\omega = \{x_2, x_3\}$

Vectors of Bits

- ▶ An interpretation can be written as a vector of 0 and 1; the order has to be fixed (usually lexicographical order)
- ▶ Example: $V = \{x_1, x_2, x_3\}$, $\omega(x_1) = 0, \omega(x_2) = \omega(x_3) = 1$ is equivalent to $\omega = 011$



- ▶ If $\omega(\varphi) = 1$, then ω is a **model** of φ . We also say that φ is **satisfied** by ω
- ▶ If $\omega(\varphi) = 0$, then ω is a counter-model of φ
- ▶ $\text{mod}(\varphi)$ is the set of models of φ
- ▶ If $\text{mod}(\varphi) = \emptyset$, then φ is **inconsistent** (or **unsatisfiable**)
- ▶ If $\text{mod}(\varphi) \neq \emptyset$, then φ is **consistent** (or **satisfiable**)
- ▶ If $\text{mod}(\varphi)$ is the set of all possible interpretations, then φ is **valid**



We define some meta-language symbols for reasoning about interpretations and formulas:

- ▶ $\omega \models \varphi$ means that (the interpretation) ω satisfies (the formula) φ , *i.e.* $\omega \in \text{mod}(\varphi)$
- ▶ $\varphi \vdash \psi$ means that (the formula) ψ is a consequence of (the formula) φ , formally defined as $\text{mod}(\varphi) \subseteq \text{mod}(\psi)$
- ▶ $\varphi \equiv \psi$ means that φ and ψ are equivalent, formally defined as $\varphi \vdash \psi$ and $\psi \vdash \varphi$ (which implies $\text{mod}(\varphi) = \text{mod}(\psi)$)



- ▶ A formula is a **literal** iff it is an atom or the negation of an atom
 x_1 and $\neg x_1$ are literals; $x_1 \vee x_2$ is not a literal



- ▶ A formula is a **literal** iff it is an atom or the negation of an atom
 x_1 and $\neg x_1$ are literals; $x_1 \vee x_2$ is not a literal
- ▶ A formula is a **clause** iff it is a disjunction of literals
 $x_1 \vee x_2 \vee \neg x_3$ is a clause ; $x_1 \vee (x_2 \wedge x_3)$ is not a clause



- ▶ A formula is a **literal** iff it is an atom or the negation of an atom
 x_1 and $\neg x_1$ are literals; $x_1 \vee x_2$ is not a literal
- ▶ A formula is a **clause** iff it is a disjunction of literals
 $x_1 \vee x_2 \vee \neg x_3$ is a clause ; $x_1 \vee (x_2 \wedge x_3)$ is not a clause
- ▶ A formula is a **cube** iff it is a conjunction of literals
 $x_1 \wedge x_2 \wedge \neg x_3$ is a cube ; $x_1 \wedge (x_2 \vee x_3)$ is not a cube



- ▶ A formula is a **literal** iff it is an atom or the negation of an atom
 x_1 and $\neg x_1$ are literals; $x_1 \vee x_2$ is not a literal
- ▶ A formula is a **clause** iff it is a disjunction of literals
 $x_1 \vee x_2 \vee \neg x_3$ is a clause ; $x_1 \vee (x_2 \wedge x_3)$ is not a clause
- ▶ A formula is a **cube** iff it is a conjunction of literals
 $x_1 \wedge x_2 \wedge \neg x_3$ is a cube ; $x_1 \wedge (x_2 \vee x_3)$ is not a cube

Conjunctive Normal Form

A formula is in Conjunctive Normal Form (CNF) iff it is a conjunction of clauses



- ▶ A formula is a **literal** iff it is an atom or the negation of an atom
 x_1 and $\neg x_1$ are literals; $x_1 \vee x_2$ is not a literal
- ▶ A formula is a **clause** iff it is a disjunction of literals
 $x_1 \vee x_2 \vee \neg x_3$ is a clause ; $x_1 \vee (x_2 \wedge x_3)$ is not a clause
- ▶ A formula is a **cube** iff it is a conjunction of literals
 $x_1 \wedge x_2 \wedge \neg x_3$ is a cube ; $x_1 \wedge (x_2 \vee x_3)$ is not a cube

Conjunctive Normal Form

A formula is in Conjunctive Normal Form (CNF) iff it is a conjunction of clauses

Disjunctive Normal Form

A formula is in Disjunctive Normal Form (DNF) iff it is a disjunction of cubes



Examples of CNF formulas

- ▶ $(x_1 \vee x_2) \wedge (x_3 \vee x_4)$
- ▶ $(x_1 \vee \neg x_2 \vee x_5) \wedge (x_3 \vee x_5)$

Examples of DNF formulas

- ▶ $(x_1 \wedge x_2) \vee (x_3 \wedge x_4)$
- ▶ $x_1 \vee (x_2 \wedge x_4) \vee (x_3 \wedge x_4 \wedge \neg x_5)$



- ▶ A clause is satisfied if at least one of its literals is satisfied
- ▶ A CNF formula is satisfied if all its clauses are satisfied

Example

With $\omega(x_1) = \omega(x_2) = 1$ and $\omega(x_3) = \omega(x_4) = 0$

- ▶ $(x_1 \vee x_2) \wedge (x_3 \vee x_4)$ is not satisfied



- ▶ A clause is satisfied if at least one of its literals is satisfied
- ▶ A CNF formula is satisfied if all its clauses are satisfied

Example

With $\omega(x_1) = \omega(x_2) = 1$ and $\omega(x_3) = \omega(x_4) = 0$

- ▶ $(x_1 \vee x_2) \wedge (x_3 \vee x_4)$ is not satisfied

With $\omega(x_1) = \omega(x_3) = 1$ and $\omega(x_2) = \omega(x_4) = 0$

- ▶ $(x_1 \vee x_2) \wedge (x_3 \vee x_4)$ is satisfied



- ▶ A cube is satisfied if all its literals are satisfied
- ▶ A DNF formula is satisfied if at least one of its cubes is satisfied

Example

With $\omega(x_1) = \omega(x_3) = 1$ and $\omega(x_2) = \omega(x_4) = 0$

- ▶ $(x_1 \wedge x_2) \vee (x_3 \wedge x_4)$ is not satisfied



- ▶ A cube is satisfied if all its literals are satisfied
- ▶ A DNF formula is satisfied if at least one of its cubes is satisfied

Example

With $\omega(x_1) = \omega(x_3) = 1$ and $\omega(x_2) = \omega(x_4) = 0$

- ▶ $(x_1 \wedge x_2) \vee (x_3 \wedge x_4)$ is not satisfied

With $\omega(x_1) = \omega(x_2) = 1$ and $\omega(x_3) = \omega(x_4) = 0$

- ▶ $(x_1 \wedge x_2) \vee (x_3 \wedge x_4)$ is satisfied