

Problem 1

a) For a function $f: \mathbb{R}^n \rightarrow \mathbb{R}$, the gradient is

$$\nabla f(x) = \begin{bmatrix} \frac{\partial f}{\partial x_1} \\ \frac{\partial f}{\partial x_2} \\ \vdots \\ \frac{\partial f}{\partial x_n} \end{bmatrix} = \left[\frac{\partial f}{\partial x} \right]^T$$

The direction of the gradient vector can be interpreted as pointing in the direction of steepest ascent.

b) For a function $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$, the Jacobian is

$$\frac{df}{dx} = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \dots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \dots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}$$

c) $f(x): \mathbb{R}^n \rightarrow \mathbb{R} \Leftrightarrow x \in \mathbb{R}^n, f \in \mathbb{R}$
Size of ∇f : $n \times 1$

d) $f(x): \mathbb{R}^n \rightarrow \mathbb{R}^m \Leftrightarrow x \in \mathbb{R}^n, f \in \mathbb{R}^m$
Size of $\frac{df}{dx}$: $m \times n$

Problem 2

$$f(x) = Ax = \begin{bmatrix} a_{11}x_1 + a_{12}x_2 \\ a_{21}x_1 + a_{22}x_2 \end{bmatrix}$$

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} \quad x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$a) \quad \frac{df}{dx} = \begin{bmatrix} \frac{df_1}{dx_1} & \frac{df_1}{dx_2} \\ \frac{df_2}{dx_1} & \frac{df_2}{dx_2} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \underline{\underline{A}}$$

This is the Jacobian

$$b) \quad \frac{d(Ax)}{dx} = A \frac{dx}{dx} = \underline{\underline{A}}$$

↑
move constants out

Problem 3

$$f(x, y) = x^T G y$$

$$x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}_{2 \times 1} \quad G = \begin{bmatrix} g_{11} & g_{12} & g_{13} \\ g_{21} & g_{22} & g_{23} \end{bmatrix}_{2 \times 3} \quad y = \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix}_{3 \times 1}$$

$$a) \quad f(x, y): (2 \times 1)^T \cdot 2 \times 3 \cdot 3 \times 1 = \underline{\underline{1 \times 1}} \Leftrightarrow \underline{\underline{\text{scalar}}}$$

$$\nabla_x f(x, y): \text{gradient wrt. } x$$

$$= \left[\frac{\partial f}{\partial x} \right]^T$$

$$\text{No, } \nabla_x f(x, y) = \left[\frac{\partial f(x, y)}{\partial x} \right]^T$$

$$b) f(x, y) = [x_1 \ x_2] G \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = [x_1 \ x_2] \begin{bmatrix} g_{11} y_1 + g_{12} y_2 + g_{13} y_3 \\ g_{21} y_1 + g_{22} y_2 + g_{23} y_3 \end{bmatrix}$$

$$= x_1 (g_{11} y_1 + g_{12} y_2 + g_{13} y_3) + x_2 (g_{21} y_1 + g_{22} y_2 + g_{23} y_3)$$

$$\nabla_x f(x, y) = \begin{bmatrix} \frac{\partial f}{\partial x_1} \\ \frac{\partial f}{\partial x_2} \end{bmatrix} = \begin{bmatrix} g_{11} y_1 + g_{12} y_2 + g_{13} y_3 \\ g_{21} y_1 + g_{22} y_2 + g_{23} y_3 \end{bmatrix} = \underline{\underline{G y}}$$

$$c) f(x, y) = y_1 (x_1 g_{11} + x_2 g_{21}) + y_2 (x_1 g_{12} + x_2 g_{22}) + y_3 (x_1 g_{13} + x_2 g_{23})$$

$$\nabla_y f(x, y) = \begin{bmatrix} \frac{\partial f}{\partial y_1} \\ \frac{\partial f}{\partial y_2} \\ \frac{\partial f}{\partial y_3} \end{bmatrix} = \begin{bmatrix} x_1 g_{11} + x_2 g_{21} \\ x_1 g_{12} + x_2 g_{22} \\ x_1 g_{13} + x_2 g_{23} \end{bmatrix} = \underline{\underline{x^T G}} = \underline{\underline{G^T x}}$$

$$d) f(x) = x^T H x \quad x \in \mathbb{R}^n \quad H \in \mathbb{R}^{n \times n}$$

$$\nabla f(x) = \nabla_{x^T} f(x) + \nabla_x f(x) = Hx + H^T x = \underline{\underline{(H + H^T)x}}$$

If H is symmetric: $H = H^T$

$$\Rightarrow \nabla f(x, y) = \underline{\underline{2Hx}}$$

Problem 4

Page 4/4

- $\mathcal{L}(\dots) \in \mathbb{R}$
- $G = G^T$ (symmetric)
- $x \in \mathbb{R}^n : n \times 1$

$$\begin{aligned}\mathcal{L}(x, \lambda, \mu) &= x^T G x + \lambda^T (Cx - d) + \mu^T (Ex - h) \\ &= x^T G x + \lambda^T C x - \lambda^T d + \mu^T E x - \mu^T h\end{aligned}$$

$$a) \nabla_x \mathcal{L}(x, \lambda, \mu) = \underline{\underline{2Gx + \lambda^T C + \mu^T E}}$$

$$b) \nabla_\lambda \mathcal{L}(x, \lambda, \mu) = \underline{\underline{Cx - d}}$$

$$c) \nabla_\mu \mathcal{L}(x, \lambda, \mu) = \underline{\underline{Ex - h}}$$