# Project 1 in FYS4150

### Eirik Ramsli Hauge

September 9, 2016

#### Abstract

This is a place to write something really deep. Like: The future is a really big place. Wow...

## Introduction

### Theory

$$\mathbf{A} \times \mathbf{v} = \mathbf{p}$$

$$\begin{pmatrix} b_1 & c_1 & 0 & 0 \\ a_2 & b_2 & c_2 & 0 \\ 0 & a_3 & b_3 & c_3 \\ 0 & 0 & a_4 & b_4 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \end{pmatrix} = \begin{pmatrix} p_1 \\ p_2 \\ p_3 \\ p_4 \end{pmatrix}$$

This gives us the following equations:

$$\begin{split} \mathrm{I} : v_1b_1 + c_1v_2 &= p_1 \\ \mathrm{II} : a_2v_1 + b_2v_2 + c_2v_3 &= p_2 \\ \mathrm{III} : a_3v_2 + b_3v_3 + c_3v_4 &= p_3 \\ \mathrm{IV} : a_4v_3 + b_4v_4 &= p_4 \end{split}$$

We want only zeroes on the left side of the diagonal. Therefore, we do as follows:

$$p_2^* = p_2 - p_1 \cdot \frac{a_2}{b_1}$$

$$= a_2 v_1 + b_2 v_2 + c_2 v_3 - v_1 a_2 - v_2 \frac{c_1 a_2}{b_1}$$

$$= v_2 (b_2 - \frac{c_1 a_2}{b_1}) + c_2 v_3$$

$$= v_2 b_2^* + c_2 v_3$$

Now we have the following matrix:

$$\hat{\mathbf{A}} = \begin{pmatrix} b_1 & c_1 & 0 & 0 \\ 0 & b_2^* & c_2 & 0 \\ 0 & a_2 & b_3 & c_3 \\ 0 & 0 & a_3 & b_4 \end{pmatrix}$$

As we can see, the a disappers from the second row. Just as we wanted. We do the same for the other rows

$$p_3^* = p_3 - p_2 \cdot \frac{a_3}{b_2^*}$$

$$= v_3(b_3 - \frac{c_2 a_3}{b_2^*}) + c_3 v_4$$

$$= v_3 b_3^* + c_3 v_3$$

General:

$$p_n^* = p_n - p_{n-1} \cdot \frac{a_n}{b_{n-1}^*} \tag{1}$$

$$b_n^* = b_n - \frac{c_{n-1}a_n}{b_{n-1}^*} \tag{2}$$

$$v_n = \frac{p_n^* - c_n v_{n+1}}{b_n^*} \tag{3}$$

**Programs** 

Results

Discussion

Conclusion