

BEILINSON MONAD

1. SETUP

We index homologically throughout. Fix a toric variety X with Cox ring $S = k[x_0, \dots, x_n]$ graded by $\text{Pic}(X) = \mathbb{Z}^{\oplus r}$. Let \mathcal{F} be a coherent sheaf on X . Write E for the Koszul dual of S . We equip E with a $\text{Pic}(X) \times \mathbb{Z}$ -grading such that $|e_i| = (-|x_i|, 1)$. Write $\omega := E^\vee$.

Remark 1.1. Before getting started, we record the following elementary observations. Of course, ω is an E -module with k -basis given by exterior polynomials in the e_i^* . Note that $|e_i^*| = (|x_i|, -1)$. The action of E on ω is by contraction. The $x_i \in S$ are also duals of the e_i , but we use different notation for the basis of ω to prevent confusion.

Another technical note: all E -modules are right modules. In particular, entries of matrices over E act on the right. This is also Macaulay2's convention. Note that this is the only way to make sense of the definition of the \mathbf{R} -functor in the EFS paper; if we apply the definition to a left E -module M , the maps in the complex $\mathbf{R}(M)$ are not E -linear. Nevertheless, sometimes we will multiply elements of E -modules on the left by elements of e (for instance, in the definition of the \mathbf{L} -functor below). When we do this, here is what we mean. When we write em for $e \in E$ and $m \in M$, where M is a right E -module, we mean $(-1)^{|e||m|}me$, where $|-|$ denotes the degree with respect to the second (standard) grading.

2. THE TATE RESOLUTION

Define $\mathcal{O}_{X \times E}$ to be the sheaf of algebras on X given by

$$U \mapsto \Gamma(U, \mathcal{O}_X) \otimes_k E.$$

Let $\text{Com}(X \times E)$ denote the category of complexes of $\mathcal{O}_{X \times E}$ -modules.

Example 2.1. Define an object $\kappa^*(\mathcal{F}) \in \text{Com}(X \times E)$ given by

$$\kappa^*(\mathcal{F})_i = \bigoplus_{\ell \in \text{Pic}(X)} \mathcal{F}(-\ell) \otimes \omega(\ell, -i)$$

with differential given by $\sum_i x_i \otimes e_i$.

Fix once and for all an affine open cover $\{V_0, \dots, V_t\}$ of X . Given a sheaf \mathcal{M} on X , denote by $\mathcal{C}^{\mathcal{M}}$ the Čech complex of \mathcal{M} corresponding to this open cover. Recall that we're indexing homologically, so the Čech complex is concentrated in nonpositive degrees.

Given $\mathcal{G} \in \text{Com}(X \times E)$, define a bicomplex as follows. The p^{th} column is given by $\mathcal{C}^{\mathcal{G}_p}$, but with the vertical differential multiplied by $(-1)^p$. The horizontal differential is induced by the differential on \mathcal{G} . Denote by $\tau_*(\mathcal{G}) \in \text{Com}(E)$ the direct sum totalization of this bicomplex. In other words, $\tau_*(\mathcal{G})$ is given as follows: take an explicit Čech model of the derived global sections of the complex \mathcal{G} , considered as a complex of \mathcal{O}_X -modules, and then remember the E -module structure on the terms of the resulting complex.

Definition 2.2. The *Tate resolution* of \mathcal{F} , denoted $\text{Tate}(\mathcal{F})$, is given by $(\tau_* \circ \kappa^*)(\mathcal{F})$.

Remark 2.3. Probably this isn't the right definition; instead, we want to replace this with a homotopy equivalent complex using Lemma 3.5 of EFS. But for the purpose of the Beilinson monad theorem, I think this distinction is irrelevant (assuming the \mathbf{U} -functor preserves homotopy).

Remark 2.4. Set

$$T(\mathcal{F}) := \bigoplus_{i=0}^t \bigoplus_{l \in \text{Pic}(X)} \mathcal{C}_{-i}^{\mathcal{F}(-l)} \otimes \omega(l, -i).$$

Unravelling the definitions, one sees that $\text{Tate}(\mathcal{F})$ has the form

$$\cdots \xrightarrow{\partial} T(0, -1) \xrightarrow{\partial} T \xrightarrow{\partial} T(0, 1) \xrightarrow{\partial} \cdots.$$

3. THE \mathbf{U} -FUNCTOR

We recall the definition of the functor

$$\mathbf{L} : \text{Com}(E) \rightarrow \text{Com}(S)$$

from Daniel's notes. For an E -module M concentrated in degree 0, $\mathbf{L}(M)$ is the complex with

$$\mathbf{L}(M)_q = \bigoplus_{d \in \text{Pic}(X)} M_{(-d, -q)} \otimes_k S(d)$$

and differential

$$(1) \quad m \otimes s \mapsto \sum_{i=0}^n e_i m \otimes x_i s.$$

For a general complex $(C, \partial) \in \text{Com}(E)$, we form the bicomplex

$$(2) \quad \begin{array}{ccccc} & & \downarrow & & \downarrow \\ \cdots & \xleftarrow{\partial} & \mathbf{L}(C_p)_q & \xleftarrow{\partial} & \mathbf{L}(C_{p+1})_q & \xleftarrow{\partial} \cdots \\ & & \downarrow & & \downarrow \\ \cdots & \xleftarrow{\partial} & \mathbf{L}(C_p)_{q-1} & \xleftarrow{\partial} & \mathbf{L}(C_{p+1})_{q-1} & \xleftarrow{\partial} \cdots \\ & & \downarrow & & \downarrow \end{array}$$

and apply $\text{Tot}^\oplus(-)$, where the vertical differential $\mathbf{L}(C_p)_q \rightarrow \mathbf{L}(C_p)_{q-1}$ is the dual Koszul map (1) multiplied by $(-1)^p$.

Let $\mathcal{L}(C)$ denote the bicomplex of \mathcal{O}_X -modules given by applying the associated sheaf functor to the bicomplex (2). Let $\mathcal{L}'(C)$ be the sub-bicomplex of $\mathcal{L}(C)$ given by taking summands of the form $C_{p,(-d,-q)} \otimes_k \mathcal{O}(d)$ with d effective. From now on, we'll write " $d \geq 0$ " for " d effective". Here, p denotes homological degree, and $(-d, -q)$ denotes internal degree. We define a functor

$$\mathbf{U} : \text{Com}(E) \rightarrow \text{Com}(\mathbb{P})$$

to be given by $C \mapsto \text{Tot}^\oplus(\mathcal{L}'(C))$.

♣♣♣ Michael: [Why Tot^\oplus and not Tot^Π ? Why $d \geq 0$ and not $d \leq 0$? Give clean conceptual explanation for definition of \mathbf{U} -functor.]

Proposition 3.1. *The above definition of the \mathbf{U} -functor agrees with Daniel's.*

Proof. Daniel's definition is given by

$$\omega(i, j) \mapsto \mathcal{L}(\omega_{\leq i})(i)[-j].$$

(We're abusing notation here by identifying the 1-column bicomplex $\mathcal{L}(\omega_{\leq i})(i)$ with its totalization.) We have

$$\begin{aligned} (\mathcal{L}(\omega_{\leq i})(i)[-j])_q &= \mathcal{L}(\omega_{\leq i})_{-j+q}(i) \\ &= \bigoplus_d (\omega_{\leq i})_{(-d, j-q)} \otimes \mathcal{O}(d+i) \\ &= \bigoplus_d (\omega_{\leq i})_{(i-d, j-q)} \otimes \mathcal{O}(d) \\ &= \bigoplus_{d \geq 0} (\omega_{\leq i})_{(i-d, j-q)} \otimes \mathcal{O}(d) \\ &= \bigoplus_{d \geq 0} \omega_{(i-d, j-q)} \otimes \mathcal{O}(d) \\ &= \mathbf{U}(\omega(i, j))_q. \end{aligned}$$

And of course the maps in both complexes are identical as well. □

Question 3.2. *Does the \mathbf{U} -functor preserve homotopy? Check this.*

4. BEILINSON MONAD

Recall that $\text{Tate}(\mathcal{F})$ is of the form

$$\cdots \rightarrow T(0, -1) \rightarrow T \rightarrow T(0, 1) \rightarrow \cdots,$$

where T is as described in Remark 2.4. The (p, q) term of $\mathcal{L}'(\text{Tate}(\mathcal{F}))$ is

$$\bigoplus_{d \geq 0} T(0, -p)_{(-d, -q)} \otimes \mathcal{O}(d) = \bigoplus_{d \geq 0} T_{(-d, -p-q)} \otimes \mathcal{O}(d),$$

and so

$$\mathbf{U}(\text{Tate}(\mathcal{F}))_m = \bigoplus_{p \in \mathbb{Z}} \bigoplus_{d \geq 0} T_{(-d, -m)} \otimes \mathcal{O}(d).$$

Equip each $\mathbf{U}(\text{Tate}(\mathcal{F}))_m$ with a $k[u]$ -module structure determined by the following “shift” operation: if $t = (\dots, t_{-1}, t_0, t_1, \dots) \in \bigoplus_{p \in \mathbb{Z}} \bigoplus_{d \geq 0} T_{(-d, -m)} \otimes \mathcal{O}(d)$,

$$u(t)_p = (-1)^{m-p} t_{p-1}.$$

Proposition 4.1. *The differential on $\mathbf{U}(\text{Tate}(\mathcal{F}))$ is $k[u]$ -linear.*

Proof. I'm writing down the proof to make sure I got the sign right in the definition of the u -action. We prove that the action of u commutes with both horizontal and vertical

differentials. Write d_T for the differential on $\text{Tate}(\mathcal{F})$ and d_K for the dual Koszul differential. We have

$$\begin{aligned} d_{\text{hor}}(u \cdot (\dots, t_{-1}, t_0, t_1, \dots)) &= d_{\text{hor}}(\dots, (-1)^{m-1}t_{-2}, (-1)^mt_{-1}, (-1)^{m-1}t_0, \dots) \\ &= (\dots, (-1)^md_T(t_{-3}), (-1)^{m-1}d_T(t_{-2}), (-1)^md_T(t_{-1}), \dots) \\ &= u \cdot (\dots, d_T(t_{-2}), d_T(t_{-1}), d_T(t_0), \dots) \\ &= u \cdot d_{\text{hor}}(\dots, t_{-1}, t_0, t_1, \dots), \end{aligned}$$

and

$$\begin{aligned} d_{\text{ver}}(u \cdot (\dots, t_{-1}, t_0, t_1, \dots)) &= d_{\text{ver}}(\dots, (-1)^{m-1}t_{-2}, (-1)^mt_{-1}, (-1)^{m-1}t_0, \dots) \\ &= (\dots, (-1)^md_K(t_{-2}), (-1)^md_K(t_{-1}), (-1)^md_K(t_0), \dots) \\ &= u \cdot (\dots, -d_K(t_{-1}), d_K(t_0), -d_K(t_1), \dots) \\ &= u \cdot d_{\text{ver}}(\dots, t_{-1}, t_0, t_1, \dots). \end{aligned}$$

□

So, $\mathbf{U}(\text{Tate}(\mathcal{F}))$ is a complex of $X \times \mathbb{A}^1$ -modules. In fact, since the action of u is invertible, $\mathbf{U}(\text{Tate}(\mathcal{F}))$ is a complex of $X \times \mathbb{G}_m$ -modules. Define

$$\mathbf{BM}(\mathcal{F}) := \mathbf{U}(\text{Tate}(\mathcal{F})) / (u - 1).$$

We have an isomorphism

$$(3) \quad \mathbf{BM}(\mathcal{F})_m \cong \bigoplus_{d \geq 0} T_{(-d, -m)} \otimes \mathcal{O}(d)$$

given by representing each class in $\mathbf{BM}(\mathcal{F})_m$ by an element concentrated in the $p = 0$ summand of $\bigoplus_{p \in \mathbb{Z}} \bigoplus_{d \geq 0} T_{(-d, -m)} \otimes \mathcal{O}(d)$. Via this isomorphism, the differential on $\mathbf{BM}(\mathcal{F})$ is given by $ud_T + d_K$, where d_T is the Tate differential and d_K is the dual Koszul differential.

Theorem 4.2. *The complex $\mathbf{BM}(\mathcal{F})$ is a monad with homology \mathcal{F} .*

Example 4.3. Take $X = \mathbb{P}(a)$, with a some positive integer, and take $\mathcal{F} = \mathcal{O}$. So X is a stacky point. If we set $T = \bigoplus_{i \in \mathbb{Z}} \omega(i, 0)$, then

$$\dots \xrightarrow{e} T(0, -1) \xrightarrow{e} T \xrightarrow{e} T(0, 1) \xrightarrow{e} \dots$$

is the Tate resolution of \mathcal{O} . We therefore have

$$\mathbf{U}(\text{Tate}(\mathcal{O}))_m = \bigoplus_{p \in \mathbb{Z}} \bigoplus_{d \geq 0} \bigoplus_{i \in \mathbb{Z}} \omega_{(i-d, -m)} \otimes \mathcal{O}(d) = \begin{cases} \bigoplus_{p \in \mathbb{Z}} \bigoplus_{d \geq 0} \omega_{(0,0)} \otimes \mathcal{O}(d), & m = 0; \\ \bigoplus_{p \in \mathbb{Z}} \bigoplus_{d \geq 0} \omega_{(a,-1)} \otimes \mathcal{O}(d), & m = 1; \\ 0, & \text{else.} \end{cases}$$

Taking coinvariants of the \mathbb{G}_m -action and applying the isomorphism (3) gives the complex

$$0 \rightarrow \bigoplus_{d \geq 0} \omega_{(a,-1)} \otimes \mathcal{O}(d) \xrightarrow{\begin{pmatrix} -1 & 0 & 0 & 0 & \dots \\ x & -1 & 0 & 0 & \dots \\ 0 & x & -1 & 0 & \dots \\ 0 & 0 & x & -1 & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}} \bigoplus_{d \geq 0} \omega_{(0,0)} \otimes \mathcal{O}(d) \rightarrow 0$$

whose homology is \mathcal{O} in degree 0 and 0 elsewhere, as expected.

5. PROOF OF THEOREM 4.2

Let $\mathcal{R} \in \text{Com}(X \times X)$ be Daniel's resolution of the diagonal. To prove Theorem 4.2, it suffices to give a homotopy equivalence

$$\mathbf{BM}(\mathcal{F}) \simeq \mathbf{R}\pi_{2*}(\pi_1^* \mathcal{F} \otimes \mathcal{R}).$$

The notation $\mathbf{R}\pi_{2*}$ is misleading: we are not working at the level of derived categories. Rather, we use the notation $\mathbf{R}\pi_{2*}$ to denote the Čech model for $\mathbf{R}\pi_{2*}$ induced by the affine open cover $\{V_0, \dots, V_t\}$ of X chosen above.

The rough idea is to define a map

$$\text{id} \times \overline{\mathbf{U}} : \text{Com}(X \times E) \dashrightarrow \text{Com}(X \times X)$$

such that the diagram

$$\begin{array}{ccccc} & & \text{Com}(X \times E) & \xrightarrow{\tau_*} & \text{Com}(E) \\ & \nearrow \kappa^* & \downarrow \text{id} \times \overline{\mathbf{U}} & & \downarrow \overline{\mathbf{U}} \\ \text{coh}(X) & & & & \\ & \searrow \pi_1^*(-) \otimes \mathcal{R} & \downarrow & \xrightarrow{\mathbf{R}\pi_{2*}} & \downarrow \\ & & \text{Com}(X \times X) & & \text{Com}(X) \end{array}$$

commutes up to homotopy, where $\overline{\mathbf{U}}$ denotes the functor given by applying the \mathbf{U} -functor and modding out by the relation $u - 1$, as discussed above. Here is how to define $\text{id} \times \overline{\mathbf{U}}$ on the image of κ^* (which is all we need). Recall that

$$\kappa^*(\mathcal{F})_i = \bigoplus_{\ell \in \text{Pic}(X)} \mathcal{F}(-\ell) \otimes \omega(\ell, -i)$$

and the differential is the dual Koszul map. We apply “ $\text{id} \times \mathbf{U}$ ” to $\kappa^*(\mathcal{F})$ to get the complex whose m^{th} term is

$$\bigoplus_{p \in \mathbb{Z}} \bigoplus_{d \geq 0} \bigoplus_{\ell \in \text{Pic}(X)} \omega_{(\ell-d, -m)} \otimes \mathcal{F}(-\ell) \boxtimes \mathcal{O}(d)$$

with differential $\sum_{i=0}^n e_i \otimes x_i + (-1)^p e_i \otimes y_i$. This complex has \mathbb{G}_m -action just as the Tate resolution does. Taking coinvariants and applying an isomorphism similar to (3), we arrive at the complex with m^{th} term

$$\bigoplus_{d \geq 0} \bigoplus_{\ell \in \text{Pic}(X)} \omega_{(\ell-d, -m)} \otimes \mathcal{F}(-\ell) \boxtimes \mathcal{O}(d)$$

and m^{th} differential $\sum_{i=0}^n (-1)^m e_i \otimes x_i + e_i \otimes y_i$.

Proposition 5.1. *$(\text{id} \times \overline{\mathbf{U}})(\kappa^*(\mathcal{O}))$ coincides with Daniel's resolution of the diagonal.*

Proof. Fill in. Should be easy. □

Example 5.2. Let's check this for $X = \mathbb{P}(a)$. Reading off the formula, we get

$$0 \rightarrow \bigoplus_{d \geq 0} \omega_{(a,-1)} \otimes \mathcal{O}(d, -(d+a)) \xrightarrow{\begin{pmatrix} y & 0 & 0 & 0 & \dots \\ -x & y & 0 & 0 & \dots \\ 0 & -x & y & 0 & \dots \\ 0 & 0 & -x & y & \dots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{pmatrix}} \bigoplus_{d \geq 0} \omega_{(0,0)} \otimes \mathcal{O}(d, -d) \rightarrow 0.$$

The homology is $\mathcal{O} \oplus \dots \oplus \mathcal{O}(a-1, -(a-1))$, and this is indeed the diagonal in this case (this is not entirely trivial to check). Notice this precisely recovers Daniel's "1-variable" example when $a = 1$.

Proposition 5.3. $(\mathbf{R}\pi_{2*} \circ (\text{id} \times \overline{\mathbf{U}}) \circ \kappa^*)(\mathcal{F}) = (\overline{\mathbf{U}} \circ \text{Tate})(\mathcal{F})$.

Proof. Let's start by computing the right hand side. As in Remark 2.4, let

$$T = \bigoplus_{i=0}^t \bigoplus_{l \in \text{Pic}(X)} \mathcal{C}_{-i}^{\mathcal{F}(-l)} \otimes \omega(l, -i),$$

so that

$$(\overline{\mathbf{U}} \circ \text{Tate})(\mathcal{F}) = \overline{\mathbf{U}}(\dots \rightarrow T(0, -1) \rightarrow T \rightarrow T(0, 1) \rightarrow \dots).$$

We have

$$\begin{aligned} (\overline{\mathbf{U}} \circ \text{Tate})(\mathcal{F})_m &= \bigoplus_{d \geq 0} T_{(-d, -m)} \otimes \mathcal{O}(d) \\ &= \bigoplus_{d \geq 0} \bigoplus_{i=0}^t \bigoplus_{l \in \text{Pic}(X)} \omega(l-d, -i-m) \otimes \mathcal{C}_{-i}^{\mathcal{F}(-l)} \otimes \mathcal{O}(d), \end{aligned}$$

and the differential is $ud_T + d_K$. On the other hand $(\mathbf{R}\pi_{2*} \circ (\text{id} \times \overline{\mathbf{U}}) \circ \kappa^*)(\mathcal{F})$ is the sheaf given as follows. Let W be an open set in X . We abuse notation slightly and write $\mathcal{C}^{\mathcal{O}(d)|_W}$ for the Čech complex on $\mathcal{O}(d)|_W$ corresponding to the open cover

$$\{V_0 \cap W, \dots, V_t \cap W\}.$$

We recall that the natural map

$$\mathcal{C}^{\mathcal{F}(-l)} \otimes \mathcal{C}^{\mathcal{O}(d)|_W} \rightarrow \mathcal{C}^{\mathcal{F}(-l) \boxtimes \mathcal{O}(d)|_W}$$

is a homotopy equivalence, where the target is the Čech complex associated to the open cover $\{V_i \times (V_j \cap W)\}_{0 \leq i, j \leq t}$. Form a bicomplex with p^{th} column given by

$$\bigoplus_{d \geq 0} \bigoplus_{l \in \text{Pic}(X)} \omega(l-d, -p) \otimes \mathcal{C}^{\mathcal{F}(-l)} \otimes \mathcal{C}^{\mathcal{O}(d)|_W},$$

with vertical differential multiplied by $(-1)^p$, and horizontal differential induced by the differential on $((\text{id} \times \overline{\mathbf{U}}) \circ \kappa^*)(\mathcal{F})$. Applying Tot^\oplus to this complex gives the value of the

$(\mathbf{R}\pi_{2*} \circ (\text{id} \times \overline{\mathbf{U}}) \circ \kappa^*)(\mathcal{F})$ at W , up to homotopy equivalence. Explicitly, the value at W is the complex whose m^{th} term is

$$\bigoplus_{i=0}^{t^2} \bigoplus_{d \geq 0} \bigoplus_{l \in \text{Pic}(X)} \omega_{(l-d, -i-m)} \otimes (\mathcal{C}^{\mathcal{F}(-l)} \otimes \mathcal{C}^{\mathcal{O}(d)|_W})_{-i},$$

with the induced differential. It suffices to check that if we sheafify the presheaf

$$W \mapsto \bigoplus_{i=0}^{t^2} \omega_{(l-d, i-n)} \otimes (\mathcal{C}^{\mathcal{F}(-l)} \otimes \mathcal{C}^{\mathcal{O}(d)|_W})_{-i},$$

we get

$$\bigoplus_{i=0}^t \omega_{(l-d, i-n)} \otimes \mathcal{C}_{-i}^{\mathcal{F}(-l)} \otimes \mathcal{O}(d).$$

And of course we need to check the differentials coincide as well. Need to fill in the rest of the details, but I think this is clear. \square

Proof of Theorem 4.2. Combine Propositions 5.1 and 5.3. \square