

TITLE

CONTENTS

1. Introduction	1
2. Differential modules	1
2.1. Expansion	2
2.2. Projective flag resolutions	2
2.3. \otimes and $\underline{\text{Hom}}$ for differential modules	6
2.4. Minimal free resolutions	7
3. The toric BGG correspondence	12
3.1. The toric BGG functors	13
3.2. Properties of the functors \mathbf{L} and \mathbf{R}	16
4. Pushforward of differential modules	18
5. Tate modules	20
5.1. Exactness properties of $\mathbf{T}(\mathcal{E})$	22
5.2. Algebraic characterization of $\mathbf{T}(\mathcal{E})$	23
6. Applications	25
6.1. Toric Syzygy Theorem	25
7. The Tate resolution and A_∞ -operations on sheaf cohomology	26
7.1. A_∞ operations and the tensor coalgebra	26
7.2. Perturbations and A_∞ -structures	27
7.3. The Tate differential and A_∞ operations	27
References	28

1. INTRODUCTION

Notational conventions:

- We index homologically throughout.

2. DIFFERENTIAL MODULES

Let A be an abelian group, and let R be an A -graded ring (for instance, A could be 0). All modules over R are right modules. ♣♣♣ Michael: [We work with right modules because our main example will be $R = E$, and in Macaulay2, entries of matrices over E act on the right. This is the same reason I'm working with homological indexing as opposed to cohomological: I'm trying to match M2.]

Definition 2.1. Let $a \in A$. A *degree a differential R -module* is a pair (D, ∂_D) , where D is an A -graded module, and

$$\partial : D \rightarrow D(a)$$

is an R -linear map such that $\partial^2 = 0$. When the fixed element a of A is clear, we will just call (D, ∂_D) a *differential module*. A morphism $(D, \partial) \rightarrow (D', \partial')$ of degree a differential modules is a map $f : D \rightarrow D'$ satisfying $f \circ \partial = \partial' \circ f$.

For the rest of section, fix an element $a \in A$. Let $\text{DM}(R, a)$ denote the category of degree a differential R -modules. The *homology* of an object $(D, \partial) \in \text{DM}(R, a)$ is the subquotient

$$\ker(\partial : D \rightarrow D(a)) / \text{im}(\partial : D(-a) \rightarrow D),$$

denoted $H(D, \partial)$. A morphism in $\text{DM}(R, a)$ is a *quasi-isomorphism* if it induces an isomorphism on homology. A *homotopy* of morphisms $f, f' : (D, \partial) \rightarrow (D', \partial')$ in $\text{DM}(R, a)$ is a morphism $h : D \rightarrow D'(-a)$ of A -graded R -modules such that $f - f' = h\partial + \partial'h$. The *mapping cone* of a morphism $f : (D, \partial) \rightarrow (D', \partial')$ in $\text{DM}(R, a)$ is the object $(D \oplus D'(-a), \begin{pmatrix} -\partial & 0 \\ f & \partial' \end{pmatrix})$.

2.1. Expansion. Let $\text{Com}_{\text{per}}(R, a)$ denote the category of complexes of A -graded R -modules satisfying

$$D[j] = D(-ja)$$

for all $j \in \mathbb{Z}$, with morphisms given by maps of complexes that are identical in each homological degree. ♣♣♣ David: [nonsense for $j = 0$, assuming that morphisms are degree 0 homogeneous. What's really meant?] ♣♣♣ Michael: [Fixed.] There is an equivalence of categories

$$\text{Ex} : \text{DM}(R, a) \xrightarrow{\cong} \text{Com}_{\text{per}}(R, a)$$

given by sending the differential module (D, ∂) to the complex

$$\cdots \xrightarrow{\partial} D(-a) \xrightarrow{\partial} D \xrightarrow{\partial} D(a) \xrightarrow{\partial} \cdots$$

Following [ABI07, Section 1.4], we call $\text{Ex}(D, \partial)$ the *expansion* of (D, ∂) . The above notions of homology, quasi-isomorphism, homotopy, and mapping cone for differential modules all correspond to the usual notions via expansion. ♣♣♣ Michael: [I got rid of the signs in the differential of the expansion. I think it's better to follow ABI precisely here.]

2.2. Projective flag resolutions. We are interested in differential modules equipped with a filtration, in the following sense (cf. [ABI07, 2.1]).

Definition 2.2 (cf. [ABI07] Section 2.1). A *flag* is an object $(D, \partial) \in \text{DM}(R, a)$ equipped with a filtration $\mathcal{F}_\bullet D$ such that

- $\mathcal{F}_i D \subseteq \mathcal{F}_{i+1} D$
- $\partial(\mathcal{F}_i D) \subseteq \mathcal{F}_{i-1} D$,
- $\bigcup_i \mathcal{F}_i D = D$, and
- $\mathcal{F}_{<0} D = 0$.

We say a flag is *locally finitely generated* if each component of the associated graded module is finitely generated. A *split flag* is a differential module (D, ∂) equipped with a decomposition $D = \bigoplus_{j \in \mathbb{Z}} D_j$ such that the filtration $\mathcal{F}_i D = \bigoplus_{j < i} D_j$ makes (D, ∂) a flag. A *projective (resp. free) split flag* is a split flag such that each D_j is projective (resp. free).

Remark 2.3. A split flag (D, ∂) such that $\partial(D_i) \subseteq D_{i-1}$ is the same thing as a chain complex of R -modules that is concentrated in nonnegative degrees. ♣♣♣ David: [maybe $\partial(D_i) \subseteq D_{i-1}$] ♣♣♣ Michael: [Fixed.]

Definition 2.4. Let $(D, \partial_D) \in \text{DM}(R, a)$, and let $(P, \partial_P) \in \text{DM}(R, a)$ be a projective (resp. free) split flag. A quasi-isomorphism $\epsilon : (P, \partial_P) \rightarrow (D, \partial_D)$ is called a *projective flag resolution* (resp. *free flag resolution*). A projective (resp. free) flag resolution is called *locally finitely generated* if the flag P is such.

Proposition 2.5. *Every $(D, \partial) \in \text{DM}(R, a)$ admits a free flag resolution.*

Proof. Choose a set of cycles in D that descends to a generating set of $H(D)$, and let F_0 be a free R -module with basis indexed by this set. Let $\epsilon_0 : (F_0, 0) \rightarrow (D, \partial_D)$ be the morphism of differential modules that sends each basis element to its associated cycle. Next, choose a set of cycles in $\text{cone}(\epsilon_0)$ that descends to a generating set of $H(\text{cone}(\epsilon_0))$, let F_1 be a free R -module with basis indexed by this set, and define $\epsilon_1 : (F_1, 0) \rightarrow \text{cone}(\epsilon_0)$ as before. Iterating this process, we obtain a flag $(F = \bigoplus_{i \geq 0} F_i, \partial_F)$ and a quasi-isomorphism $\epsilon : (F, \partial_F) \xrightarrow{\sim} (D, \partial_D)$. \square

♣♣♣ Michael: [I don't think the Cartan-Eilenberg resolution yields a flag, so the killing cycles algorithm seems to be our only method for building free flag resolutions.]

As in classical homological algebra, morphisms of differential modules may be lifted to projective flag resolutions in a unique way, up to homotopy. More generally, we have the following

Proposition 2.6. *Let $(D, \partial_D), (D', \partial_{D'}) \in \text{DM}(R, a)$, and suppose we have morphisms $\epsilon : (P, \partial_P) \rightarrow (D, \partial_D)$, $\epsilon' : (P', \partial_{P'}) \rightarrow (D', \partial_{D'})$, where (P, ∂_P) is a projective split flag, and ϵ' is a quasi-isomorphism. Given a morphism $f : (D, \partial_D) \rightarrow (D', \partial_{D'})$ of differential modules, there exists a morphism*

$$\tilde{f} : \text{cone}(\epsilon) \rightarrow \text{cone}(\epsilon')$$

of differential modules of the form

$$(1) \quad \begin{pmatrix} \alpha & 0 \\ \rho & f \end{pmatrix}.$$

In particular, the entry $\alpha : P \rightarrow P'$ of (1) is a morphism of differential modules. Moreover, given two such lifts

$$\tilde{f}_1 = \begin{pmatrix} \alpha_1 & 0 \\ \rho_1 & f \end{pmatrix}, \tilde{f}_2 = \begin{pmatrix} \alpha_2 & 0 \\ \rho_2 & f \end{pmatrix} : \text{cone}(\epsilon) \rightarrow \text{cone}(\epsilon'),$$

there is a homotopy

$$h = \begin{pmatrix} h_1 & 0 \\ h_2 & 0 \end{pmatrix} : P \oplus D \rightarrow P'(-a) \oplus D'(-a).$$

between \tilde{f}_1 and \tilde{f}_2 . In particular, h_1 is a homotopy between α_2 and α_1 .

Remark 2.7. It need not be the case that $\epsilon'\alpha = f\epsilon$. For instance, ♣♣♣ Michael: [Fill in.]

Proof. Set $\tilde{P} := \text{cone}(\epsilon)$ and $\tilde{P}' := \text{cone}(\epsilon')$. We begin by defining $g_0 : P_0 \rightarrow \tilde{P}'$ such that the map

$$\tilde{f}_0 : P_0 \oplus D \rightarrow \tilde{P}'$$

given by $(p, d) \mapsto g_0(p) + (0, f(d))$ is a morphism of differential modules, where $P_0 \oplus D$ is equipped with the differential $\begin{pmatrix} 0 & 0 \\ \epsilon & \partial_D \end{pmatrix}$, i.e. the restriction of $\partial_{\tilde{P}}$ to $P_0 \oplus D$. We have a diagram

$$\begin{array}{ccc} & & \tilde{P}' \\ & & \downarrow \partial_{\tilde{P}'} \\ P_0 & \xrightarrow{\beta} & \text{im}(\partial_{\tilde{P}'}) = \ker(\partial_{\tilde{P}'}), \end{array}$$

where $\beta(p) = (0, (f\epsilon)(p))$. Note that β does indeed land in $\ker(\partial_{\tilde{P}'})$: we have

$$(\partial_{\tilde{P}'}\beta)(p) = (0, (\partial_{D'}f\epsilon)(p)) = (0, (f\partial_D\epsilon)(p)) = 0;$$

the last equality holds since $\partial_P|_{P_0} = 0$, and $\epsilon\partial_P = \partial_D\epsilon$. Since P_0 is projective, we get an induced map

$$g_0 : P_0 \rightarrow \tilde{P}'$$

making the diagram commute. One easily checks that g_0 has the desired property: if $(p, d) \in P_0 \oplus D$,

$$\begin{aligned} (\tilde{f}_0\partial_{\tilde{P}})(p, d) &= (0, (f\epsilon)(p)) + (0, (f\partial_D)(d)) \\ &= \beta(p) + (0, (\partial_{D'}f)(d)) \\ &= (\partial_{\tilde{P}'}g_0)(p) + \partial_{\tilde{P}'}(0, f(d)) \\ &= (\partial_{\tilde{P}'}\tilde{f}_0)(p, d). \end{aligned}$$

Now, suppose $n > 0$, and assume we have

$$g_i : P_{\leq i} \rightarrow \tilde{P}'$$

for all $i < n$, such that

- the map $\tilde{f}_i : P_{\leq i} \oplus D \rightarrow \tilde{P}'$ given by $(p, d) \mapsto g_i(p) + (0, f(d))$ is a morphism of differential modules (where $P_{\leq i} \oplus D$ is equipped with the differential given by the restriction of $\partial_{\tilde{P}}$), and
- $g_i|_{P_{\leq j}} = g_j$ for all $j < i$.

We have a diagram

$$\begin{array}{ccc} & & \tilde{P}' \\ & & \downarrow \partial_{\tilde{P}'} \\ P_n & \xrightarrow{\gamma} & \text{im}(\partial_{\tilde{P}'}) = \ker(\partial_{\tilde{P}'}), \end{array}$$

where $\gamma(p) = (\tilde{f}_{n-1}\partial_{\tilde{P}})(p, 0)$; the map γ lands in $\ker(\partial_{\tilde{P}'})$, since

$$(\partial_{\tilde{P}'}\tilde{f}_{n-1}\partial_{\tilde{P}})(p, 0) = (\tilde{f}_{n-1}\partial_{\tilde{P}}\partial_{\tilde{P}})(p, 0) = 0.$$

Since P_n is projective, we obtain a map $\tilde{\gamma} : P_n \rightarrow \tilde{P}'$ making the diagram commute. We define $g_n : P_{\leq n} \rightarrow \tilde{P}'$ to be the map

$$(g_{n-1} \quad \tilde{\gamma}) : P_{\leq n-1} \oplus P_n \rightarrow \tilde{P}'.$$

We now verify that the map

$$\tilde{f}_n : P_{\leq n} \oplus D \rightarrow \tilde{P}',$$

given by $(p, d) \mapsto g_n(p) + (0, f(d))$, is a morphism of differential modules. Let $(p, d) \in P_{\leq n} \oplus D$. We have:

$$\begin{aligned} (\tilde{f}_n \partial_{\tilde{P}})(p, d) &= g_n(-\partial_P(p)) + (0, (f\epsilon)(p) + (f\partial_D)(d)) \\ &= \tilde{f}_n(-\partial_P(p), \epsilon(p)) + (0, (\partial_{D'}f)(d)) \\ &= (\tilde{f}_n \partial_{\tilde{P}})(p, 0) + (\partial_{\tilde{P}'} \tilde{f}_n)(0, d), \end{aligned}$$

so it suffices to show

$$(\tilde{f}_n \partial_{\tilde{P}})(p, 0) = (\partial_{\tilde{P}'} \tilde{f}_n)(p, 0).$$

To see this, write $p = p' + p''$, where $p' \in P_{\leq n-1}$ and $p'' \in P_n$, and notice that

$$\begin{aligned} (\tilde{f}_n \partial_{\tilde{P}})(p, 0) &= (\tilde{f}_{n-1} \partial_{\tilde{P}})(p, 0) \\ &= (\tilde{f}_{n-1} \partial_{\tilde{P}})(p', 0) + (\tilde{f}_{n-1} \partial_{\tilde{P}})(p'', 0) \\ &= (\partial_{\tilde{P}'} \tilde{f}_{n-1})(p', 0) + \gamma(p'') \\ &= (\partial_{\tilde{P}'} \tilde{f}_n)(p', 0) + (\partial_{\tilde{P}'} g_n)(p'') \\ &= (\partial_{\tilde{P}'} \tilde{f}_n)(p', 0) + (\partial_{\tilde{P}'} \tilde{f}_n)(p'', 0) \\ &= (\partial_{\tilde{P}'} \tilde{f}_n)(p, 0). \end{aligned}$$

Let g be the colimit of the g_i , and take $\tilde{f} : \tilde{P} \rightarrow \tilde{P}'$ to be given by $(p, d) \mapsto g(p) + (0, f(d))$. We now show our lift \tilde{f} is unique up to homotopy. Without loss, assume $f = 0$; we will show \tilde{f} is null homotopic. We again proceed by induction. We have a diagram

$$\begin{array}{ccc} & \tilde{P}' & \\ & \downarrow \partial_{\tilde{P}'} & \\ P_0 & \xrightarrow{g_0} & \ker(\partial_{\tilde{P}'}), \end{array}$$

since $(\partial_{\tilde{P}'} g_0)(p) = \beta(p) = 0$ for all $p \in P_0$. Since P_0 is projective, we obtain a map $s_0 : P_0 \rightarrow \tilde{P}'$ making the diagram commute. Let $n > 0$, and suppose we have maps $s_i : P_{\leq i} \rightarrow \tilde{P}'$ for $i < n$ such that

- $g_i = \partial_{\tilde{P}'} s_i - s_{i-1} \partial_P$ (set $s_{<0} := 0$), and
- $s_i|_{P_{\leq j}} = s_j$ for all $j < i$.

In particular, let's record the relation

$$(2) \quad g_{n-1} = \partial_{\tilde{P}'} s_{n-1} - s_{n-2} \partial_P.$$

We have a diagram

$$\begin{array}{ccc} & \tilde{P}' & \\ & \downarrow \partial_{\tilde{P}'} & \\ P_{\leq n} & \xrightarrow{g_n + s_{n-1} \partial_P} & \ker(\partial_{\tilde{P}'}), \end{array}$$

since, by (2), we have

$$\begin{aligned}\partial_{\tilde{P}'}(g_n + s_{n-1}\partial_P) &= \partial_{\tilde{P}'}g_n + (g_{n-1} + s_{n-2}\partial_P)\partial_P \\ &= \partial_{\tilde{P}'}g_n + g_{n-1}\partial_P,\end{aligned}$$

and

$$\begin{aligned}(\partial_{\tilde{P}'}g_n)(p) &= (\partial_{\tilde{P}'}\tilde{f}_n)(p, 0) \\ &= (\tilde{f}_n\partial_{\tilde{P}})(p, 0) \\ &= \tilde{f}_n(-\partial_P(p), \epsilon(p)) \\ &= -(g_{n-1}\partial_P)(p).\end{aligned}$$

Define $s_n : P_{\leq n} \rightarrow \tilde{P}'$ making the diagram commute. Let s denote the colimit of the s_i . We have

$$g = \partial_{\tilde{P}'}s - s\partial_P.$$

Now take $h : \tilde{P} \rightarrow \tilde{P}'$ to be the map given by $(p, d) \mapsto s(p)$, and observe that

$$\begin{aligned}\tilde{f}(p, d) &= g(p) \\ &= (\partial_{\tilde{P}'}s)(p) - (s\partial_P)(p) \\ &= (\partial_{\tilde{P}'}h)(p, d) + (h\partial_{\tilde{P}})(p, d).\end{aligned}$$

□

2.3. \otimes and $\underline{\text{Hom}}$ for differential modules. As stated in [ABI07], there is no tensor product for general differential modules. However, suppose A is of the form $B \times \mathbb{Z}$ for some abelian group B , and let $a \in A$ be of the form $(b, \pm 1)$ for some $b \in B$. We can use the \mathbb{Z} -grading to define notions of tensor product and internal Hom for degree a differential R -modules in the following way. First, if N is an A -graded R -module, and $n \in N$ is homogeneous with respect to the \mathbb{Z} -grading, define $\|n\| \in \mathbb{Z}$ to be the \mathbb{Z} -degree of n . Let $(D, \partial_D), (D', \partial_{D'}) \in \text{DM}(R, a)$, and assume D and D' are equipped with left R -actions making them R - R -bimodules. Define

$$D \otimes_R^{\text{DM}} D' = (D \otimes_R D', d \otimes d' \mapsto \partial_D(d) \otimes d' + (-1)^{\|d\|} d \otimes \partial_{D'}(d')) \in \text{DM}(R, a)$$

and

$$\underline{\text{Hom}}_R^{\text{DM}}(D, D') = (\underline{\text{Hom}}_R(D, D'), f \mapsto f \circ \partial_D - (-1)^{\|f\|} \partial_{D'} \circ f) \in \text{DM}(R, a).$$

Let $(D, \partial_D) \in \text{DM}(R, a)$. It's clear that there is an adjunction

$$-\otimes_R^{\text{DM}} D : \text{DM}(R, a) \rightleftarrows \text{DM}(R, a) : \underline{\text{Hom}}_R^{\text{DM}}(D, -).$$

Remark 2.8. The ring R , thought of as a B -graded ring, may be considered as a dg-algebra (over \mathbb{Z}) with trivial differential and homological grading induced by the \mathbb{Z} -grading. When $a = (0, -1)$, the category $\text{DM}(R, a)$ is equivalent (in fact, isomorphic) to the category of dg-modules over this dg-algebra, and the notions of tensor product and internal Hom described above correspond to the usual ones for dg-modules.

Let F be a free flag resolution of D , and let $(D', \partial_{D'})$ be another object in $\mathrm{DM}(R, a)$. We define

$$\mathrm{Tor}_{\mathrm{DM}}^R(D, D') = H(F \otimes_R^{\mathrm{DM}} D')$$

and

$$\mathrm{Ext}_R^{\mathrm{DM}}(D, D') = H(\mathrm{Hom}_R^{\mathrm{DM}}(F, D')).$$

It follows easily from Proposition 2.6 that these definitions do not depend on the choice of free flag resolution.

Remark 2.9. Suppose A is an arbitrary grading group, $a \in A$, and $(D, \partial_D), (D', \partial_{D'}) \in \mathrm{DM}(R, a)$, where D, D' are R - R -bimodules. When $\partial_{D'} = 0$, the definitions of $D \otimes_R^{\mathrm{DM}} D'$, $\mathrm{Hom}_R^{\mathrm{DM}}(D, D')$, $\mathrm{Tor}_{\mathrm{DM}}^R(D, D')$, and $\mathrm{Ext}_R^{\mathrm{DM}}(D, D')$ still make sense.

2.4. Minimal free resolutions. From now on, assume either that

- (1) the grading group A is trivial and R is local, or
- (2) the set $\bigoplus_{a \neq 0} R_a$ is a maximal ideal of R .

♣♣♣ **Michael:** [Are these the right assumptions? Double check. We just need graded Nakayama.] ♣♣♣

Daniel: [Maybe you just need that R_0 is a local ring?] Denote the unique (homogeneous, in the second case) maximal ideal by \mathfrak{m} . We say a morphism $f : M \rightarrow N$ of A -graded R -modules is *minimal* if $f(M) \subseteq \mathfrak{m}N$.

We now wish to define a notion of minimal free resolutions for differential R -modules. It is tempting to define such a resolution to be a minimal free resolution. But, we will see in Example 2.17 that such resolutions do not exist in general. Instead, we proceed as follows.

Definition 2.10. A *trivial* differential R -module is a direct sum of objects in $\mathrm{DM}(R, a)$ of the form

$$R(b) \oplus R(b-a) \xrightarrow{\begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}} R(b+a) \oplus R(b)$$

for some $b \in A$.

Remark 2.11. A free differential module (F, ∂_F) is isomorphic to a trivial differential module if and only if it is *contractible*, i.e. the identity map on F is null-homotopic.

♣♣♣ **Michael:** [Double check this remark.]

Proposition 2.12. Let (F, ∂_F) be either a finitely generated free differential module or a locally finitely generated free split flag. There is an automorphism A of F such that

$$(F, A\partial_F A^{-1}) = (T, \partial_T) \oplus (M, \partial_M),$$

where (T, ∂_T) is trivial and (M, ∂_M) is minimal.

Proof. Suppose first that F is finitely generated. Choose a basis of F , and view ∂_F as a matrix with respect to this basis. If ∂_F has no unit entries, then it is minimal and we are done. Otherwise, the condition $\partial_F^2 = 0$ forces ∂_F to have a unit entry u that does not lie on the diagonal. Without loss of generality, we can assume that this entry is in the first column and second row. Let B_1 be the matrix corresponding to the row operations that zero out

all other entries in the first column of ∂_F . This is an identity matrix, except in the second column. It follows that $B_1\partial_FB_1^{-1}$ has the form:

$$B_1\partial_FB_1^{-1} = \begin{pmatrix} 0 & a'_{1,2} & a'_{1,3} & \cdots \\ u & a_{2,2} & a_{2,3} & \cdots \\ 0 & a'_{3,2} & a'_{3,3} & \cdots \\ 0 & a'_{4,2} & a'_{4,3} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

Let B_2 the matrix corresponding the column operations which zero out all the other entries in the second row of ∂_F . This is an identity matrix, except in the top row. It follows that $B_2^{-1}B_1\partial_FB_1^{-1}B_2$ has the form

$$B_2^{-1}B_1\partial_FB_1^{-1}B_2 = \begin{pmatrix} 0 & a''_{1,2} & a''_{1,3} & \cdots \\ u & 0 & 0 & \cdots \\ 0 & a''_{3,2} & a''_{3,3} & \cdots \\ 0 & a''_{4,2} & a''_{4,3} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix}.$$

The first column of ∂_F^2 equals the second column of ∂_F multiplied by u . Since $\partial_F^2 = 0$, this means that the entire second column is zero. Similarly, the second row of ∂_F^2 is the first row of ∂_F multiplied by u , and thus the entire first row of ∂_F must be zero. We conclude that

$$(F, B_2^{-1}B_1\partial_FB_1^{-1}B_2) = (T, \partial_T) \oplus (D, \partial_D),$$

where T is a rank 2 free R -module, and $\partial_T = \begin{pmatrix} 0 & 0 \\ u & 0 \end{pmatrix}$. Without loss, we can assume $u = 1$.

Now apply the same argument to (D, ∂_D) . Since F is finitely generated, this process eventually terminates.

Suppose now that F is a locally finitely generated free split flag. We can apply the above argument to each summand F_i , yielding automorphisms A_i such that $(F_i, A_i\partial_FA_i^{-1}) = (M_i, \partial_{M_i}) \oplus (T_i, \partial_{T_i})$ for all i . Moreover, the above argument shows that we can choose the trivial summands to be compatible for increasing i : that is, we may assume that there are inclusions $T_i \rightarrow T_{i+1}$ for all i , such that

$$\begin{array}{ccc} T_i & \longrightarrow & F_i \\ \downarrow & & \downarrow \\ T_{i+1} & \longrightarrow & F_{i+1} \end{array}$$

commutes for all i . These diagrams induce maps $M_i \rightarrow M_{i+1}$, which may not be inclusions. We let (M, ∂_M) be the colimit of the (M_i, ∂_{M_i}) , and similarly for (T, ∂_T) . Since colimits commute with coproducts, $F = T \oplus M$. It is clear from the construction that T is trivial. Since each ∂_{M_i} factors through $\mathfrak{m}M_i$, $\text{colim } \partial_{M_i}$ factors through $\mathfrak{m} \text{colim } M_i = \mathfrak{m}M$. \square

Definition 2.13. Let $(D, \partial_D) \in \text{DM}(R, a)$. A *stably free flag resolution* of (D, ∂_D) is a free differential module (G, ∂_G) such that there is a free flag resolution

$$(F, \partial_F) \xrightarrow{\epsilon} (D, \partial_D)$$

8

and a trivial differential module (T, ∂_T) satisfying $(F, \partial_F) = (G, \partial_G) \oplus (T, \partial_T)$. We say (G, ∂_G) is a *minimal stably free flag resolution* if ∂_G is minimal. We will shorten “minimal stably free flag resolution” to “minimal free resolution” from now on. ♣♣♣ Daniel: [Should we call this a “flag retract” or something?] ♣♣♣ Michael: [I think not, because retract just means summand of a flag, but we want a summand of a flag *whose complement is trivial*. So I think “stable” really is the right term here.] ♣♣♣ Daniel: [At least let’s add a remark comparing this with retract.]

Proposition 2.14. *Every $(D, \partial_D) \in \text{DM}(R, a)$ has a minimal free resolution.*

Proof. Combine Propositions 2.5 and 2.12. □

To prove uniqueness of minimal free resolutions, we will need the following

Lemma 2.15. *Let $(M, \partial_M), (M', \partial_{M'})$ be minimal differential modules. If a morphism*

$$f : (M, \partial_M) \rightarrow (M', \partial_{M'})$$

factors through a trivial differential module, f is minimal.

Proof. Suppose we have a factorization

$$(M, \partial_M) \xrightarrow{g} (T, \partial_T) \xrightarrow{h} (M', \partial_{M'})$$

of f , where (T, ∂_T) is trivial. Let $m \in M$, and choose a basis $\{e_i\}_{i \in I}$ of T . We can write $g(m)$ as

$$r_1 e_{i_1} + \cdots + r_n e_{i_n}.$$

Suppose $r_j \notin \mathfrak{m}$. Since $r_1 \partial_T(e_{i_1}) + \cdots + r_n \partial_T(e_{i_n}) = \partial_T(g(m)) = g(\partial_M(m)) \in \mathfrak{m}T$, and ∂_T is a matrix with at most a single 1 in each row and 0’s elsewhere, we have $\partial_T(e_{i_j}) = 0$. Using that T is exact, choose an element $t \in T$ such that $e_{i_j} = \partial_T(t)$. Then

$$h(e_{i_j}) = h(\partial_T(t)) = \partial_{M'}(h(t)) \in \mathfrak{m}M'.$$

We conclude that $f(m) \in \mathfrak{m}M'$. □

Theorem 2.16. *Let $(D, \partial_D) \in \text{DM}(R, a)$, and let*

$$\begin{array}{ccc} (M, \partial_M) \oplus (T, \partial_T) & & (M', \partial_{M'}) \oplus (T', \partial_{T'}) \\ & \searrow \epsilon & \swarrow \epsilon' \\ & (D, \partial_D) & \end{array}$$

be locally finite free flag resolutions, where $(M, \partial_M), (M', \partial_{M'})$ are minimal and $(T, \partial_T), (T', \partial_{T'})$ are trivial. There is an isomorphism $(M, \partial_M) \cong (M', \partial_{M'})$. In particular, minimal free resolutions of differential modules are unique up to isomorphism.

Proof. Applying Proposition 2.6 to the identity map on D , we may choose morphisms

$$\begin{aligned} \alpha &= \begin{pmatrix} \alpha_1 & \alpha_2 \\ \alpha_3 & \alpha_4 \end{pmatrix} : (M, \partial_M) \oplus (T, \partial_T) \rightarrow (M', \partial_{M'}) \oplus (T', \partial_{T'}) \\ \alpha' &= \begin{pmatrix} \alpha'_1 & \alpha'_2 \\ \alpha'_3 & \alpha'_4 \end{pmatrix} : (M', \partial_{M'}) \oplus (T', \partial_{T'}) \rightarrow (M, \partial_M) \oplus (T, \partial_T) \end{aligned}$$

of differential modules and homotopies

$$h = \begin{pmatrix} s_1 & s_2 \\ s_3 & s_4 \end{pmatrix} : M \oplus T \rightarrow M(0, -1) \oplus T(0, -1)$$

$$h' = \begin{pmatrix} s'_1 & s'_2 \\ s'_3 & s'_4 \end{pmatrix} : M' \oplus T' \rightarrow M'(0, -1) \oplus T'(0, -1)$$

such that

$$\alpha' \alpha - \text{id}_F = h \partial_{M \oplus T} + \partial_{M \oplus T} h$$


$$\alpha \alpha' - \text{id}_{F'} = h' \partial_{M' \oplus T'} + \partial_{M' \oplus T'} h'$$

Reading off the top-left entry of the matrices on each side of these equations, we get

$$\alpha'_1 \alpha_1 + \alpha'_2 \alpha_3 - \text{id}_F = h_1 \partial_M + \partial_M h_1$$

$$\alpha_1 \alpha'_1 + \alpha_2 \alpha'_3 - \text{id}_{F'} = h'_1 \partial_{M'} + \partial_{M'} h'_1.$$

By Lemma 2.15, $\alpha'_2 \alpha_3$ and $\alpha_2 \alpha'_3$ are minimal. We conclude that $\alpha'_1 \alpha_1 = \text{id}_M$ and $\alpha_1 \alpha'_1 = \text{id}_{M'}$ modulo \mathfrak{m} .

Now, assume R is nontrivially graded. By the graded version of Nakayama's Lemma and the local finiteness of M and M' , it follows that $\alpha'_1 \alpha_1$ and $\alpha_1 \alpha'_1$ are automorphisms. In particular, α_1 is injective and surjective. In the case where R is trivially graded... 
Michael: [Not sure how this is going to work in non-graded case. May just need to assume M and M' are finitely generated in this case.] \square

Example 2.17. We now give an example of a differential module with no minimal free flag resolution. Take $A = \mathbb{Z}$, $a = 2$, and $R = k[x, y]$, where $|x| = 1 = |y|$. Let $D = R^{\oplus 2}$, and take

$$\partial_D : R^{\oplus 2} \rightarrow R(2)^{\oplus 2}$$

to be

$$\begin{pmatrix} xy & -x^2 \\ y^2 & -xy \end{pmatrix}.$$

Since (D, ∂_D) does not admit a flag structure ([ABI07]), it suffices, by Theorem 2.16, to show that (D, ∂_D) is the minimal free resolution of itself.

We use the Cartan-Eilenberg construction to produce a free flag resolution of (D, ∂_D) . The cycles are the rank 1 free submodule of R^2 generated by $\begin{pmatrix} x \\ y \end{pmatrix}$, so Z is resolved by $G := [R(-1)]$. The boundaries B are the image of the above matrix, and so $B(-2)$ is resolved by $H := [R(-2)^2 \leftarrow R(-3)]$. Using this, we can produce a Cartan-Eilenberg resolution of (D, ∂_D) given by

$$F = G_0 \oplus H_0 \oplus H_1(2) = R(-1) \oplus R(-2) \oplus R(-2) \oplus R(-1),$$

$$\partial_F = \begin{pmatrix} 0 & -y & -x & 1 \\ 0 & 0 & 0 & -x \\ 0 & 0 & 0 & y \\ 0 & 0 & 0 & 0 \end{pmatrix},$$

and

$$\epsilon = \begin{pmatrix} x & -1 & 0 & 0 \\ y & 0 & 1 & 0 \end{pmatrix} : F \rightarrow D.$$

Now, take $A = \begin{pmatrix} -1 & 0 & 0 & 0 \\ x & -1 & 0 & 0 \\ y & 0 & 1 & 0 \\ 0 & -y & -x & -1 \end{pmatrix}$, so that

$$A\partial_F A^{-1} = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & xy & -x^2 & 0 \\ 0 & y^2 & -xy & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}.$$

Observe that $(F, A\partial_F A^{-1}) \cong (F, \partial_F)$ is a direct sum of (D, ∂_D) and a trivial object.

Example 2.18. In the category of complexes over a graded or local ring, any complex with bounded homology will admit a minimal free resolution. If you start with a complex which is free, minimal, and bounded, then it will equal its own minimal free resolution. However, if the original complex was free and minimal, but not bounded above, then it will generally not equal its minimal free resolution. For instance, if $R = k[x]/(x^2)$ and one considers the complex F :

$$0 \rightarrow R \xrightarrow{x} R(1) \xrightarrow{x} R(2) \xrightarrow{x} \cdots,$$

then the minimal free resolution will be the complex F' :

$$\cdots \xrightarrow{x} R(-3) \xrightarrow{x} R(-2) \xrightarrow{x} R(-1) \rightarrow 0.$$

And of course F' is not isomorphic to F .

A similar phenomenon occurs for differential modules: namely one can find a free, minimal differential module which is not isomorphic to its minimal free resolution. Essentially the same example works. Consider the differential module (D, ∂_D) obtained from the complex F by forgetting the homological grading. Let (D', ∂'_D) be the differential module obtained from F' in a similar way. The homological grading on F' can realize F' as a flag, and the quasi-isomorphism of complexes $F' \xrightarrow{x} F$ shows that (D', ∂'_D) is a flag, free resolution of D . Of course, D' is not isomorphic to D because the underlying graded modules are distinct. By the uniqueness of minimal free resolutions, it follows that D is not a minimal free resolution of itself.

Remark 2.19. Results similar to these are proven Avramov-Foxby-Halperin's unpublished notes, but with boundedness assumptions that are not satisfied in our setting. ♣♣♣ Michael: [Fill this in.]

Example 2.20. When R is a graded algebra over a field hypotheses as above, $\text{Tor}(M, k)$ computes

Lemma 2.21. *Let R be a non-negatively \mathbb{Z} -graded ring. and let (F, ∂) be a free graded differential R -module. Assume $\partial F \subseteq R_+ F$. For any integer d , (F, ∂) may be realized as an extension of free differential modules*

$$(F_{<d}, \partial|_{F_{<d}}) \xrightarrow{\epsilon} (F_{\geq d}, \partial|_{F_{\geq d}}).$$

The differential module $(F_{<d}, \partial|_{F_{<d}})$ has a natural split flag structure where \mathcal{F}^i consists of all summands of the form $R(-i)$. In particular, if F has at most finitely many generators of degree i for each i , then $(F_{<d}, \partial|_{F_{<d}})$ is the minimal free resolution of $(F_{\geq d}, \partial|_{F_{\geq d}})$.

A similar statement holds for non-positively \mathbb{Z} -graded rings.

Proof. The differential ∂ has a block decomposition of the form:

$$\begin{pmatrix} \partial|_{F < d} & \epsilon \\ 0 & \partial|_{F \geq d} \end{pmatrix}$$

The rest of the statement is straightforward, though we need a hypothesis to guarantee uniqueness of minimal free resolutions. \square

3. THE TORIC BGG CORRESPONDENCE

Let k be a field and V a k -vector space with basis $\{e_0, \dots, e_n\}$. Denote by x_0, \dots, x_n the corresponding basis elements of $W = \text{Hom}_k(V, k)$. Let $E = \bigwedge(V)$ and $S = \text{Sym}(W)$. Equip S (resp. E) with a \mathbb{Z} -grading such that $|x_i| = 1$ (resp. $|e_i| = -1$) for all i . The following theorem is called the *Bernstein-Gel'fand-Gel'fand correspondence*:

Theorem 3.1 ([BGG78]). *Let $\text{Com}(S)$ (resp. $\text{Com}(E)$) denote the category of complexes of graded S -modules (resp. E -modules). There is an adjunction*

$$\mathbf{L}_{\text{st}} : \text{Com}(E) \rightleftarrows \text{Com}(S) : \mathbf{R}_{\text{st}}$$

that induces an equivalence

$$\text{D}^b(E) \simeq \text{D}^b(S).$$

The subscript “st” stands for “standard” and is intended to distinguish these classical BGG functors from the analogous functors introduced below in the toric setting.

Remark 3.2. All E -modules are right modules. However, a right E -module M can be considered as a left E -module with action $em = (-1)^{|e||m|}me$, and vice versa.

We recall the definitions of the functors \mathbf{L}_{st} and \mathbf{R}_{st} . If N is a graded E -module, thought of as an object in $\text{Com}(E)$ concentrated in degree 0, $\mathbf{L}_{\text{st}}(N)$ is the complex with

$$\mathbf{L}_{\text{st}}(N)_j = N_m \otimes_k S(-j)$$

and differential given by multiplication on the right by $\sum_{i=0}^n e_i \otimes x_i$. The functor \mathbf{L}_{st} is extended to complexes by applying the above formula to each term and taking the direct sum totalization of the resulting bicomplex. If M is a graded S -module, the complex $\mathbf{R}_{\text{st}}(M)$ has terms

$$\mathbf{R}_{\text{st}}(M)_j = M_{-j} \otimes_k \omega(j),$$

where $\omega = \text{Hom}_k(E, k)$. Note that ω is (non-canonically) isomorphic to $E(-n-1)$. The differential on $\mathbf{R}_{\text{st}}(M)$ is multiplication on the right by $\sum_{i=0}^n x_i \otimes e_i$. One extends \mathbf{R}_{st} to complexes in the same way as \mathbf{L}_{st} .

Now, let A be an abelian group, let $a_0, \dots, a_n \in A$, and equip $S = \text{Sym}(W)$ with the A -grading given by $|x_i| = a_i$. We wish to formulate a “toric” BGG correspondence involving the category $\text{Com}(S)$ of complexes of A -graded S -modules. This requires a bit of care, as the following example illustrates:

Example 3.3. Suppose $S = k[x_0, x_1]$ is equipped with the \mathbb{Z} -grading such that $|x_0| = 1$ and $|x_1| = 2$. Take $E = \Lambda(e_0, e_1)$, \mathbb{Z} -graded such that $|e_0| = -1$ and $|e_1| = -2$. Let M be a graded S -module, and take $\mathbf{R}(M) = \bigoplus_{j \in \mathbb{Z}} M_{-j} \otimes_k \omega(j)$. Notice that the square-zero endomorphism $\partial_{\mathbf{R}} = x_0 \otimes e_0 + x_1 \otimes e_1$ of $\mathbf{R}(M)$ does not respect the homological grading $\mathbf{R}(M)_j = M_{-j} \otimes_k \omega(j)$. One has the same problem defining the functor \mathbf{L} .

Our solution to the problem in Example 3.3 is as follows:

- (1) We equip the exterior algebra $E = \Lambda(V)$ with an $A \times \mathbb{Z}$ -grading given by $|e_i| = (-a_i, -1)$, and we only consider $A \times \mathbb{Z}$ -graded E -modules. We call the additional \mathbb{Z} -grading the *auxiliary grading* of E . This extra grading allows us to define a homological grading on the output of the functor \mathbf{L} . ♣♣♣ Daniel: [I'm tempted to use a notation like $(a; 1)$ instead of $(a, 1)$, though I could be talked out of this.] ♣♣♣ Michael: [I don't have a strong opinion on this.]
- (2) We want to consider all A -graded S -modules, and not only those equipped with an additional auxiliary grading. We therefore do not impose a homological grading on the image of the functor \mathbf{R} ; instead, we allow \mathbf{R} to take values in the category $\mathrm{DM}(E) := \mathrm{DM}(E, (0, -1))$ of degree $(0, -1)$ differential E -modules.

We now discuss the toric BGG correspondence in detail.

♣♣♣ Daniel: [I think we want to include the functor \mathbf{R}_I where $I \subseteq \{1, \dots, n\}$ and the differential is $\sum_{i \in I} x_i \otimes e_i$. These restricted differentials play a key role in the Tate resolution stuff.] ♣♣♣ Michael: [I agree. Not sure we should expect these functors to be adjoints, by the way. The map \mathbf{R}_I is given by extending scalars to the smaller set of variables, applying \mathbf{R} , and then restricting scalars back to the larger set of variables; the problem is that extension of scalars is a left adjoint, while \mathbf{R} /restriction of scalars are right adjoints.]

3.1. The toric BGG functors. We start by defining functors

$$\mathbf{L} : \mathrm{DM}(E) \rightleftarrows \mathrm{Com}(S) : \mathbf{R}.$$

If $(N, \partial_N) \in \mathrm{DM}(E)$, $\mathbf{L}(N, \partial_N)$ is the complex with terms

$$\mathbf{L}(N, \partial_N)_j = \bigoplus_{a \in A} N_{(a, j)} \otimes_k S(-a)$$

and differential $\partial_{\mathbf{L}} = \sum_{i=0}^n e_i \otimes x_i + (-1)^j \partial_N$.

If M is an S -module concentrated in degree 0, the differential module $\mathbf{R}(M)$ has underlying module

$$\bigoplus_{d \in A} M_{-d} \otimes \omega(d, 0),$$

where, as in the classical BGG correspondence, ω denotes the E -module $\mathrm{Hom}_k(E, k)$. Here, ω is (non-canonically) isomorphic to $E(-\sum_{i=0}^n a_i, -n-1)$. The differential on $\mathbf{R}(M)$ is given by $\partial_{\mathbf{R}} = \sum_{i=0}^n x_i \otimes e_i$. Given an object $C \in \mathrm{Com}(S)$, we define $\mathbf{R}(C)$ as follows: form a bicomplex with q^{th} row given by the expansion $\mathrm{Ex}(\mathbf{R}(C_q))$ and p^{th} vertical differential given by $(-1)^p \partial_C$, apply $\mathrm{Tot}^{\oplus}(-)$ to get an object in $\mathrm{Com}_{\mathrm{per}}(E, (0, -1))$, and apply the equivalence in Section 2.1 to obtain an object in $\mathrm{DM}(E)$. Explicitly: $\mathbf{R}(C)$ has underlying module $\bigoplus_{j \in \mathbb{Z}} \mathbf{R}(C_j)(0, -j)$ and differential $\partial_{\mathbf{R}}$ that acts by $\sum_{i=0}^n x_i \otimes e_i + (-1)^j \partial_C$ on the summand $\mathbf{R}(C_j)(0, -j)$.

Remark 3.4. Our toric BGG functors are related to the classical ones in the following way. Suppose $A = \mathbb{Z}$ and $a_i = 1$ for all i . Recall that $\mathrm{Com}(E)$ denotes the category of complexes of \mathbb{Z} -graded E -modules, where E is equipped with the \mathbb{Z} -grading given by $|e_i| = -1$; while $\mathrm{DM}(E)$ is the category of degree $(0, -1)$ differential E -modules, where E is $\mathbb{Z} \times \mathbb{Z}$ -graded such that $|e_i| = (-1, -1)$. In this case, there is an equivalence (in fact, an isomorphism) of categories

$$\mathrm{Com}(E) \simeq \mathrm{DM}(E)$$

given as follows. Noting that any \mathbb{Z} -graded E -module N may be considered as a $\mathbb{Z} \times \mathbb{Z}$ -graded E -module with components $N_{(i,i)} = N_i$ and $N_{(i,j)} = 0$ for $i \neq j$, we define a functor

$$\text{Fold} : \text{Com}(E) \rightarrow \text{DM}(E)$$

given by $(\cdots \xrightarrow{\partial_C} C_j \xrightarrow{\partial_C} C_{j-1} \xrightarrow{\partial_C} \cdots) \mapsto (\bigoplus_{j \in \mathbb{Z}} C_j(0, -j), \partial_C)$. If $(N, \partial_N) \in \text{DM}(E)$, we set

$$N_j = \{n \in N : |n| = (a, i), \text{ where } i - a = j\}.$$

Notice that N_j is a submodule of N . Since ∂_N is a map from N to $N(0, -1)$, ∂_N induces a map from N_j to N_{j-1} for all j . Noting that any $\mathbb{Z} \times \mathbb{Z}$ -graded E -module M can be considered as a \mathbb{Z} -graded E -module with components $M_a = \bigoplus_{i \in \mathbb{Z}} M_{(a,i)}$, we define a functor

$$\text{Unfold} : \text{DM}(E) \rightarrow \text{Com}(E)$$

by $(N, \partial_N) \mapsto (\cdots \xrightarrow{\partial_N} N_j \xrightarrow{\partial_N} N_{j-1} \xrightarrow{\partial_N} \cdots)$. It's easy to check that Fold and Unfold are inverses. Moreover, we have $\mathbf{L}_{\text{st}} = \mathbf{L} \circ \text{Fold}$ and $\mathbf{R}_{\text{st}} = \text{Unfold} \circ \mathbf{R}$.

Proposition 3.5. *The functors*

$$\mathbf{L} : \text{DM}(E) \rightleftarrows \text{Com}(S) : \mathbf{R}$$

form an adjunction.

Proof. To start, let M be an S -module, and let (N, ∂_N) be a differential E -module. We have

$$\begin{aligned} \text{Hom}_S(\mathbf{L}(N)_j, M) &= \text{Hom}_S\left(\bigoplus_{a \in A} N_{(a,j)} \otimes_k S(-a), M\right) \\ &= \prod_{a \in A} \text{Hom}_k(N_{(a,j)}, M_a) \end{aligned}$$

Now, let C be a complex of S -modules. By the above reasoning, $\text{Hom}_{\text{Com}(S)}(\mathbf{L}(N, \partial_N), C)$ is the subspace of

$$\prod_{a \in A, j \in \mathbb{Z}} \text{Hom}_k(N_{(a,j)}, (M_j)_a)$$

given by morphisms that commute with the differentials. On the other hand, the space $\text{Hom}_{\text{DM}(E)}((N, \partial_N), \mathbf{R}(C))$ is the subspace of

$$\begin{aligned} \text{Hom}_E\left(N, \bigoplus_{a \in A, j \in \mathbb{Z}} (M_j)_{-a} \otimes_k \omega(a, -j)\right) &= \prod_{a \in A, j \in \mathbb{Z}} \text{Hom}_E(N, \text{Hom}_k(E(-a, j), (M_j)_{-a})) \\ &= \prod_{a \in A, j \in \mathbb{Z}} \text{Hom}_k(N(-a, j), (M_j)_{-a}) \\ &= \prod_{a \in A, j \in \mathbb{Z}} \text{Hom}_k(N_{(-a,j)}, (M_j)_{-a}) \end{aligned}$$

given by morphisms that commute with the differentials; the first equality holds because each $(M_j)_{-a} \otimes_k \omega(a, -j)$ is 0 in all but finitely many degrees. Reindexing by replacing a with $-a$, we get

$$\prod_{a \in A, j \in \mathbb{Z}} \text{Hom}_k(N_{(a,j)}, (M_j)_a),$$

as desired. Finally, one checks that the requirements imposed by compatibility with the differentials coincide. \square

♣♣♣ **Michael:** [under construction from here.] The functors \mathbf{L} and \mathbf{R} are not inverses, but they are inverses up to quasi-isomorphism:

Proposition 3.6. *For any $C \in \text{Com}(S)$ and $N \in \text{DM}(E)$, we have equalities*

$$\begin{aligned} (\mathbf{L} \circ \mathbf{R})(C) &= C \otimes_k (\mathbf{L} \circ \mathbf{R})(k) \\ (\mathbf{R} \circ \mathbf{L})(N) &= N \otimes_k^{\text{DM}} (\mathbf{R} \circ \mathbf{L})(k). \end{aligned}$$

In particular, \mathbf{L} and \mathbf{R} are inverses up to natural quasi-isomorphism.

Proof. We have:

$$\begin{aligned} ((\mathbf{L} \circ \mathbf{R})(C)_i)_a &= \bigoplus_{d \in A} \mathbf{R}(C)_{(d,i)} \otimes_k S_{a-d} \\ &= \bigoplus_{d \in A} \bigoplus_{j \in \mathbb{Z}} \mathbf{R}(C_j)(0, -j)_{(d,i)} \otimes_k S_{a-d} \\ &= \bigoplus_{d \in A} \bigoplus_{j \in \mathbb{Z}} \bigoplus_{b \in A} (C_j)_{-b} \otimes_k \omega_{(b+d, i-j)} \otimes_k S_{a-d} \\ &= \bigoplus_{j \in \mathbb{Z}} \bigoplus_{d \in A} \bigoplus_{b \in A} (C_j)_b \otimes_k \omega_{(d, i-j)} \otimes_k S(-d)_{a-b} \\ &= \bigoplus_{j \in \mathbb{Z}} \bigoplus_{d \in A} (C_j \otimes_k \omega_{(d, i-j)} \otimes_k S(-d))_a \\ &= \bigoplus_{j \in \mathbb{Z}} (C_j \otimes_k (\mathbf{L} \circ \mathbf{R})(k)_{i-j})_a \\ &= ((C \otimes_k (\mathbf{L} \circ \mathbf{R})(k)))_i. \end{aligned}$$

This gives the first equality. As for the second:

$$\begin{aligned} (\mathbf{R} \circ \mathbf{L})(N)_{(a,i)} &= \bigoplus_{j \in \mathbb{Z}} (\mathbf{R}(\mathbf{L}(N)_j)(0, -j))_{(a,i)} \\ &= \bigoplus_{j \in \mathbb{Z}} \bigoplus_{d \in A} (\mathbf{L}(N)_j)_{-d} \otimes_k \omega_{(a+d, i-j)} \\ &= \bigoplus_{j \in \mathbb{Z}} \bigoplus_{d \in A} \bigoplus_{b \in A} N_{(b,j)} \otimes_k S_{-b-d} \otimes_k \omega_{(a+d, i-j)} \\ &= \bigoplus_{(a,j) \in A \times \mathbb{Z}} N_{(b,j)} \otimes_k (\mathbf{R} \circ \mathbf{L})(k)_{(a-b, i-j)} \\ &= N \otimes_k (\mathbf{R} \circ \mathbf{L})(k). \end{aligned}$$

For the last statement, observe that $(\mathbf{L} \circ \mathbf{R})(k)$ is the Koszul complex on the variables in S , and \square

$$d_L + (-1)^i d_C d_L + (-1)^{|c|} d_C$$

The derived category $D_{\text{DM}}(E)$ is obtained by inverting quasi-isomorphisms in $\text{DM}(E)$. Denote by $D_{\text{DM}}^b(E)$ the subcategory of $D_{\text{DM}}(E)$ given by objects with finitely generated homology.

We observe that the functors \mathbf{L} and \mathbf{R} preserve objects with finitely generated homology, as well as mapping cones and homotopy equivalences. It follows that \mathbf{L} and \mathbf{R} induce functors on bounded derived categories.

Corollary 3.7. *The induced maps*

$$\mathbf{L} : D_{\text{DM}}^b(E) \rightleftarrows D^b(S) : \mathbf{R}$$

are inverse equivalences.

Given any $A \times \mathbb{Z}$ -graded ring R , define a functor

$$\text{Fold} : \text{Com}(R) \rightarrow \text{DM}(R, a)$$

given by

$$(\text{Fold}(C))_i = \bigoplus_{j \in \mathbb{Z}} C_j(0, j - i)$$

and with i^{th} differential given by $(-1)^i d^C$.

3.2. Properties of the functors \mathbf{L} and \mathbf{R} . We record several properties of the toric BGG functors. Each is a direct analogue of a result in the classical BGG story and is proven in essentially the same way.

Proposition 3.8 (cf. [EFS03] Proposition 2.3). *Let $C \in \text{Com}(S)$, and let N be a finitely generated $A \times \mathbb{Z}$ -graded E -module, considered as an object in $\text{DM}(E)$ with trivial differential.*

(a) *We have*

$$H(\mathbf{R}(C))_{(a,j)} = H_j(C \otimes_S^{\mathbb{L}} k)_a.$$

In particular, if C is concentrated in degree 0, $H(\mathbf{R}(C))_{(a,j)} = \text{Tor}_j^S(C, k)_a$

(b) *Letting $w = \sum_{i=0}^n a_i \in A$, we have*

$$H_j(\mathbf{L}(N))_a = \text{Ext}_E^{\text{DM}}(k, N)_{(a,j)}.$$

♣♣♣ Michael: [The finiteness hypothesis on N is not present in [EFS03, Proposition 2.3], but I think it may be necessary here. See my comment in the proof. Please double check me on this; perhaps I'm wrong and the finiteness assumption isn't necessary!]

Proof. We can view the Koszul complex K on the variables of S as the complex of A -graded S -modules with homological degree j component $\bigoplus_{d \in A} S(-d) \otimes_k \omega_{(d,j)}$ and differential given

by multiplication by $\sum_{i=0}^n x_i \otimes e_i$. We have:

$$\begin{aligned}
\mathbf{R}(C)_{(a,j)} &= (\bigoplus_{i \in \mathbb{Z}} \mathbf{R}(C_i)(0, -i))_{(a,j)} \\
&= (\bigoplus_{i \in \mathbb{Z}} \bigoplus_{d \in A} (C_i)_{-d} \otimes_k \omega(d, -i))_{(a,j)} \\
&= \bigoplus_{i \in \mathbb{Z}} \bigoplus_{d \in A} (C_i)_{-d} \otimes_k \omega(d+a, j-i) \\
&= \bigoplus_{i \in \mathbb{Z}} \bigoplus_{d \in A} (C_i(-d))_a \otimes_k \omega(d, j-i) \\
&= (\bigoplus_{i \in \mathbb{Z}} C_i \otimes_S K_{j-i})_a \\
&= ((C \otimes_S K)_j)_a.
\end{aligned}$$

This equality identifies cycles in $\mathbf{R}(P)_{(a,j)}$ with j -cycles in $(C \otimes_S K)_a$, and similarly for boundaries. This proves (a). As for (b):

$$\begin{aligned}
(\mathbf{L}(N)_j)_a &= (\bigoplus_{d \in A} N_{(d,j)} \otimes_k S(-d))_a \\
&= \bigoplus_{d \in A} N_{(d,j)} \otimes_k S_{a-d} \\
&= \bigoplus_{d \in A} S_{-d} \otimes_k N_{(d+a,j)} \\
&= \bigoplus_{d \in A} S_{-d} \otimes_k N(-w+d, -n-1)_{(w+a,j+n+1)} \\
&= (\bigoplus_{d \in A} S_{-d} \otimes_k \omega(d, 0) \otimes_E N)_{(w+a,j+n+1)} \\
&= (\mathbf{R}(S) \otimes_E N)_{(w+a,j+n+1)} \\
&= \text{Hom}_E(\mathbf{R}(S)^\vee, N)_{(w+a,j+n+1)}.
\end{aligned}$$

The last equality follows since N is finitely generated. ♣♣♣ Michael: [The reason finitely generated is necessary is that pulling a direct sum out of the first component of Hom gives a product. Even the fact that we're in the graded setting doesn't seem to help us avoid this. Note that, in [EFS03], there is a homological grading, and each term of the complex $\mathbf{R}(S)^\vee$ is finitely generated, so this problem doesn't arise. Maybe we should include a comment about this?] As above, this equality identifies j -cycles in $\mathbf{L}(N)_a$ with cycles in $\text{Hom}_E(\mathbf{R}(S)^\vee, N)_{(w+a,j+n+1)}$. Finally, note that $\mathbf{R}(S)^\vee(w, n+1)$ is a free flag resolution of the residue field k , considered as an object in $\text{DM}(E)$ with trivial differential. \square

Corollary 3.9. *If M is a finitely generated A -graded S -module, $\dim_k H(\mathbf{R}(M)) < \infty$.*

Definition 3.10. A complex of free modules over S (resp. free differential E -module) is *linear* if each of its differentials can be expressed as a matrix whose entries are linear forms (resp. if it is of the form $(\bigoplus_{d \in A} \omega(d, 0), \partial)$, where ∂ is a matrix whose entries are linear forms).

Let $\text{Lin}(S)$ (resp. $\text{Lin}_{\text{DM}}(E)$) denote the category of linear free complexes of S -modules (resp. linear free differential E -modules).

Proposition 3.11 ([EFS03] Proposition 2.1). *We have induced equivalences*

$$\mathbf{L} : \text{Mod}(E) \xrightarrow{\cong} \text{Lin}(S)$$

and

$$\mathbf{R} : \text{Mod}(S) \xrightarrow{\cong} \text{Lin}_{\text{DM}}(E).$$

Proof. We prove the first functor is an equivalence; one uses a similar argument to prove the second functor is an equivalence. Let U be a k -vector space and $e \in V$. Identifying V with $\text{Hom}_k(W, k)$, we get an induced map

$$e : W \otimes U \rightarrow U$$

given by $w \otimes u \mapsto e(w)u$. Now, let

$$\cdots \xrightarrow{d} \bigoplus_{d \in A} S(d) \otimes_k N_{d,i} \xrightarrow{d} \bigoplus_{d \in A} S(d) \otimes_k N_{d,i-1} \xrightarrow{d} \cdots$$

be an object in $\text{Lin}(S)$. Let $N = \bigoplus_{d \in A, i \in \mathbb{Z}} N_{d,i}$. Define an E -module structure on N as follows. If $n \in N_{d,i}$ and $e \in V$, $e \cdot n = e(d(1 \otimes n)) \in \bigoplus_{d \in A} N_{d,i-1}$. We consider N as an $A \times \mathbb{Z}$ -graded E -module by defining $N_{(d,i)} = N_{d,i}$. The relation $d^2 = 0$ implies the relations N must satisfy to be an E -module. ♣♣♣ Michael: [Check this last statement, and check that the proof of the second equivalence really is the same.] \square

♣♣♣ Daniel: [We need to state and prove reciprocity theroem. For that, we'll need to define an injective resolution of differential modules. That's not difficult, but it does need to be added.]

4. PUSHFORWARD OF DIFFERENTIAL MODULES

♣♣♣ Daniel: [This section is in rough shape.] Here we give the explicit construction of the pushforward functor τ_* . We also prove results about a pushforward functor $\text{DM}(X) \rightarrow \text{DM}(k)$.

Lemma 4.1. *Let X be a toric variety over k with Cox ring S , graded by $\text{Pic}(X)$. Let E be the exterior algebra on the dual variables, with the $\text{Pic}(X) \oplus \mathbb{Z}$ -grading described above.*

- (1) *There is an additive functor: $\pi_* : \text{DM}(X) \rightarrow \text{DM}^\circ(k)$ which preserves exactness and where $|\pi_* \mathcal{D}| = \bigoplus_{i=0}^{\dim X} H^i(X, |\mathcal{D}|)$, namely: the underlying module of $\pi_* \mathcal{D}$ is the total sheaf cohomology of the underlying module of \mathcal{D} .*
- (2) *There is an additive functor: $\tau_* : \text{DM}_{\text{free}}(\mathbb{P}_E^n) \rightarrow \text{DM}^\circ(E)$ which preserves exactness and where the underlying module of $\tau_* \mathcal{D}$ satisfies: If $|\mathcal{D}| = \bigoplus_{\ell \in \text{Pic}(X)} \mathcal{E}_\ell \otimes_k E(\ell; 0)$ then*

$$(3) \quad |\tau_* \mathcal{D}| = \bigoplus_{i=0}^{\dim X} \bigoplus_{\ell \in \text{Pic}(X)} H^i(\mathcal{E}_\ell) \otimes_k E(\ell; -i).$$

Proof. ♣♣♣ Daniel: [Some notational bugs here. I ran out of time but will fix soon. All ideas correct.] We first prove (1). Start with a differential X -module whose underlying sheaf is \mathcal{F} . By unfolding, we can realize this in $\mathbf{D}(\mathbb{P}_E^n)$ as

$$\cdots \xrightarrow{\partial} \mathcal{F} \xrightarrow{\partial} \mathcal{F} \xrightarrow{\partial} \mathcal{F} \xrightarrow{\partial} \cdots$$

Let $C^0 \rightarrow C^1 \rightarrow \dots \rightarrow C^n$ be the Čech resolution for \mathcal{F} , with C^i an S -module. We build a double complex F_j^i where $F_j^i = C^i$ (in particular F_j^i only depends on j); the columns are Čech resolutions and the horizontal maps are localizations of the differential ∂ .

Each column of the double complex is the same, and taking vertical homology in the i 'th spot gives $H^i(X, \mathcal{F})$. By choosing a splitting of $H^i(\mathcal{E}_\ell)$ into the Čech complex for \mathcal{F} , we get a splitting for the vertical homology in this double complex. We can then apply [EFS03, Lemma 3.5], to get a complex of E -modules:

$$\dots \xrightarrow{\partial} G \xrightarrow{\partial} G \xrightarrow{\partial} G \xrightarrow{\partial} \dots$$

where G in column 0 is

$$G = \bigoplus_{i=0}^{\dim X} H^i(F_i^*) = \bigoplus_{i=0}^{\dim X} H^i(\mathcal{F}).$$

A different splitting will induce the same complex, up to homotopy (since both would be homotopic to the original total complex, again by [EFS03, Lemma 3.5]). Applying Remark ?? in reverse, G induces a well-defined element of $\mathrm{DM}^\circ(k)$ as desired.

Exactness is nearly immediate from the construction. By construction, the i th homology of the complex

$$\dots \xrightarrow{\partial} \pi_* \mathcal{D} \xrightarrow{\partial} \pi_* \mathcal{D} \xrightarrow{\partial} \pi_* \mathcal{D} \xrightarrow{\partial} \dots$$

in $\mathbf{D}(k)$ is the hypercohomology of the complex

$$\dots \xrightarrow{\partial} \mathcal{D} \xrightarrow{\partial} \mathcal{D} \xrightarrow{\partial} \mathcal{D} \xrightarrow{\partial} \dots$$

But by the standard hypercohomology spectral sequence $E_2^{i,j} = H^i(H^j -) \Rightarrow H^{i+j}(-)$, if (the unfolded complex for) \mathcal{D} is exact then so is (the unfolded complex for) $\pi_* \mathcal{D}$.

The proof of (2) is nearly identical as the proof of (1), though we need to track the gradings in more detail. Starting with an object $\mathcal{F} := \bigoplus_\ell \mathcal{E}_\ell \otimes_k E(\ell; 0)$ from $\mathrm{DM}_{\mathrm{free}}(\mathbb{P}_E^n)$, we get the unfolded complex:

$$\dots \xrightarrow{\partial} \bigoplus_\ell \mathcal{E}_\ell \otimes_k E(\ell; -1) \xrightarrow{\partial} \bigoplus_\ell \mathcal{E}_\ell \otimes_k E(\ell; 0) \xrightarrow{\partial} \bigoplus_\ell \mathcal{E}_\ell \otimes_k E(\ell; 1) \xrightarrow{\partial} \dots$$

For each ℓ , we take a separate Čech resolution of \mathcal{E}_ℓ , tensor with $- \otimes_k E(\ell; 0)$, and then take the direct sum of these to get one of our vertical columns, C_0^* . We then let $C_i^* = C_0^* \otimes_E E(0; i)$. We again build a double complex from these, where the columns are twists of Čech resolutions, and the horizontal maps are localizations of the differential ∂ .

Each column is the same (up to a twist of the E -part), and taking vertical homology in the i 'th spot gives $H(C_j^*) = \bigoplus_\ell H^i(\mathcal{E}_\ell) \otimes_k E(\ell; -j)$. Since the vertical differentials are defined entirely over k , we can still choose a splitting of the homology into the Čech complex, and thus obtain a splitting for the vertical homology in this double complex. We apply [EFS03, Lemma 3.5], to get a complex of E -modules:

$$\dots \xrightarrow{\partial} G(0; -1) \xrightarrow{\partial} G \xrightarrow{\partial} G(0; 1) \xrightarrow{\partial} \dots$$

where G in homological degree 0 is

$$G = \bigoplus_{i=0}^{\dim X} H^i(F_i^*) = \bigoplus_{i=0}^{\dim X} \bigoplus_{\ell \in \mathrm{Pic}(X)} H^i(\mathcal{E}_\ell) \otimes_k E(\ell; -i).$$

We thus obtain a well-defined element of $\mathrm{DM}^\circ(E)$ as desired. To check exactness, we can consider the commutative square

$$\begin{array}{ccc} \mathrm{DM}_{\mathrm{free}}(\mathbb{P}_E^n) & \longrightarrow & \mathrm{DM}(\mathbb{P}_k^n) \\ \tau_* \downarrow & & \downarrow \pi_* \\ \mathrm{DM}^\circ(E) & \longrightarrow & \mathrm{DM}^\circ(k) \end{array}$$

where the horizontal arrows simply forget the E -module structure. Since forgetful maps respect exactness, exactness for part (2) follows from exactness for part (1). \square


Example 4.2. Let's work on \mathbb{P}^1 . Let $\mathcal{D} := \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}^2(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(-2)$ with the Koszul differential on \mathcal{D} , so that \mathcal{D} is exact. Let $C^0 \rightarrow C^1$ be the Cech resolution of \mathcal{D} . Under the construction above, we end up with a total complex $\cdots \xrightarrow{\partial} G \xrightarrow{\partial} G \xrightarrow{\partial} G \xrightarrow{\partial} \cdots$ where

$$G \cong H^0(\mathbb{P}^1, \mathcal{O}) \oplus H^1(\mathbb{P}^1, \mathcal{O}(-2)).$$

While G is isomorphic to the total cohomology of C^\bullet , note that the H^0 and H^1 come from different copies of C^\bullet in the total complex. Namely, if we are looking at the copy of G in position 0, then the H^0 is the 0'th homology of the C^\bullet in column 0 whereas the H^1 part is the 1'st homology of C^\bullet in column 1. That is to say, if we write C_i^\bullet for the Cech complex in column i , and we write G_i for the copy of G in position i , then:

$$G_i = H^0(C_i^\bullet) \oplus H^1(C_{i+1}^\bullet).$$

So to get the map $H^1 \rightarrow H^0$ we would require at least 3 copies of the Cech complex. This explains why we can't just define the differential as "the pushforward of the map ∂ ".

Remark 4.3. Rouquier also uses differential modules in [Rou06, Section 4] to provide an analogue of the BGG correspondence for non-graded polynomial and exterior algebras. 

Michael: [Also mention Baranovsky here.]

5. TATE MODULES

Let X be a projective toric variety and let \mathcal{E} be a coherent sheaf on X . Our main result is a proof that, over the exterior dual of the Cox ring of X , there exists a free differential module $\mathbf{T}(\mathcal{E})$ which combines all of the sheaf cohomology groups of \mathcal{E} into a single exact differential module.

Theorem 5.1. *Let X be a simplicial toric variety and let \mathcal{E} be a coherent sheaf on X . There exists a free, exact differential module $(\mathbf{T}(\mathcal{E}), \partial)$ whose underlying module is:*

$$\mathbf{T}(\mathcal{E}) = \bigoplus_{i=0}^{\dim X} \bigoplus_{\ell \in \mathrm{Pic}(X)} H^i(X, \mathcal{E}(-\ell)) \otimes_k \omega_E(\ell; -i).$$

The key idea in the proof is to realize $\mathbf{T}(\mathcal{E})$ by developing a theory of pushforwards for differential \mathcal{O}_X -modules. The following push-pull diagram summarizes our definition of the Tate module:

$$\begin{array}{ccc}
\mathrm{Coh}(X) & \xrightarrow{\kappa^*} & \mathrm{DM}_{\mathrm{free}}(X_E) \\
& \searrow \mathbf{T} & \downarrow \tau_* \\
& & \mathrm{DM}^\circ(E)
\end{array}$$

The arrow κ^* sends the sheaf \mathcal{E} to the Koszul complex of $\widehat{\mathcal{E}}$ with respect to the variables x_0, x_1, \dots, x_n , which can naturally be understood as a differential \mathcal{O}_{X_E} -module. The vertical arrow τ_* is the natural (derived) pushforward functor for these categories of differential modules. So in essence, as with an integral transform, we are pulling back \mathcal{E} to X_E ; then tensoring with the Koszul complex; then pushing forward.

This process recovers the known Tate resolutions on projective space and on products of projective spaces (see Example 5.6 for the precise statement), but via a very different approach. More importantly, this definition allows us to highlight subtle exactness properties of these Tate modules, which are controlled by the irrelevant ideal of S .

We prove two other key results in this section. Theorem 5.8 shows that the Tate module satisfies even stronger exactness properties, which are encoded by the irrelevant ideal of X . Theorem 5.9 gives an algebraic characterization of the Tate module, which is more closely related to the approach of [EFS03, EES15].

We start making this precise. Let $\mathrm{Mod}(X_E)$ denote the category of $\mathcal{O}_X \otimes_k E$ -modules which are graded with respect to the E -grading. We let $\mathrm{DM}(X_E)$ be the category of differential $\mathcal{O}_X \otimes_k E$ -modules which have degree $(0; 1)$ in the $\mathrm{Pic}(X) \oplus \mathbb{Z}$ grading on E .¹ We first observe that the Koszul complex of x_1, \dots, x_n is naturally an object in $\mathrm{DM}(X_E)$. To get the grading right, we start with the $\mathcal{O}_X \otimes_k E$ module:

$$\bigoplus_{d \in \mathrm{Pic}(X)} \mathcal{O}_X(d) \otimes_k \omega_E(-d; 0).$$

and we endow this with the differential given by $\sum_{i=1}^n x_i \otimes e_i$. We refer to this differential $\mathcal{O}_X \otimes_k E$ -module as \mathcal{K} . In a similar way, for any subset $I \subseteq \{1, \dots, n\}$, we can endow the same underlying module with the differential $\sum_{i \in I} x_i \otimes e_i$. We refer to this as \mathcal{K}_I . There is a natural quotient $\mathcal{K} \rightarrow \mathcal{K}_I$ obtained by sending $e_i \mapsto 0$ for $i \notin I$. ♣♣♣ Daniel: [Check this.]

Definition 5.2. We define $\kappa^* : \mathrm{Coh}(X) \rightarrow \mathrm{DM}(X_E)$ as the composition of functors:

$$\begin{array}{ccccc}
& & \kappa^* & & \\
& \nearrow & & \searrow & \\
\mathrm{Mod}(X) & \xrightarrow{\otimes_k E} & \mathrm{Mod}(X_E) & \xrightarrow{\otimes \mathcal{K}} & \mathrm{DM}(X_E)
\end{array}$$

$$\mathcal{E} \longmapsto \mathcal{E} \otimes_k E \longmapsto (\mathcal{E} \otimes_k E) \otimes \mathcal{K}$$

We define κ_I^* similarly, but tensoring by \mathcal{K}_I in the second step.

Remark 5.3. We can think of $\kappa^* \mathcal{E}$ in more concrete terms as follows. The underlying module of $\kappa^* \mathcal{E}$ is $\bigoplus_{d \in \mathrm{Pic}(X)} \mathcal{E}(d) \otimes_k \omega_E(-d; 0)$ and the differential is multiplication by $\sum_{i=1}^n x_i \otimes e_i$.

¹We could also consider differential modules of degree d for any $d \in \mathrm{Pic}(X) \oplus \mathbb{Z}$, but degree $(0; 1)$ is the only one we will require.

Theorem 5.1 will follow easily once we have constructed the pushforward functor τ_* , which sends exact differential modules to exact differential modules. This functor also determines a functor on the underlying submodules $\tau_*^{\text{Mod}} : \text{Mod}(X_E) \rightarrow \text{Mod}(E)$ and such a functor is in term determined by its effect on objects of the form $\mathcal{E} \otimes_k \omega_E(d; j)$. In our case, this functor will essentially send a sheaf to its sheaf cohomology, with appropriate E -gradings:

$$\tau_*^{\text{Mod}} : \mathcal{E} \otimes_k \omega_E(d; j) \mapsto \bigoplus_{i=0}^{\dim X} H^i(X, \mathcal{E}) \otimes_k E(d; j - i).$$

The details involve homological perturbation applied to complexes of the form $C^\bullet \otimes_k E$, where C^\bullet is the Cech complex of a sheaf on X , and these will be covered in §4. However, the main result that we need is captured by the following proposition:

Proposition 5.4. *There is an additive functor τ_* from $\text{DM}(X_E)$ to the homotopy category of $\text{DM}(E)$, which preserves exactness and where the induced functor $\tau_*^{\text{Mod}} : \text{Mod}(X_E) \rightarrow \text{Mod}(E)$ on underlying modules is determined by*


$$\mathcal{E} \otimes_k \omega_E(d; j) \mapsto \bigoplus_{i=0}^{\dim X} H^i(X, \mathcal{E}_\ell) \otimes_k E(d; j - i).$$

We postpone the proof until §4, but note that it immediately implies Theorem 5.1.

Definition 5.5. We define $(\mathbf{T}(\mathcal{E}), \partial)$ as the differential E -module $\tau_* \kappa^*(\mathcal{E} \otimes \mathcal{K})$. For any $I \subseteq \{1, 2, \dots, n\}$, we define $(\mathbf{T}(\mathcal{E}), \partial_I)$ as the differential E -module $\tau_* \kappa^*(\mathcal{E} \otimes \mathcal{K})$.

Proof of Theorem 5.1. The homology of \mathcal{K} is supported on $V(x_1, \dots, x_n)$ which is the empty set. It follows that the same statement holds for $\kappa^* \mathcal{E}$. In other words, $\kappa^* \mathcal{E}$ is exact. Thus $\tau_* \kappa^* \mathcal{E} = \mathbf{T}(\mathcal{E})$ is exact. Proposition 5.4 confirms that the underlying module of $\tau_* \kappa^* \mathcal{E}$ is as stated. \square

We next observe that when $X = \mathbb{P}^n$ this recovers the Tate resolution of [EFS03].

Example 5.6. Let $X = \mathbb{P}^n$ and let \mathcal{E} be a coherent sheaf on \mathbb{P}^n . Let M be a graded S -module such that $\widetilde{M} = \mathcal{E}$. Let $\mathbf{T}'\mathcal{E}$ denote the differential module obtained from the Tate resolution as defined in [EFS03]. By Remark 3.4, it suffices to show that $\mathbf{T}'\mathcal{E}$ is isomorphic to the Tate module $\mathbf{T}\mathcal{E} = \tau_* \kappa^* \mathcal{E}$ as defined as above. For starters, both Tate resolutions agree with $\mathbf{R}M$ in degrees $\geq d$ for some d . Write T for the tail of $\tau_* \kappa^* \mathcal{E} \rightarrow \mathbf{R}M_{\geq d}$ and T' for the tail of $\mathbf{T}'\mathcal{E} \rightarrow \mathbf{R}M_{\geq d}$. Then T and T' are both minimal free resolutions of the differential module $\mathbf{R}M_{\geq d}$ and thus they are isomorphic by the uniqueness of minimal free resolutions of differential modules.  Daniel: [Need to add reference to the equivalence between $\text{DM}(E)$ and $\text{Com}(E)$ in this case.]

5.1. Exactness properties of $\mathbf{T}(\mathcal{E})$. Using nearly identical methods reveals deeper exactness properties of these Tate resolutions.

Definition 5.7. Given $I \subseteq \{0, 1, \dots, n\}$, we say that I is **irrelevant** if the ideal $\langle x_i \text{ where } i \in I \rangle$ contains the irrelevant ideal. Following Batyryev (see [CLS, p. 304]), we say that I is **primitive** if it is irrelevant but if no proper subset of I is irrelevant.

One crucial feature of Tate resolutions over other toric varieties is that they satisfy more complicated exactness properties, which are encoded by the irrelevant ideal.

Theorem 5.8 (Exactness Properties). *If $I \subseteq \{0, 1, \dots, n\}$ is irrelevant, then $(\mathbf{T}(\mathcal{E}), \partial_I)$ is exact.*

Proof. The proof of Theorem 5.1 goes through almost verbatim. The homology of \mathcal{K}_I is supported on P_I . But I is irrelevant which implies that \mathcal{K}_I is exact, which implies that κ_I^* is exact, which implies that $\tau_*\kappa_I^*\mathcal{E}$ is exact. \square

These nuanced exactness properties were not present for the Tate resolutions in [EFS03]: the irrelevant ideal equals the maximal ideal on projective space, so there are no interesting choices for I in that case. But for products of projective spaces, these exactness properties are equivalent to the the exact “rows and columns” which played a key role in [EES-products].

5.2. Algebraic characterization of $\mathbf{T}(\mathcal{E})$. The exactness properties lead to an algebraic characterization of $\mathbf{T}(\mathcal{E})$ as a differential module. The following theorem shows that, if M is a multigraded S -module, then the Tate resolution of \widetilde{M} can be determined by the algebraic data of: $\mathbf{R}(M)$ and the exactness properties of Theorem 5.8. This is something like a parallel of the fact that a toric variety may be determined by two pieces of algebraic data: its multigraded Cox ring S and its irrelevant ideal. Namely, $\mathbf{R}(M)$ is a differential module which is entirely determined by the Cox ring S , but passing from $\mathbf{R}(M)$ to $\mathbf{T}(\widetilde{M})$ requires the exactness properties, which are determined by the irrelevant ideal.

Theorem 5.9 (Algebraic Characterization of $\mathbf{T}(\mathcal{E})$). *Up to isomorphism, $\mathbf{T}(\widetilde{M}, \partial)$ is the unique minimal, free differential module which equals $\mathbf{R}M_{\geq d}$ for all $d \gg 0$ and which satisfies the exactness properties of Theorem 5.8.*

Remark 5.10. This theorem quickly implies that the Tate resolutions of [EFS03] and [EES15] agree with the Tate modules constructed in this paper.

The proof involves some facts about the minimal primes of the irrelevant ideal of X .

Lemma 5.11. *Let $\mu : \text{Pic}(X) \rightarrow \mathbb{Z}$ be a linear functional such that $\mu \geq 0$ is one of the minimal defining halfspaces of $\text{Eff}(X)$. There exists a primitive subset $I \subseteq \{1, \dots, n\}$ such that $\mu(\deg x_i) > 0$ for all $i \in I$.*

Proof. The nef cone $\overline{\text{NE}}(X)$ belongs to the effective cone $\text{Eff}(X)$, so the intersection of $\mu = 0$ and $\overline{\text{NE}}(X)$ will lie inside of some (and possibly more than one) facet of $\overline{\text{NE}}(X)$. Let $\tau : \text{Pic}(X) \rightarrow \mathbb{Z}$ be the defining functional of that facet. We thus have:

$$\text{Eff}(X) \setminus \{\mu = 0\} \supseteq \overline{\text{NE}}(X) \setminus \{\tau = 0\}.$$

By [CLS, Proof of Theorem 6.4.11] (see also the citations to Cox-von Resse and Kresch from CLS...), we conclude that the set I of i such that $\tau(\deg x_i) > 0$ forms a primitive collection. It follows that $\mu(\deg x_i) > 0$ for each $i \in I$. Namely, $\mu(\deg x_i)$ since $\deg x_i \in \text{Eff}(X)$ and $\tau(\deg x_i) \neq 0 \Rightarrow \mu(\deg x_i) \neq 0$ by the displayed inclusion. \square

Example 5.12. Consider the Hirzebruch surface \mathbb{F}_3 with Cox ring $S = k[x_0, x_1, x_2, x_3]$, irrelevant ideal $(x_0, x_2) \cap (x_1, x_3)$ and degrees of the variables $\deg(x_0) = \deg(x_2) = (1, 0)$, $\deg(x_1) = (-3, 1)$ and $\deg(x_3) = (0, 1)$. The defining halfspaces of $\text{Eff}(X)$ are determined are give $\mu_i \geq 0$ for $i = 1, 2$ where $\mu_i : \mathbb{Z}^2 \rightarrow \mathbb{Z}$ is $\mu_1(a, b) = a + c$ and $\mu_2(a, b) = b$. Note that μ_1 is strictly positive on x_0, x_2 while μ_2 is strictly positive on x_1, x_3 .

Sketch of proof of Theorem 5.9. Fix some $d \in \text{Pic}(X)$ which is sufficiently large, in a sense to be made precise as we proceed. Fix some facet $\text{Eff}(X)$, defined by a functional $\mu :$

$\text{Pic}(X) \rightarrow \mathbb{Z}$ which is nonnegative on $\text{Eff}(X)$. By Lemma 5.11, we can find a primitive collection $I \subseteq \{0, 1, \dots, n\}$ where $\mu(\deg x_i)$ is strictly positive for all $i \in I$.

We use μ to flatten down to a \mathbb{Z} -grading on E . Namely, we define $\deg_\mu(e_i) = -\mu(x_i) \in \mathbb{Z}$. With respect to the \deg_μ -grading, we can apply Lemma 2.21 to $(\mathbf{T}(\mathcal{E}), \partial_I)$ to obtain:

$$(\mathbf{T}(\mathcal{E})^{\text{I-tail}}, \partial_I) \rightarrow (\mathbf{T}(\mathcal{E})^{\text{I-head}}, \partial_I)$$

where the I -head consists of all factors of the form $\omega_E(a; b)$ such that $\deg_\mu(a) \leq e$; the I -tail consists of all factors of the form $\omega_E(a; b)$ such that $\deg_\mu(a) > e$; and the differentials are the restrictions of ∂_I .

We next claim that: the I -tail of the Tate resolution, and the restriction of ∂ to this I -tail, can be recovered entirely from the I -head, its differential.

Since Lemma 2.21 implies that the I -tail is the minimal free resolution of the I -head, the uniqueness of minimal free resolutions implies that we can recover the tail (and its differential ∂_I) entirely from the head. This recovers the free module $\mathbf{T}(\mathcal{E})^{\text{I-tail}}$ as well as the part of the restriction of ∂ involving variables e_i with $i \in I$. We next claim that we can also recover the restriction of $\partial_I^C := \partial - \partial_I$ to the tail from ∂ on the head.

Since $\deg_\mu(x_i) \geq 0$ for all $1 \leq i \leq n$, it follows that ∂_I^C restricts to an endomorphism $\partial_I^C|_{\text{head}}$ of the I -head. Since we have a map $(\mathbf{T}(\mathcal{E}), \partial_I) \rightarrow (\mathbf{T}(\mathcal{E})^{\text{I-head}}, \partial_I)$, we can apply Proposition 2.6 to lift this to an endomorphism $(\mathbf{T}(\mathcal{E}), \partial_I)$. But by uniqueness of lifts, we see that any lift will agree with ∂_I^C , up to homotopy. We have thus recovered the entire Tate module $(\mathbf{T}(\mathcal{E}), \partial)$ from $(\mathbf{T}(\mathcal{E})^{\text{I-head}}, \partial)$, which is like “half” of the Tate module.

Now, we will iterate this argument, with two replacements. We first replace the total Tate module $(\mathbf{T}(\mathcal{E}), \partial)$ by the head half of the Tate module $(\mathbf{T}(\mathcal{E})^{\text{I-head}}, \partial)$. We then replace τ by another defining facet. Iterating in this way, we eventually conclude that, by using exactness properties, the full Tate module can be recovered entirely from the restriction of the Tate module to summands of the form $\omega_E(a; b)$ where $\deg_\mu(a) \geq \deg_\mu(d)$ for all functionals μ defining $\text{Eff}(X)$. We conclude that, using the exactness relations, the entire Tate resolution can be recovered from the Tate module in very positive degrees. That is, $(\mathbf{T}(\mathcal{E})_{\geq d}, \partial)$ determines the entire Tate resolution, for any degree d .

Finally, we want to observe that all of the previous arguments go through with a global upper bound on the $\text{Pic}(X)$ -grading. ♣♣♣ Daniel: [I'm struggling to phrase this precisely, but I don't think there are any serious mathematical obstacles here. Big idea: everything of degree $\leq d'$ only depends on a finite window of degrees. And for d large enough, we can assume that for degrees between d and d' , the Tate resolution agrees with R.M.] For a differential module F over E and degree $e \in \text{Pic}(X)$ we will write $F_{(e;*)}$ for the differential module obtained by considering in the $(e; j)$ part of F , for all j . Since we are working with a differential of degree $(0; 1)$, this is a differential module of k -vector spaces. And of course any differential module F is entirely determined by its $(e : *)$ -strands.

Moreover, since the elements of E have $\text{Pic}(X)$ -degrees between 0 and $w_E := \sum_i \deg(x_i)$, we see that the strand $F_{(e;*)}$ only depends on summands of the form $\omega_E(a; j)$ where $e \geq a \geq e + w_E$. ♣♣♣ Daniel: [Check the signs!!!] It follows that the degree $(e; *)$ strand of F is determined by the subquotient of F obtained by summing only over the free summands $\omega_E(-a; j)$ where $e \geq a \geq e + w_E$.

So if we write $F_{\geq d}^{\leq d'}$ for the natural subquotient of F determined by restricting attention to summands of the form $\omega_E(a; j)$ where $d \leq -a \leq d'$, then $F_{\geq d}^{\leq d'}$ will have the same $(e; *)$ -strand as F , as long as $\clubsuit\clubsuit\clubsuit$ Daniel: [some condition on e, d, d' and $w_E \dots$]

In summary, the degree $(e; *)$ strands of $(\mathbf{T}(\mathcal{E}), \partial)$ for $e \leq d' + w_E$ will be determined by the degree $(e; *)$ strands of $(\mathbf{T}(\mathcal{E})_{\geq d}^{\leq d'}, \partial)$ for any d , as long as $d' - d \geq w_E$ (or something similar). But for any finite window of degrees, like the window between d and $d + w_E$, we can choose $d \gg 0$ so that \mathcal{E} has no higher cohomology in this window, and in this case $(\mathbf{T}(\mathcal{E})_{\geq d}^{\leq d'}, \partial)$ is just $(\mathbf{R}M, \partial)_{\geq d}^{\leq d'}$.

□

Remark 5.13. Running this argument in reverse gives a sketch of an algorithm for computing Tate resolutions, which would largely parallel the algorithm of [EFS03] and [eisenbud-decker]. There are some delicate issues to address, though, as any algorithm would only work with finite windows of $\mathbf{R}(M_{\geq d})$, and so one would need to understand how such truncations affect the process of computing minimal free resolutions, lifting endomorphisms, and more.

6. APPLICATIONS

6.1. Toric Syzygy Theorem.

Definition 6.1. We define the exterior irrelevant ideal of E as the ideal generated by monomials $e_{i_1}e_{i_2} \cdots e_{i_s}$ such that $x_{i_1}x_{i_2} \cdots x_{i_s}$ lies in the irrelevant ideal of S .

Theorem 6.2. *Let M be a finitely generated, graded S -module. Then M admits a virtual resolution of length at most $\dim X$. More specifically, for any sufficiently ample degree $d \in \text{Pic}(X)$, the projective dimension of $M_{\geq d}$ is at most $\dim X$.*

Proof. By applying Theorem 3.8(a) in the case $P = M_{\geq d}$, it will suffice to prove that the homology of $\mathbf{R}M_{\geq d}$ lies in degrees (a, j) with $-\dim X \leq j \leq 0$. We let $(F, \partial) = \mathbf{R}M_{\geq d}$. By the “lift” of an element $\alpha \in H(F, \partial)$ we will mean an element of F , lying in the kernel of ∂ and whose image in $H(F, \partial)$ is α . Let $B \subseteq E$ be the exterior irrelevant ideal. We will show that any lift of any element of $H(F, \partial)$ lies inside $B \cdot F$.

By Theorem 3.8(a), the homology of $\mathbf{R}M$ is supported in finitely many distinct degrees. It follows that, for any degree e which is away from those finitely many degrees, the strand $(\mathbf{R}M, \partial)_{e,*}$ will be exact.

Suppose that we choose some degree d which is greater than all of the degrees a where (a, j) is in the support of $H\mathbf{R}M$. We consider $(\mathbf{R}M_{\geq d}, \partial) \rightarrow (\mathbf{R}M, \partial)$. Let ζ be the lift of an element of $H(\mathbf{R}M_{\geq d}, \partial)$. Since this is a map of differential modules, ζ maps to a cycle of $(\mathbf{R}M, \partial)$; but for degree reasons, this cannot be a homology element, and thus ζ lies in the image of ∂ . More precisely, we can see that if $\omega = \sum_{i=1}^n \deg(x_i)$, then ζ must be a boundary of $(\mathbf{R}M_{\geq d-\omega}, \partial)$.

Now we choose d to be sufficiently ample so that \widetilde{M} has no higher cohomology in all degrees in the range between $d - \omega$ and d . It follows that, for any degree e in this range, and for any subset $I \subseteq \{1, \dots, n\}$: $(\mathbf{R}M, \partial_I)_{e,*}$ equals $(\mathbf{T}\mathcal{E}, \partial_I)_{e,*}$ and this has no higher cohomology. (We use Theorem 5.8 here.)

Recalling that $F = \mathbf{R}M_{\geq d}$, we consider (F, ∂_I) . Let ζ be the lift of a homology element in some degree $(e, *)$ with $d - \omega \leq e \leq d$. By the previous paragraph, we know that ζ lies in

the image of the differential ∂_I on $(\mathbf{R}M, \partial_I)$. In particular, if $L_I = \langle e_i \text{ where } i \in I \rangle$, then ζ lies in $L_I \cdot F$.

Next we observe that the full differential $\partial : F \rightarrow F$ is the sum of two differentials. Since we have truncated in a very positive degree, we have $\partial = \partial_I + \partial_{I^c}$ each of which is a differential by ♣♣♣ Daniel: [We need to add this somewhere earlier, but it's purely formal.] Using a spectral sequence argument ♣♣♣ Daniel: [fill in], we conclude that any lift of a homology element of (F, ∂) must also lie in $L_I \cdot F$.

Now, this has to hold for every minimal prime of the irrelevant ideal. The intersection of L_I over all such sets I is, by definition essentially, our exterior irrelevant ideal. It follows that the lift of any element of $H(F, \partial)$ lies in $B \cdot F$, as claimed. Recall that F is a direct sum of modules over the form $\omega_E(a; 0)$, which is nonzero only in degree (a, j) where auxiliary degrees $-n \leq j \leq 0$. By Lemma 6.3, every element of L_I is a product of at least r variables. We thus conclude that the homology of (F, ∂) lies entirely in degrees of the form (a, j) with $-n + r \leq i \leq 0$. Finally, we have that $\dim X = n - r$, completing the proof. \square

Lemma 6.3. *Let X be a projective simplicial toric variety and let $r = \text{rank Pic}(X)$. Every minimal generator of the irrelevant ideal of X is a product of r distinct monomials.*

Proof. The irrelevant ideal is generated by monomials corresponding to the product of the rays in the fan of X that are complements to a max face [CLS, p. 207]. Since r is the number of rays minus the dimension of the ambient lattice, it follows that each generator is a product of r variables. \square

♣♣♣ Daniel: [A remark: Imagine we have fixed S and M but not X . (In other words, imagine we have to toric varieties with the same cox ring S . The choice of d depends on X . But the statement about the length of the minimal free resolution of X does not. In other words, all we seem to need is that d is sufficiently ample for SOME X whose Cox ring S . This is a much less restrictive condition, which might be equivalent to being sufficiently far from the boundary of the effective cone. But I'm not sure.]

7. THE TATE RESOLUTION AND A_∞ -OPERATIONS ON SHEAF COHOMOLOGY

♣♣♣ Michael: [I don't know if we should include this, because I see no application...]

7.1. A_∞ operations and the tensor coalgebra. Let R be a ring and M an R -module. The tensor coalgebra $T^c(M)$ is the sum of all the tensor powers of M , equipped with the comultiplication

$$x_1 \otimes \cdots \otimes x_r \mapsto \sum_{i=1}^r (x_1 \otimes \cdots \otimes x_i) \otimes (x_{i+1} \otimes \cdots \otimes x_r) \in T^c(M) \otimes T^c(M).$$

We recall that a *coderivation* on an R -coalgebra (C, δ) is an R -linear map $d : C \rightarrow C$ satisfying the co-Leibniz rule $\Delta \circ d = (d \otimes 1 + 1 \otimes d) \circ \Delta$. A *dg-coalgebra over R* is a \mathbb{Z} -graded R -coalgebra equipped with a square zero coderivation of degree -1 .

Suppose S is a \mathbb{Z} -graded R -module. Then $T^c(S)$ inherits a \mathbb{Z} -grading, and the datum of an A_∞ -algebra structure on S is exactly the datum of a square 0 coderivation on $T^c(S)$. In more detail: any coderivation d on $T^c(S)$ is determined by its projection $T^c(S) \xrightarrow{d} T^c(S) \twoheadrightarrow S$ onto the first tensor power. So, a coderivation is the same thing as a degree -1 map $d : T^c(S) \rightarrow S$.

The components of this map are the higher multiplications on S , and the requirement that $d^2 = 0$ encodes the A_∞ relations.

7.2. Perturbations and A_∞ -structures. Let R be a ring, and suppose A is an augmented, connected, unital differential graded R -algebra. Moreover, suppose A contracts onto its homology, i.e. there are maps

$$\beta : A \rightrightarrows H_*(A) : \alpha$$

of graded R -modules such that $\beta\alpha = \text{id}_{H_*(A)}$ and $\alpha\beta$ is homotopic to id_A . We get induced maps of tensor coalgebras

$$(4) \quad T^c(\beta) : T^c(A) \rightrightarrows T^c(H_*(A)) : T^c(\alpha).$$

We observe that $T^c(A)$ is again a dg-coalgebra with differential induced by the one on A :

$$x_1 \otimes \cdots \otimes x_r \mapsto \sum_{i=1}^r \pm x_1 \otimes \cdots \otimes d(x_i) \otimes \cdots \otimes x_r$$

(note: this is the coderivation determined by the degree -1 map $T^c(A) \rightarrow A$ given by $x_1 \otimes \cdots \otimes x_r \mapsto \sum_{i=1}^r \pm d(x_i)$). We will abuse notation and call this induced differential d . The homotopy between $\alpha\beta$ and id_A induces a homotopy between $T^c(\alpha)T^c(\beta)$ and $\text{id}_{T^c(A)}$. That is, the maps in (4) give a contraction of $T^c(A)$ onto $T^c(H_*(A))$ (where the latter has no differential).

Now, suppose we perturb the differential on $T^c(A)$ via some second differential ∂ so that $d + \partial$ remains a coderivation. For instance, we could do this by perturbing the differential on A so that A remains a dga. The so-called Coalgebra Perturbation Lemma ([HK91] (2.1*)) says that the differential on $T^c(H_*(A))$ obtained from the usual perturbation lemma is a coderivation. We therefore obtain an induced A_∞ structure on $H_*(A)$.

Example 7.1. Suppose R is a field. Then one of course one can always give an R -linear contraction of A onto $H_*(A)$. We can perturb the differential on $T^c(A)$ by adding in the *bar differential*

$$a_1 \otimes \cdots \otimes a_r \mapsto \left(\sum_{i=1}^{r-1} \pm a_1 \otimes \cdots \otimes a_i a_{i+1} \otimes \cdots \otimes a_r \right) \pm (a_r a_1 \otimes \cdots \otimes a_{r-1}).$$

We obtain from this an A_∞ -structure on $H_*(A)$. Moreover, $H_*(A)$ is quasi-isomorphic to A as an A_∞ -algebra (the point is that the A_∞ -structure on A induced by the perturbed coderivation on $T^c(A)$ yields the original dg-algebra structure on A). That is, we recover the usual statement that a dga over a field can be recovered from its homology equipped with certain A_∞ -operations.

In summary: a perturbation ∂ of the coderivation d on $T^c(A)$ *such that $d + \partial$ is also a coderivation* determines an A_∞ -algebra structure on $H_*(A)$. In particular, a perturbation ∂ of the differential on A *such that $d + \partial$ is also a derivation* determines such a structure.

7.3. The Tate differential and A_∞ operations. We wish to apply the above formalism to the Tate resolution. Let X be a toric variety, and let \mathcal{K} denote the exact differential module $\bigoplus_{\ell \in \text{Pic}(X)} \mathcal{O}(-\ell) \otimes_k \omega(\ell, 0)$ with differential $\sum_{i=0}^n x_i \otimes e_i$. For the purpose of making it more

transparent that the above dg-methods extend to our setting, expand \mathcal{K} into a 1-periodic complex:

$$(5) \quad \cdots \rightarrow \mathcal{K}(0, 1) \rightarrow \mathcal{K} \rightarrow \mathcal{K}(0, -1) \rightarrow \cdots$$

Form a bicomplex \mathcal{B} whose columns are the Čech complexes of the terms of (5), with trivial horizontal differentials. Notice that the columns of \mathcal{B} split E -linearly. The complex $A = \text{Tot}^\oplus(\mathcal{B})$ therefore contracts onto its homology. Note also that the Čech cup product makes A a dga over the ring E . ♣♣♣ Michael: [Or maybe ω and not E ...of course the difference just amounts to a grading twist. A possibly more significant issue is that A is not a connected dga, because it's 1-periodic.]

We can perturb the differential on A by adding in the horizontal differentials. The result remains a dga. To see this, recall the general fact that, given two complexes \mathcal{F} and \mathcal{G} of sheaves, there is a map

$$\mathcal{C}(\mathcal{F}) \otimes \mathcal{C}(\mathcal{G}) \rightarrow \mathcal{C}(\mathcal{F} \otimes \mathcal{G})$$

of Čech complexes ♣♣♣ Michael: [tensoring differential modules is subtle, so this may require some more care]. Now note that we have a natural map $\mathcal{C}(\mathcal{K} \otimes \mathcal{K}) \rightarrow \mathcal{C}(\mathcal{K})$, since \mathcal{K} is a dga.

Applying the formalism in the previous section, we obtain an induced A_∞ -structure on the homology of A . Note that, of course, $H_*(A)$ is just the sheaf cohomology of the twists of \mathcal{O} in each degree.

Finally, recall that the perturbed differential on $H_*(A)$ is the Tate differential. On the other hand, the A_∞ -operations we constructed are built out of the corresponding perturbation of the differential on $T^c(H_*(A))$. So, these A_∞ -operations should be encoding the Tate differential. There is just a little fuzziness here concerning the relationship between the perturbed differential on $H_*(A)$ and the perturbed differential on $T^c(H_*(A))$. Modulo this, it's clear how the above A_∞ -operations are induced by the Tate differential.

To perform this construction with a sheaf other than \mathcal{O} , we need a version of the results in the previous section for dg- and A_∞ -modules, rather than algebras. I suspect this is possible, but I can't seem to find it written down (perhaps because it's too obvious to write down...).

REFERENCES

- [ABI07] L. L. Avramov, R.-O. Buchweitz, and S. Iyengar, *Class and rank of differential modules*, *Inventiones mathematicae* **169** (2007), no. 1, 1–35.
- [BGG78] I. N. Bernstein, I. M. Gel'fand, and S. I. Gel'fand, *Algebraic bundles over \mathbb{P}^n and problems of linear algebra*, *Funct Anal. and Appl* **12** (1978), 212–214.
- [EES15] D. Eisenbud, D. Erman, and F.-O. Schreyer, *Tate resolutions for products of projective spaces*, *Acta Mathematica Vietnamica* **40** (2015), no. 1, 5–36.
- [EFS03] D. Eisenbud, G. Floystad, and F.-O. Schreyer, *Sheaf cohomology and free resolutions over exterior algebras*, *Transactions of the American Mathematical Society* **355** (2003), no. 11, 4397–4426.
- [HK91] J. Huebschmann and T. Kadeishvili, *Small models for chain algebras*, *Mathematische Zeitschrift* **207** (1991), no. 1, 245–280.
- [Rou06] R. Rouquier, *Representation dimension of exterior algebras*, *Inventiones mathematicae* **165** (2006), no. 2, 357–367.