Hemodynamic Disorders
SGL 7
Embolism & infarction

Ass. Prof. Dr. Tenya Tariq Abdulhameed

M.B.Ch.B, M.Sc, Ph.D (HISTOPATHOLOGY)

College of Medicine, HMU.

2026-2027

OUTLINE:

Embolism

- Definition of an embolus
- Types of emboli

I.Pulmonary Thromboembolism

II.Systemic Thromboembolism


Infarction

- Introduction
- Causes of Infarction
- Morphology
- Factors That Influence Infarct Development


Summary

Embolism:

An **embolus** is a detached intravascular solid, liquid, or gaseous mass that is carried by the blood from its point of origin to a distant site, **where it often causes tissue dysfunction or infarction.**



Types of emboli:



1. The vast majority of emboli derive from a dislodged thrombus—hence the term thromboembolism.

2. **Fat embolism** can occur after crushing injuries to the bones, **(eg, due to a long bone fracture)** release microscopic fat globules into the circulation




3. **Amniotic fluid embolism** rarely may follow childbirth.
4. **Air embolism** can occur upon rapid decompression, most commonly in divers
5. **Less commonly,** emboli are composed of bubbles of nitrogen, atherosclerotic debris (cholesterol emboli), tumor fragments and bits of bone marrow.

the clinical outcomes are best understood from the standpoint of whether emboli lodge in the **pulmonary or systemic circulations:**


I. Pulmonary Thromboembolism
II. Systemic Thromboembolism

1) Pulmonary Thromboembolism


- Pulmonary emboli originate from deep venous thromboses and are responsible for the most common form of thromboembolic disease.
- The incidence of pulmonary embolism is 2 to 4 per 1000 hospitalized patients
- It causes about **100,000 deaths** per year in the United States.


Deep Vein Thrombosis : DVT

How a Pulmonary Embolism Forms







Embolus derived from a lower-extremity deep venous thrombus lodged in a pulmonary artery branch.

Clinical and pathologic features:

The consequences of embolization depend on:

1.the caliber of the occluded vessel

2.the collateral supply

3.and the affected tissue’s vulnerability to anoxia.

1. Most pulmonary emboli (60%–80%) are small and clinically silent. With time, they undergo organization and become incorporated into the vascular wall
2. At the other end of the spectrum, a **large embolus** that blocks a major pulmonary artery can cause sudden death, if lodge at the bifurcation of the right and left pulmonary arteries called **(saddle embolus)**



3. Embolic obstruction of **medium-sized arteries** and subsequent rupture of downstream capillaries rendered anoxic can cause pulmonary hemorrhage.
4. Embolism to **small end-arteriolar pulmonary branches** usually causes infarction.
5. **Multiple emboli** occurring through time can cause pulmonary hypertension and right ventricular failure (cor pulmonale).

2) Systemic Thromboembolism:

- Most systemic emboli (80%) arise from intracardiac mural thrombi.
- By contrast with venous emboli, which lodge primarily in the lung, arterial emboli can travel virtually **anywhere**


- Common arteriolar embolization sites include the lower extremities (75%) and central nervous system (10%); intestines, kidneys, and spleen are less common targets.
- arterial emboli often lodge in end arteries and cause infarction.

INFARCTION:




- An infarct is an area of ischemic necrosis caused by **occlusion of the vascular supply** to the affected tissue.
- Infarction primarily affecting the **heart and the brain** is a common and extremely important cause of clinical illness.

- Roughly **40%** of all deaths in the United States are a consequence of cardiovascular disease, with most of these deaths stemming from myocardial or cerebral infarction.



- **Pulmonary infarction** is a common clinical complication
- **bowel infarction** often is fatal





- ischemic necrosis of distal extremities **(gangrene)** causes substantial morbidity in the diabetic population.



Causes of Infarction:




- Arterial thrombosis or arterial embolism underlies the vast majority of infarctions.

Can venous thrombosis cause infarction?

Morphology:

- Infarcts are classified based on their **color** (reflecting the amount of hemorrhage)
- Thus, infarcts may be either red (hemorrhagic) **or white (anemic)**



A



B

A/Red infarction in brain,
B/ white infarction in spleen



Causes of red infarcts:

1. as a result of **venous occlusions** (such as in ovarian torsion)
2. in **loose tissues** (e.g., lung) where blood can collect in infarcted zones
3. in tissues with **dual circulations** such as lung and small intestine
4. **in previously congested tissues**
5. when flow is reestablished after infarction has occurred (e.g., after angioplasty of an arterial obstruction).


Cause of white infarcts:

White infarcts occur with arterial occlusions in **solid organs with end-arterial circulations** (e.g., heart,spleen, and kidney)



white infarction in kidney

Grossly: Infarcts tend to be wedge shaped, with the occluded vessel at the apex and the organ periphery forming the base




Microscopically: In most tissues, the main histologic finding associated with infarcts is ischemic coagulative necrosis:

I. An **inflammatory response** begins to develop along the margins of infarcts within a few hours and usually is well defined within 1 to 2 days.

II. Eventually, inflammation is followed by **repair,** beginning in the preserved margins


III. Most infarcts are ultimately replaced by **scar.**



Stages of ischemic necrosis in myocardium

The brain is an exception to these generalizations; ischemic tissue injury in the central nervous system results in **liquefactive necrosis**






Factors That Influence Infarct Development:


- The range of outcomes is influenced by the following three variables:

1. Anatomy of the vascular supply
2. Rate of occlusion
3. Tissue vulnerability to hypoxia

1. Anatomy of the vascular supply:


- The presence or absence of an alternative blood supply is the most important factor.
- The dual supply of the lung, **liver, hand and forearm**, with are rather resistant to infarction.
- By contrast, **the kidney and the spleen** both have end-arterial circulations, and arterial obstruction generally leads to infarction in these tissues.


Hepatic Vein
Oxygen-poor blood
Hepatic Artery
Oxygen-rich blood
Bile Duct
Hepatic Portal Vein
Nutrient-rich blood
FROM BODY
TO BODY
Gastrointestinal Tract


Arcuate artery
Interlobar artery
Descending aorta
Segmental artery
Renal artery

2. **Rate of occlusion**:

Slowly developing occlusions are **less likely** to cause infarction because they allow time for the development of collateral blood supplies.



3. **Tissue vulnerability to hypoxia:**

Neurons undergo irreversible damage when deprived of their blood supply for only **3 to 4 minutes.**

Myocardial cells, although hardier than neurons, still die after **only 20 to 30 minutes of ischemia**.

Summary:

- An embolus is a solid, liquid, or gaseous mass carried by the blood to a site distant from its origin; most are dislodged thrombi.
- Pulmonary emboli derive primarily from lower-extremity deep vein thrombi. Systemic emboli derive primarily from cardiac mural or valvular thrombi, aortic aneurysms, or atherosclerotic plaques
- Infarcts are areas of ischemic necrosis most commonly caused by arterial occlusion
- Infarcts caused by venous occlusion or occurring in spongy tissues typically are hemorrhagic (red);those caused by arterial occlusion in compact tissues typically are **pale (white).**