

Quiz 8

Math 54-Lec 3, Linear Algebra, Fall 2017

SECTION:

NAME:

You have 30 minutes to complete this quiz. To receive full credit, justify your answers.

Problem 1. (5 points) Let $A = \begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix}$. Find the singular values of A . Hence, what is the maximum value of the quadratic form $Q(\vec{x}) = \vec{x}^T A^T A \vec{x}$ subject to the constraint that \vec{x} is a unit vector?

$$A^T A = \begin{bmatrix} 2 & 0 \\ 3 & 2 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix} = \begin{bmatrix} 4 & 6 \\ 6 & 13 \end{bmatrix}$$

$$\det(A^T A - \lambda I) = \det \begin{pmatrix} 4-\lambda & 6 \\ 6 & 13-\lambda \end{pmatrix} = \lambda^2 - 17\lambda + 16 = (\lambda - 16)(\lambda - 1)$$

$\therefore A^T A$ has e-values 16, 1 \Rightarrow singular values of A are 4, 1.

Recall $\max \|A\vec{x}\| = \sigma_1$, the largest singular value of A .

$$\therefore \max \|A\vec{x}\| = 4$$

$$\therefore \max \|A\vec{x}\|^2 = (A\vec{x})^T (A\vec{x}) = \vec{x}^T A^T A \vec{x} = 4^2 = \boxed{16}$$

Note: all max are under constraint $\|\vec{x}\| = 1$.

Problem 2. (6 points) Consider the quadratic form: $Q(\vec{x}) = 2x_1^2 + 6x_1x_2 - 6x_2^2$. Find the matrix form of Q . That is, write $Q(\vec{x}) = \vec{x}^T A \vec{x}$ for some symmetric matrix A . Is Q positive definite, negative definite, or indefinite?

$$Q(\vec{x}) = \vec{x}^T A \vec{x} \quad \text{for } A = \begin{bmatrix} 2 & 3 \\ 3 & -6 \end{bmatrix}$$

$$\text{Observe, } Q\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}\right) = 2 \quad \& \quad Q\left(\begin{bmatrix} 0 \\ 1 \end{bmatrix}\right) = -6,$$

so Q is indefinite b/c it attains positive and negative values.

Quiz 8

Math 54-Lec 3, Linear Algebra, Fall 2017

Problem 3. (2 points each) Label the following statements true or false. If the statement is true, explain why. If it is false, explain why or provide a counterexample. Correct answers without justification will receive no credit.

(a.) Let A be an $n \times n$ orthogonally diagonalizable matrix. If A is invertible, then $A^{-1} = A^T$.

(b.) The expression $\|x\|^2$ is quadratic form.

(a) a matrix, A , is orthogonally diagonalizable iff it is symmetric. \therefore It suffices to find an invertible, symmetric matrix s.t. $A^{-1} \neq A^T$. Consider:

$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$, it is clearly symmetric. It is also invertible b/c its columns are linearly independent.

$$\text{However } A^{-1} = \frac{1}{\det(A)} \begin{bmatrix} 1 & -2 \\ -2 & 1 \end{bmatrix} = \frac{-1}{3} \begin{bmatrix} 1 & -2 \\ -2 & 1 \end{bmatrix} = \begin{bmatrix} -1/3 & 2/3 \\ 2/3 & -1/3 \end{bmatrix}$$

so $A^{-1} \neq A^T$. Hence, the statement is false.

(b) True, $\|\vec{x}\|^2 = \vec{x}^T \vec{x} = \vec{x}^T I \vec{x}$ where I is the $n \times n$ identity matrix.