

MATH 392 Problem Set 2

EJ Arce

2 February 2018

3.4

a


```
FlightDelays <- FlightDelays %>%
  mutate(Delayed20 = ifelse(Delay > 20, 1, 0))

# Calculating observed difference in the two groups' mean proportions
xobs <- mean(FlightDelays$Delayed20[FlightDelays$Carrier == "UA"] -
  mean(FlightDelays$Delayed20[FlightDelays$Carrier=="AA"]))

# Running simulation for hypothesis testing
nsim <- 10000
means <- rep(NA, nsim)
for(i in 1:nsim){
  perm <- sample(FlightDelays$Carrier, replace=F)
  means[i] <- mean(FlightDelays$Delayed20[perm=="UA"] -
    mean(FlightDelays$Delayed20[perm=="AA"]))
}
simdf <- data.frame(means)

# Plotting the simulated null distribution
ggplot(simdf, aes(x = means)) +
  geom_density() +
  geom_vline(xintercept = xobs, col = "red") +
  ggtitle("Density plot of observed mean differences from 10000 simulations")
```

Density plot of observed mean differences from 10000 simulations



The red vertical line indicates the difference in proportions observed in the actual dataset. The p-value for a two-tailed test is calculated below.

```
(sum(means > xobs) + 1)/(length(means) + 1) * 2
```

```
## [1] 0.0009999
```

b

We are testing to see if the variance in flight delays for United Airlines is greater than the variance for American Airlines. Thus, we are conducting a one-tailed significance test. Specifically,

$$H_0 : \rho_{UA} \leq \rho_{AA}$$

$$H_A : \rho_{UA} > \rho_{AA}$$

```
# Variance in flight delay lengths for each carrier
varUA <- var(FlightDelays$Delay[FlightDelays$Carrier == "UA"])
varAA <- var(FlightDelays$Delay[FlightDelays$Carrier == "AA"])
varUA
```

```
## [1] 2038
```

```
varAA
```

```
## [1] 1606
```


varUA and varAA indicate the variances of United Airlines' and American Airlines' delay times, respectively. A simulation just like the last problem will be used to test if the difference in these variances is statistically significant.

```
obs.diff <- varUA-varAA

nsim <- 10000
diffs <- rep(NA, nsim)
for(i in 1:nsim){
  perm <- sample(FlightDelays$Carrier, replace=F)
  diffs[i] <- var(FlightDelays$Delay[perm=="UA"] -
                 var(FlightDelays$Delay[perm=="AA"]))
}
simdf <- data.frame(diffs)

# Plotting the simulated null distribution
ggplot(simdf, aes(x = diffs)) +
  geom_density() +
  geom_vline(xintercept = obs.diff, col = "red") +
  ggtitle("Density plot of observed variance differences from 10000 simulations")
```

Density plot of observed variance differences from 10000 simulative



```
(sum(diffs > obs.diff) + 1)/(length(diffs) + 1)
```

```
## [1] 1
```

3.16

a

```
table(GSS2002$Gender, GSS2002$Pres00)
```

```
##
##      Bush Didnt vote Gore Nader Other
## Female  459          5  492    26    3
## Male    426          5  289    31   13
```

b

```
gender <- GSS2002$Gender
pres <- GSS2002$Pres00
chisq.test(gender, pres)
```

```
##
## Pearson's Chi-squared test
##
## data:  gender and pres
## X-squared = 33, df = 4, p-value = 1e-06
```

c

```
# Remove NAs
GSS2002 <- GSS2002 %>%
  filter(!is.na(Gender),
         !is.na(Pres00))
```

```
# Chi-squared test using permutations
x2.obs <- chisq.test(gender,pres)$statistic
nsim <- 10000
x2.stats <- rep(NA,nsim)
for(i in 1:nsim){
  perm <- xtabs(~sample(Gender, replace=F) + Pres00, data = GSS2002)
  x2.stats[i] <- chisq.test(perm)$statistic
}
(sum(x2.stats > x2.obs) + 1)/(nsim+1)
```

```
## [1] 9.999e-05
```

None of the simulated χ^2 values were greater than our observed χ^2_{obs} value of 33.29, resulting in our very low p-value.

3.22

```
q <- c(.2,.4,.6,.8)
d <- data.frame("quantile" = q)
obs.stats <- c(12.57,16.87,20.73,24.66)
d<-cbind(d,obs.stats)
norm.stats <- rep(NA,4)
for(i in 1:4){
  norm.stats[i] <- qnorm(q[i],22,7)
}
d <- cbind(d,norm.stats)
d
```

```
##   quantile obs.stats norm.stats
## 1      0.2      12.57      16.11
## 2      0.4      16.87      20.23
## 3      0.6      20.73      23.77
## 4      0.8      24.66      27.89
```

```
# Calculate observed chi-squared value
((3.54^2)/16.11) + ((3.36^2)/20.23) + ((3.04^2)/23.77) + ((3.23^2)/27.89)
```


```
## [1] 2.099
```

3.31

Empirical Solution

```
nsim <- 10000
t.obs <- rnorm(nsim,0,1)
ts <- data.frame(t.obs)
ts <- ts %>%
  arrange(t.obs) %>%
  mutate(t.obs = abs(t.obs))
p.val <- rep(NA,nsim)
for(i in 1:nsim){
  p.val[i] <- (sum(ts$t.obs>ts$t.obs[i])+1)/(length(ts$t.obs)+1)
}
```

```
ts <- cbind(ts, p.val)
ggplot(ts, aes(x=p.val)) +
  geom_density()
```



As we'd expect, the simulated density plot follows a uniform distribution.

Analytical Solution

Consider a test statistic $t = T(x_1, \dots, x_n)$. Its corresponding p-value is calculated by solving

$$p = \Pr(T(X) \geq t | H_0).$$

This makes p a random variable as well, since its calculated probability depends on the random variable $T(X)$. Thus the p-value follows some probability distribution $P(T)$. Since the p-values are drawn from the distribution of $T(X)$, then the p-values have a one-to-one correspondence to each observed test statistic t . Thus,

$$\Pr(P(T) \geq p) = \Pr(T(X) \geq t) = p.$$

This shows that $P(T)$ follows a uniform distribution, where $\Pr(p) = 1/n$.

3.32

Let Z denote the standard normal random variable. Then $Z \sim N(0,1)$. Suppose $X = Z^2$. Show that $X \sim \chi^2_{df=1}$.

The pdf of Z is already known to be

$$f_Z(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}$$

$$F_Z(z) = \frac{1}{2} \left(1 + \operatorname{erf}\left(\frac{z}{\sqrt{2}}\right) \right)$$

Using the cdf method,

$$F_X(x) = P(X \leq x) = P(Z^2 \leq x) = P(Z \leq \sqrt{x}) = F_Z(\sqrt{x})$$

$$F_Z(\sqrt{x}) = \int_{-\sqrt{x}}^{\sqrt{x}} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx$$

$$F_Z(\sqrt{x}) = 2 \int_0^{\sqrt{x}} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx$$

$$F_Z(\sqrt{x}) = 2 \frac{1}{2} \operatorname{erf}\left(\frac{\sqrt{x}}{\sqrt{2}}\right)$$

$$F_Z(\sqrt{x}) = \operatorname{erf}\left(\frac{\sqrt{x}}{\sqrt{2}}\right)$$

$$f_X(x) = \frac{\partial}{\partial x} F_Z(\sqrt{x}) = f_Z(\sqrt{x}) \frac{1}{2} x^{-\frac{1}{2}}$$

$$f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{x}{2}} \frac{1}{2} x^{-\frac{1}{2}}$$

$$f_X(x) = \frac{x^{-\frac{1}{2}} e^{-\frac{x}{2}}}{2\sqrt{2\pi}}$$

Notice that

$$\Gamma(1/2) = \int_0^\infty t^{-1/2} e^{-t} dt = \sqrt{2\pi},$$

so we get

$$f_X(x) = \frac{x^{-\frac{1}{2}} e^{-\frac{x}{2}}}{2\Gamma(1/2)}$$

Thus,

$$f_X(x) \sim \chi_{df=1}^2$$