

MATH 392 Problem Set 5 (Corrected)

EJ Arce

9 March 2018

6.39

$\{\hat{\beta}_n\}$ is consistent if and only if $\lim_{n \rightarrow \infty} P(|\hat{\beta}_n - \beta| < \varepsilon) = 1 \forall \varepsilon > 0$.

We know that for any n , the distribution of X_{max} follows the distribution

$$F_{X_{max}}(x) = \begin{cases} 0 & \text{if } x > \beta \\ (\frac{x}{\beta})^n & \text{if } 0 < x < \beta \\ 0 & \text{if } x < 0 \end{cases}$$

Thus

$$\begin{aligned} \lim_{n \rightarrow \infty} P(|\hat{\beta}_n - \beta| < \varepsilon) &= \lim_{n \rightarrow \infty} P(\beta - \varepsilon < \hat{\beta}_n < \beta + \varepsilon) \\ &= \lim_{n \rightarrow \infty} P(\beta - \varepsilon < X_{max} < \beta) \\ &= \lim_{n \rightarrow \infty} \left(\frac{\beta}{\beta}\right)^n \Big|_{\beta-\varepsilon}^{\beta} \\ &= 1 - \lim_{n \rightarrow \infty} \left(\frac{\beta - \varepsilon}{\beta}\right)^n \\ &= 1 \end{aligned}$$

6.40

$\{\hat{\sigma}_n^2\}$ is consistent if and only if $\lim_{n \rightarrow \infty} P(|\hat{\sigma}_n^2 - \sigma^2| < \varepsilon) = 1 \forall \varepsilon > 0$.

From Problem Set 4,

$$E[\hat{\sigma}_n^2] - \sigma^2 = \frac{n-1}{n}\sigma^2 - \sigma^2 = \frac{-1}{n}\sigma^2$$

It suffices to show that $\lim_{n \rightarrow \infty} \left|\frac{n-1}{n} - 1\right| < \varepsilon \forall \varepsilon > 0$.

Let $\varepsilon > 0$. Then $\exists m \in \mathbb{N}$ st $\forall n \geq m, \left|\frac{n-1}{n} - 1\right| < \varepsilon$. Thus, $\forall n \geq m$,

$$\left|\frac{1}{n}\right| < \varepsilon \Rightarrow \left|\frac{1}{n}\right| < \varepsilon \Rightarrow \left|\left(\frac{n-1}{n} - \frac{n}{n}\right)\right| < \varepsilon \Rightarrow \left|\frac{n-1}{n} - 1\right| < \varepsilon \Rightarrow |E[\hat{\sigma}_n^2] - \sigma^2| < \varepsilon.$$

Thus, $\lim_{n \rightarrow \infty} P(|\hat{\sigma}_n^2 - \sigma^2| < \varepsilon) = 1 \forall \varepsilon > 0$.

7.3

a

$$X \sim N(185, 50^2), n = 100, \bar{X} = 210$$

$$P(q_{.05} < \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} < q_{.95}) = 1 - \alpha = .9$$

Solve for μ :

$$P(\bar{X} - q_{.95} \frac{\sigma}{\sqrt{n}} < \mu < \bar{X} - q_{.05} \frac{\sigma}{\sqrt{n}}) = .9$$

$$(210 - 1.645 \frac{50}{\sqrt{100}} < \mu < 210 + 1.645 \frac{50}{\sqrt{100}}) = (201.8 < \mu < 218.8)$$

Thus the 90% confidence interval is (201.8, 218.8).

b

Solve for n:

$$\bar{X} - LB_{.05} = \bar{X} - (\bar{X} - 1.96(\frac{50}{\sqrt{n}})) \leq 10$$

$$1.96 \frac{50}{10} \leq \sqrt{n}$$

$$n \geq 96.04$$

A sample size of at least 97 is needed.

c

Like b,

$$2.576 \frac{50}{10} \leq \sqrt{n}$$

$$n \geq 165.9$$

A sample size of at least 166 is needed.

7.8

For $n = 30$, the code generating the simulation is below, along with the proportion of times the confidence interval missed μ . Compared to the next three numbers for $n=60$, $n=100$, and $n=250$, the frequency of missing μ converges to the theoretical frequency of .05.

```
# Run simulation
low <- 0
high <- 0
n <- 30
q <- qt(.975, n-1)
N <- 10^5
```

```

for(i in 1:N){
  x <- rgamma(n,shape = 5, rate = 2)
  xbar <- mean(x)
  s <- sd(x)
  L <- xbar - q*s/sqrt(n)
  U <- xbar + q*s/sqrt(n)
  if(U<5/2){
    low <- low + 1
  }
  if(5/2<L){
    high <- high + 1
  }
}

(high+low)/N

```

```

## [1] 0.05396
## [1] 0.05194
## [1] 0.05169
## [1] 0.05009

```

7.34

We are given

$$X \sim \text{Gamma}(2, \lambda), f_X(x) = \frac{x e^{-x/\lambda}}{\Gamma(2)\lambda^2}$$

$$2\lambda x \sim \chi_{df=4}^2, f_{2\lambda X}(2\lambda x) = \frac{(2\lambda x)^{4/2-1} e^{-(2\lambda x)^2/2}}{2^{4/2}\Gamma(4/2)}$$

Thus, the quantiles will be calculated in terms of $2\lambda x$, using the $\chi_{df=4}^2$ distribution.

$$\begin{aligned}
 .95 &= P(q_{1,x} < \lambda < q_{2,x}) = P(q_{1,2\lambda x} < 2\lambda X < q_{2,2\lambda x}) \\
 &= P\left(\frac{q_{1,2\lambda x}}{2X} < \lambda < \frac{q_{2,2\lambda x}}{2X}\right) \\
 &= P\left(\frac{.2422}{X} < \lambda < \frac{5.572}{X}\right)
 \end{aligned}$$

Thus the 95% confidence interval for λ is $(\frac{.2422}{X}, \frac{5.572}{X})$.