

MATH 392 Power Calculations

EJ Arce

9 March 2018

Given $\mu = 100, \sigma^2 = 15^2, \bar{X} = 105, \alpha = .05$, for what sizes n will power be .7? .75? .9?

$$H_0 : \mu = 100$$

$$H_A : \mu > 100$$

Power = .7

Under each hypothesis, the sampling distributions will follow

$$X_{H_0} \sim N(\mu, \sigma/\sqrt{n})$$

$$X_{H_A} \sim N(\bar{X}_{obs}, \sigma/\sqrt{n})$$

With $\alpha = .05$, the critical value x_{crit} at which we reject the null hypothesis is $\mu + 1.645(\sigma/\sqrt{n})$. β is the area under the alternative hypothesis distribution to the left of this critical value. With $\beta = .3$, we can refer to the standard normal distribution to find how many standard errors away from the mean our critical value is:

```
qnorm(.3,0,1)
```

```
## [1] -0.5244005
```

Thus,

$$x_{crit} = 100 + 1.645 \frac{\sigma}{\sqrt{n}} = 105 - .5244 \frac{\sigma}{\sqrt{n}}$$

Now solve for n :

$$\begin{aligned} (1.645 + .5244) \frac{\sigma}{\sqrt{n}} &= 5 \\ \left(\frac{(2.189)15}{5} \right)^2 &= n \\ n &= 42.35 \end{aligned}$$

Thus for a power of at least .7, the sample size must be at least 43.

Power = .75

Generalizing the equations for x_{crit} ,

$$100 + 1.645 \frac{\sigma}{\sqrt{n}} = 105 - q_{z_\beta} \frac{\sigma}{\sqrt{n}}$$

where q_{z_β} is the quantile for the standard normal distribution under which the area to the left of the quantile is β . Thus for $\beta = .25$, q_{z_β} is

```
qnorm(.25,0,1)
```

```
## [1] -0.6744898
```

Thus

$$\begin{aligned} (1.645 + .6745) \frac{\sigma}{\sqrt{n}} &= 5 \\ \left(\frac{(2.319)15}{5} \right)^2 &= n \\ 48.4 &= n \end{aligned}$$

A sample size of at least 49 is needed for a power of .75

Power = .9

```
qnorm(.1,0,1)
```

```
## [1] -1.281552
```

$$\begin{aligned} (1.645 + 1.282) \frac{\sigma}{\sqrt{n}} &= 5 \\ \left(\frac{(2.927)15}{5} \right)^2 &= n \\ 77.11 &= n \end{aligned}$$

A sample size of at least 78 is needed for a power of .9.