POSITION PAPER · PREPRINT · NOT PEER REVIEWED

The perforator hypothesis: knots as small arteries held shut where they pierce the fascia

Eugene Jhong · Knots of Existence, independent · 2 November 2026 · draft 1

Written by AI (Claude Opus 5.5, Anthropic) under the author's direction, and refereed by a second model (Claude Fable 5.1). See Authorship and AI use.

Correspondence and objections: github.com/ejhong/knots/issues · PDF · the companion preprint

ABSTRACT

Anyone with a foam roller can find them: small, tender places in the soft tissue that persist for years yet can let go within seconds, sometimes with a tingle across a patch of skin. No mechanism for them is established, and no release has been recorded as it happens. We propose one. A knot is a cutaneous perforator, a small artery travelling with its veins and a nerve through an opening in the fascia, held shut by the smooth muscle of its own wall. Each part rests on measured physiology. A toned small artery can snap shut and stay shut until its tone falls well below the level that shut it; held shut for hours, it relaxes only slowly and incompletely; the breath moves the skin's sympathetic drive within seconds; a squeeze and release widen a small artery; a nerve deprived of blood fires when blood returns; and a vessel held shut holds shut the branches it feeds. A knot, then, is a memory with an anatomy: written by the sympathetic drive, kept by the wall, read by the perforator's own nerve. We argue that it is now the strongest account in a specific sense: it explains more of what is reported, with less that is new, than six alternatives in their strongest forms. We state what it does not explain, weigh the claim that acupuncture points sit on perforators, consider the slower fascial changes reported after intensive practice, and their dangers, and give the measurements that would refute it.

A caution. Experiences like some of those discussed here have for some people been prolonged and disabling (Section 9). This paper describes; it does not prescribe. Nothing in it is a practice or a protocol, and no one should try to bring these experiences on. Anyone in difficulty can find support from Cheetah House.

1. Introduction

1.1 What is reported

Anyone with a foam roller can find them: small, discrete, tender places in the soft tissue, dozens of them, by many accounts densest around the hips and shoulders, and more with every decade. Massage therapists and physiotherapists call them knots or trigger points; contemplative traditions speak of knots in the body's channels. Pressed, a knot announces itself unmistakably. Stretched, it gives not the sharp pain of a muscle at the end of its range but a dull block, as though layers had stopped sliding past each other.

Those who work on knots, the author among them, describe a consistent set of features (Box 1). These are impressions gathered informally, not measurements. Some are echoed in systematic work: in a study of meditation-related challenges among Western Buddhist practitioners, pressure and tension that grew more acute or released in the course of practice were one principal set of the somatic changes reported [1]. We take Box 1 as what any account of knots must explain, and no more.

Box 1. What practitioners report

- They persist. A fresh knot may yield in a session; an old one can take months.
- They let go within seconds, not days: under a hand, often as it lifts; under a roller; with the breath, by many accounts less with a single deep breath than with a subtle, complex pattern of breathing joined to fine attention to the body.
- Deeper ones let go to focused attention, sometimes without touch.
- Warmth helps: a hot shower softens many at once.
- A release can come with a small pop, or a tingle across a patch of skin.
- They limit movement: a stretch meets a dull block at the knot.
- They come in families. When one lets go, others nearby may go with it, or another may appear close by; a large one can free smaller ones around it, not the reverse.
- They gather where stress is held, and accumulate with age.
- Some come back to roughly the same place.

1.2 A state, not a structure

A knot that lets go within a breath cannot be a scar or an adhesion: what the body has built, it cannot unbuild in a breath. It must be a state the body is holding, as the first essay on these knots put it [2]: the tissue stays where it is, and the knot is a state written on it. The question is which state, held where, and by what. Several candidates have been proposed (Section 4). We argue for one, and claim for it something specific. At

present it is the strongest account in three senses: it explains more of Box 1 than any alternative; it does so with physiology that is already measured, adding a new identification rather than a new mechanism; and it is the cheapest to refute.

1.3 Where the hypothesis came from

The hypothesis grew in public. The first essay [2] followed Johnson's vasocomputation [3], in which a held vascular contraction stores a state, and placed the store in a latch in the smooth muscle of small arteries inside muscle. The second [4] moved it to the perforators, the bundles that carry an artery, its veins and a nerve through the fascia to the skin, and supposed that a collar of gelled hyaluronan around each held it; a longer preprint of the same month set out that version at length [5]. Checked against physics, the collar failed. At its most viscous, hyaluronan around a vessel pushes back about 12 times more weakly than the blood pressure holding the vessel open [6, 7, 8], and on its own, in physiological conditions, it stays a liquid rather than gelling [9]. If anything holds a perforator shut, it is the wall. That is the form argued here; where the earlier versions differ, this one replaces them. A companion preprint runs seven candidate mechanisms evenly, as models [10]; this paper argues for one.

2. The hypothesis

> *A knot is a cutaneous perforator, a small artery travelling with its veins, a nerve and a lymphatic through an opening in a fascia, held shut by the smooth muscle of its own wall. It lets go when the wall's tone falls, or the wall is moved, enough for the vessel to reopen; blood returning to the patch the vessel feeds wakes its nerve.*




Figure 1. One perforator in section. (a) Open: a small artery and its nerve rise through an opening in the deep fascia, cross the gliding layer in a sleeve of sliding tissue, and branch in the skin. (b) Held shut by its own wall at the opening: a knot. Its sleeve dries and can jam; the patch it feeds is underfed. (c) Letting go: tone falls below the band that held it, blood floods the patch, and the nerve fires.

In parts, each of which can be checked on its own:

1. **Place.** Knots sit where perforators pass through the deep fascia (the larger ones) or the superficial fascia (the smaller ones), at fixed sites. · new
2. **Hold.** A small artery with tone in its wall can close and stay closed until its tone falls well below the level that closed it. · established
3. **Set.** Held shut for hours, the wall's muscle adapts to its shortened length and relaxes only slowly and incompletely; over days the artery remodels inward. The vessel could then stay shut nearer to resting tone. · established in small arteries; in knots, new
4. **Drive.** What carries a vessel shut is the sympathetic drive to the skin's vessels, raised by stress and moved within seconds by the breath. What lets it go is that drive falling, warmth, or the wall being moved: shaken, or squeezed and released. · established physiology; its role in knots, new
5. **Nerve.** While the vessel is shut, the patch it feeds is underfed and tender; when blood returns, the nerve that went without it fires: the tingle of a release. · the tingle established; the tenderness, new
6. **Sleeve.** Each vessel crosses the gliding layer between the fasciae in a sleeve of sliding tissue that its flow keeps wet; a vessel shut for long lets its sleeve stiffen, pinning the layers there: the dull block. · hypothesis
7. **Trees.** Perforators branch from parents. A shut parent holds its children shut, and when it opens they can open with it; a child that opens cannot open its parent. · hydraulics; in knots, new

A memory with an anatomy

Read as a whole, the hypothesis describes a memory with an anatomy. In Johnson's account a held vascular contraction acts as a medium-term memory [3], and the essay that brought the idea to the perforators asked what would write, store and read such a state [4]. Here the answer is concrete. The sympathetic drive writes: stress raises it, and a perforator it carries shut stays shut as the drive eases, so long as its tone does not fall below the band; only adaptation and remodelling let it hold at rest. The writing is local. Vessels differ in the thickness of their walls, the pressure that feeds them and the load at their openings, so a rising drive carries a few of them across the band at a time, not all at once. The wall keeps: its switch holds the state, and adaptation and remodelling make it durable. The perforator's own nerve reads: as tenderness while the vessel holds,

and as a tingle when it lets go. Each perforator is a site that can be set or cleared, and a body has hundreds of large ones and many more small ones [11]: a map of held states that a hand can find, the breath can reach and an instrument could read.

3. The case

No single observation below proves the hypothesis. The argument is that one anatomical object, behaving as measured physiology says it behaves, accounts for nearly every feature in Box 1, where each alternative accounts for some.

3.1 Where knots are

Perforators are discrete, numerous and fixed. Taylor and Palmer's whole-body injection studies plotted an average of 374 major perforators in each body, the arteries closely following the connective-tissue framework [11]. They pierce the deep fascia at sites that recur from person to person: the clusters are shared, but the individual vessels scatter around their cluster's centre, in the leg with standard deviations of 6–11% of the tibia's length, roughly 2–4 cm [12], and differ between the two sides of one body [13]. That is how knots are described: in the same regions in everyone, never at quite the same point.

They are also where the layers are held. The superficial fascia is tied to the deep fascia by fibrous bands, the retinacula cutis, which work as tethers and anchors [14], and perforators cross the same gap, large and sparse where the skin is mobile, smaller and denser where it is tethered [15]. Pressure from above presses each bundle against the firm deep fascia beneath it: a hand or a roller meets a perforator where it can squeeze it.

And perforator exits are already used to find tender points. In entrapment of the cutaneous nerves of the abdominal wall, the perforating vessels' exit through the rectus fascia beneath the point of maximal pain is found by Doppler and used as the landmark for injection [16]; lidocaine injected at the point of maximal pain relieved it in 13 of 24 patients, saline in 4 of 24 [17]; and a case report describes a tender point with a pulsating perforating artery directly beneath it on Doppler, after whose release with fluid the sign placing the pain in the abdominal wall disappeared and pain and movement improved [18]. These implicate the nerve as much as the vessel. They show that the address is right.

3.2 What holds them

A small artery is a balance between the pressure inside it, pushing it open, and the tension of its muscle, pulling it shut. Burton showed that with active tension in the wall the balance has no stable open state below a critical pressure: the vessel snaps shut [19]. Whole vascular beds stop flowing at such a critical closing pressure [20], and failing skin flaps show it [21]. Because a muscle is weak when it is very short [22], a vessel that has closed needs far less tone to stay closed than it took to close it. Over a band of tone, open and shut are both stable, and a shut vessel reopens only when its tone falls below the band, when outside pressure is relieved, or when the pressure feeding it rises [23]. In living tissue, nerves hold some small arterioles fully closed at rest: in the muscle of anaesthetised rats, cutting them opened 22% more [24].

With the measured strength and wall thickness of human small arteries [22, 25], the project's model of the wall finds that band between 0.62 and 0.93 of maximal tone, with rest at about 0.19 (Figure 2). Across 49,152 plausible parameter sets, about half (52%) give the switch, and only 6% hold shut at resting tone [8]. The hold is possible but demanding: a healthy vessel holds itself shut only while its tone stays well above rest. Thicker walls, as in hypertension [25], make it easier.




Figure 2. The switch in a small artery's wall, from the project's model with measured parameters: how far shut the vessel is (up, from relaxed to shut) against the tone of its muscle (across). Over the shaded band, open and shut are both stable. Tone rising past the band snaps the vessel shut; once shut, it reopens only when tone falls below the band.

3.3 Why old knots persist

Smooth muscle adapts to the length at which it is held. Arterioles constricted for five minutes relax fully when the drive is removed; after four hours they relax slowly [26] and incompletely, their cells repositioned to hold the smaller diameter [27], though the hold still needs calcium and goes at once without it [26]. Smooth muscle held short regains much of the force it lost there [28, 29], and resistance arteries kept constricted for three days remodel inward, their relaxed lumen 10–16% smaller [30]. A vessel held shut long enough might then hold nearer to rest: the knot would set, and the setting would deepen with time, while still answering to whatever lowers the wall's calcium. That fits the difference practitioners describe between fresh knots, which yield in a session, and old ones, which take months; and it predicts something an isolated vessel can show (Section 7).

3.4 What the breath can reach, and how fast

The skin's small arteries are governed by sympathetic nerves: skin blood flow runs from almost nothing under local cooling to several litres a minute in heat [31], and about a third of the superficial fascia's nerve fibres are autonomic, gathered around its vessels [32]. The breath reaches this drive within seconds. A deep inspiration constricts the skin's vessels [33], cutting fingertip flow by 59–71% [34], beginning 3.6–3.8 s after the breath, deepest at about 5 s, and recovering over the following ten to thirty seconds [35]. Slow breathing works the other way: fifteen minutes of it lowered sympathetic activity to muscle by five bursts in every hundred heartbeats [36]. A single deep breath, then, briefly raises the drive to the

skin's vessels, which may be why releases are reported less with one deep breath than with a subtler pattern (Box 1). These measurements come from the skin of the fingers and toes, unusually rich in sympathetic control, and from the nerves to muscle; whether the breath moves the drive to the perforators of the trunk as fast, or as far, is not known. Movement is a possible second route: vibration cuts the active force of vascular smooth muscle at once [37], but at 1 to 400 cycles a second, far faster than a breath, and whether the slow stretch of breathing does the same is untested. Which route matters, and which pattern of breathing serves it, is open; the hypothesis requires only that a releasing breath lower the drive more than it raises it, or move the wall.

3.5 Why hands, rollers and warmth

A vessel pressed shut from outside cannot reopen while the pressure lasts; it can reopen as the pressure lifts [23]. A knot under a hand should therefore let go as the hand lifts or eases, not while it presses. Squeezed and released, the feed arteries of rat muscle widen within seconds, and several brief squeezes widen them more than one long one [38]: a roller's passes are a series of brief squeezes, and it shears the tissue around the vessel as well. Heat acts more slowly and more widely. Local warming raises the skin's flow most of the way to its maximum [39], but a patch of skin warms with a time constant of about 44 s even at maximal flow [8]: too slow to free a knot within a breath, fast enough over the minutes of a hot shower, and everywhere the water touches.

3.6 Why a knot is tender, and why a release tingles

While a perforator is shut, the patch it feeds is underfed, and so is the nerve that travels with it. Underfed, not starved: a perforator's territory is linked to its neighbours' by anastomoses [40], through which some blood can still come. At active trigger points the tissue is acidic and rich in the substances that sensitise pain nerves [41]; an underfed patch of skin and fascia could build the same chemistry, and its nerve would be tender to pressure. On this account a knot is tender because it is shut. When flow returns after a five-minute arterial occlusion of the forearm, the skin's flow overshoots, rising by about 187% and peaking about 11 s after the release [42], a response in which sensory nerves and BKCa channels take part [43]. A nerve that has gone without blood also fires on its own when the blood returns: recorded in human sensory nerves, this firing begins about half a minute after circulation is restored and peaks by the second minute, as the felt tingling does [44]. When a shut perforator opens, blood floods its patch and the nerve that went without it fires: on a small scale, the prickle of a limb waking after pressure. The tingle should follow the opening within about a minute and cover the vessel's territory, a patch of skin rather than a line [40]. Two cautions. Overbreathing increases such firing [44], so a tingle during forceful breathing is weak evidence by itself; and a single perforator's flood is far smaller than a cuff's, which is why a recording needs a pressed sham beside it (Section 7).

3.7 Why a dull block

Each perforator crosses the gliding layer between the superficial and deep fasciae, a web of small fluid-filled cells that lets skin, fat, fascia and vessels slide past each other [45]. The vessel's flow keeps its stretch of this web wet. Shut for long, the sleeve would dry and stiffen. This is not the collar of Section 1.3, which had to hold a vessel shut against the blood pressure and could not. The sleeve need only resist the small shear of layers sliding past each other, and for that it needs a material that holds until pushed past a threshold. The gliding layer offers candidates: crowding and immobility raise hyaluronan's concentration and viscosity and reduce the gliding of the layers, and in inflammation modified hyaluronan can form aggregates or gel-like structures [46]; materials of this kind can jam below a critical stress and give way abruptly above it [47]. A jammed sleeve pins the layers at that point, and a stretch meets a block there. Fascial layers glide less in chronic low back pain [48], and the loose sublayers of the neck's fascia are thicker in chronic neck pain [49]. The sleeve is the least established part of the hypothesis, and it predicts an order a recording could check: the flood and the tingle as the vessel opens, the sliding only once the layer is moved.

3.8 Why releases come in families

Perforators form trees: a parent vessel feeds children around it. A child cannot flow while its parent is shut, so a shut parent holds its children shut, and when the parent opens, those not held by their own walls open with it. A child that opens does nothing for a parent that stays shut. The asymmetry is hydraulic: pressure comes from the source. Signals along the wall would not supply it, for a dilation started on an arteriole spreads both ways [50], fading with a length constant of about two millimetres, and those started on daughter vessels sum in the parent [51]; over the centimetres between one perforator and the next, they would have faded. Release should then run in families: freeing a large knot frees small ones around it, and freeing a small one does not free the large one. Trigger-point practice describes a similar order: dry needling a key trigger point raised the pressure-pain threshold of satellite points in its zone of referred pain [52], though a spinal mechanism has been proposed for it.

3.9 Why more with age

Small mesenteric arteries from organ donors over sixty-five contract more strongly to adrenergic stimulation [53], and resting sympathetic activity to muscle rises with age [54]; older skin reaches a lower maximal flow when heated [55]. All three would make vessels likelier to hold and harder to flood open. But the skin's reflex vasoconstriction is blunted with age [56], so whether the drive to the skin's own vessels rises is not settled. The hypothesis predicts that the number of knots tracks the loss of the skin's microvascular reactivity, place by place, and that children have few.




Figure 3. The clocks. How fast knots are reported to change (terracotta; rough, from Box 1), beside how fast the physiology the hypothesis names is measured to change (slate). A deep breath narrows the skin's vessels from 3.6 s, most at about 5 s, recovering over the following ten to thirty seconds [35]; a reopened vessel's flow peaks about 11 s after release [42]; a patch of skin warms with a time constant of 44 s at maximal flow and 341 s at a tenth of it [8]; arterioles constricted for five minutes keep no trace of it, and after four hours relax only slowly and incompletely [26, 27]; arteries constricted for three days remodel inward [30]. The slower changes in the fascia (Section 5) are reported over days to weeks. The shaded band is a breath.

Table 1. What the hypothesis accounts for, what it rests on, and how firmly.

Reported	The perforator account	Rests on	Status
Discrete, tender places in the same regions in everyone	Perforator exits: fixed sites, shared clusters; tender because shut	[11, 12, 16, 41]	Identity untested
They persist for months	A wall that holds itself shut; muscle that adapts; an artery that remodels	[19, 27, 30]	Each measured; together untested
They let go within seconds	A shut vessel reopens as soon as its tone falls below the band	[19, 20, 23]	Established mechanism
With the breath	A deep breath raises the drive for seconds; slow breathing lowers it	[35, 36]	Measured in fingers and in nerves to muscle; in knots, untested
As a hand lifts; under a roller	A pressed vessel reopens only on release; brief squeezes widen it	[38]	Measured in feed arteries of rat muscle
A hot shower frees many	Local warming lowers the tone of the skin's vessels	[39]	Established
A tingle over a patch	A nerve that went without blood fires when it returns; reactive hyperaemia	[43, 44]	Established
A dull block to stretch	A dry sleeve jams in the gliding layer	[45, 46, 47]	Hypothesis
Families; a large one frees small ones	Trees: a shut parent holds its children shut	[11]	Hydraulics; in knots, untested
More with age	Stronger adrenergic contraction; less capacity to dilate	[53, 56]	Mixed
Deeper ones go to attention alone	No direct route	—	Not explained

4. The alternatives

Six other mechanisms have been proposed or can be assembled from established physiology (Table 2). Each has a strongest form, and most have evidence the perforator hypothesis lacks.

Table 2. Six alternatives in their strongest forms, and the trace each would leave at the moment of release.

Mechanism	What holds a knot	Best evidence	Explains, and struggles with	At release
Trigger points	An energy crisis: sustained contraction squeezes its own capillaries, and the ischaemia withholds the energy needed to relax [57]	Stiff nodules on ultrasound [58]; an acidic milieu [41]; spontaneous electrical activity [59]; lactate rises after compression [60]	A palpable nodule, a twitch, referred pain. It struggles with release in seconds to the breath and with a tingle over the skin; finding points by touch is only moderately reliable [61], and the construct is contested [62]	Stiffness falls in the muscle
Densification	Hyaluronan between fascial layers grows viscous, and in inflammation can form gel-like aggregates; the layers stop gliding [46, 63]	Thicker loose sublayers in neck pain [49]; less glide in back pain [48]	The block and the slide, and perhaps the slower changes in the fascia (Section 5). It struggles with seconds: hyaluronan on its own stays liquid [9], and warmth thins it only two- to threefold over forty degrees [46]	The layers slide again
Nerves	A cutaneous nerve sensitised where it pierces the fascia [64]	Established entrapments [17]; fibres that fire when pressed after a neuritis [65]	Tenderness, and a local anaesthetic test. It has no hold of its own, and in one preparation, in nerve-injured rats, sympathetic stimulation changed sensory firing only after about fourteen seconds [66]	The nerve's firing falls
Vascular latch	A contraction held long enough engages smooth muscle's latch and freezes a prediction in place, cut off from awareness and from updating [3]	An integrative account; untested	Release to attention alone. The latch saves energy but does not by itself remember [67]	Flow rises slowly
Motor switch	Motor units, once recruited, keep firing on persistent inward currents [68, 69]	Self-sustained firing measured in people	Release to attention and inhibition, at once. It struggles with a tingle over a patch of skin, and with a release at a site with no muscle beneath it	A motor unit falls silent
Perception	Central gain, kept up by threat, attention and poor sleep, makes ordinary input felt as a knot [70]	Central sensitisation is established in chronic pain	Variability and global effects. It struggles with discrete places that return to the same spot	Nothing changes in the tissue
Perforators	A small artery's wall holds itself shut	Its parts are measured; the identity is not	Most of Box 1; not release to attention alone	Skin flow floods in

Three points about the comparison. First, the perforator and the nerve share an address. A cutaneous nerve passes through the same opening, so finding tender points on perforators would not separate them; low flow before a release, and a flood at it, would.

Second, the perforator hypothesis has no direct route for knots that let go to attention alone, without touch. The latch or the motor switch would; so would a perforator whose tone is held in its band by a held prediction, the latch's command acting on the perforator's wall.

Third, the alternatives need not all be rivals. Knots may be more than one thing, and densification, which struggles with release in seconds, may describe the slower changes of Section 5. A recording should classify each release rather than crown one theory.

The companion study writes all seven mechanisms as models from their own physiology and runs them through the same scenes [10]. We give its comparisons little weight here: most numbers specific to knots are guesses there, the formalisations of three of the seven are the project's own and have not been checked by those who hold them, and a shared calibration gives every model knots by construction. What the case rests on is the physiology above.

5. Beyond the knots: the fascia

The knots are not the end of what is reported. Some who take this work furthest, chiefly in long and intensive contemplative practice, describe a second family of changes that follows them: layers of the fascia beneath the skin felt to peel apart along lines, places left feeling hollow, and then,

over days, a filling-in felt as re-stitching. These reports are rarer than knots, come from fewer people and are harder to check. They run on a slower clock, hours to weeks rather than seconds, and in them the order runs one way: the knots first, the layers after. They belong to a wider record of bodily experience in intensive practice, which includes pressure and tension that build and release, currents of energy and involuntary movement [1], and energy-like somatic experiences whose course, effects and interpretation have been described in their own right [71].

The perforator hypothesis bears on this family because perforators are among the things that hold the fascia down. The superficial fascia runs unbroken over the body and glides over the deep fascia on a plane of loose tissue [45, 72]; beneath a firm, protective superficial layer lies a mobile one, made for gliding [73]. It is held in place by fibrous tethers [14] and by the perforators that pass through it, each in its sleeve. If a shut perforator's sleeve jams (Section 3.7), each knot is also a point where the layers are pinned. Were many to let go across a region, and the returning flood to wet the gliding plane, the layers could part along the lines between the remaining tethers, and close again as the fluid is taken up and the sleeves hold once more. This is a hypothesis built on a hypothesis, and it is not yet modelled.

There is a strong alternative. Intensive practice brings long hours of attention to the body, and often overbreathing, strain and short sleep, each of which can produce vivid bodily sensation with no change in the tissue; attention to the body is not always benign, and past some point its effects can turn negative [74]. A reading in which these experiences are changes in how the body is perceived, the perception account of Section 4, is the natural first reading for a clinician. Practitioners in difficulty found grounding, calming and embodying activities, sleep, diet and exercise helpful; body-based treatments, massage and acupuncture among them, were reported as helpful by some and not by others [1].

The two readings predict different things, and ultrasound can tell them apart. The glide between fascial layers can be measured [48]. If the layers are changing, glide should differ at the sites where the changes are reported, follow the knots rather than precede them, and settle over the days the reports give; if only perception is changing, the tissue should look as it did. On the first reading, densification, which cannot explain release within a breath, may be right about this slower family: hyaluronan in the gliding layer is where slow change in the fascia would live [46, 63].

6. Acupuncture points

Chinese medicine placed points along channels, and also took the tender spot itself as a point, the *ashi* point. If knots are perforators, it is natural to ask whether the classical points are too. Heine reported that about 80% of acupoints lie where a vessel–nerve bundle perforates the fascia [75]; the figure is widely repeated, but its method cannot be checked from the record. Dissection beneath 201 points found a vessel, mostly a vein, at about seven in ten, but no one structure common to all [76], and a multicentre dissection of four bodies and two lower legs found vessel–nerve bundles at only a few points [77]. Most points lie on the connective-tissue planes between muscles [78], which is where many perforators run. The one study we found with control points detected perforators by thermography at 87.5% of 144 lower-limb points and at 18.1% of matched control points, in six people, with no blinding described [79]; a dissection of two arms found a perforating vessel at all 30 points studied [80]. Control points a few centimetres away are not a measure of chance where perforators cluster, and none of these studies asked how often a point placed at random would land as near a vessel. Both maps are also fuzzy: 71 doctors' marks for one point spread over as much as 41 cm² [81], and perforator clusters spread by centimetres between people [12].

A first look against chance used five limb lines where a measured perforator cluster and a run of WHO points share an intermuscular septum [82]. The perforator density at the 21 points was about 1.23 times that at random points on the same lines, not reliably above chance (p 0.03–0.40 across 144 variants of the unstated details), where a map placed on the clusters would score 1.8–3.0 times. Some points sit on clusters, SP6 near the lowest posterior tibial perforator [83] and LU6 and LU7 near the radial artery's two clusters [84]; others do not. The correspondence is therefore not shown, and not excluded, and the hypothesis does not depend on it. Its nearest kin in the traditions is the *ashi* point, defined by tenderness alone.

7. Predictions, and what would refute them

Table 3. What the hypothesis predicts, how to look, and the result that would refute it.

Prediction	How to look	Refuted if
1. Tender points sit on perforators	Blinded palpation against a blinded Doppler or thermal map; distances against chance at several radii, with random and shifted maps	No excess over chance
2. A person has fixed stations	One person's knots marked over months against their own perforator map	New knots at arbitrary places, or continuous drift
3. Low flow before a release	Laser speckle over a knot's patch against its neighbours	Flow no lower than nearby
4. A flood at release	Laser speckle or a thermal camera at each felt release, with a pressed sham beside it	Release with no local rise, or a rise no larger than at a pressed sham
5. Release as pressure lifts	The same, with calibrated pressure	Releases under steady pressure, with flow rising while compressed
6. An order: flood, tingle, then sliding	Laser speckle, a button at the felt release, ultrasound of the layers sliding	Sliding returns before the flood

Prediction	How to look	Refuted if
7. Families	Freeing an upstream site against a downstream one	Small frees large as often as large frees small
8. Setting	Isolated small arteries carried shut for four hours against five minutes, then returned to resting tone	Those held for hours reopen at rest as fully as those held for minutes
9. Age	A census of knots, with the skin's microvascular reactivity at the same sites, across ages	No relation; children as knotted as adults
10. The fascia follows the knots	Ultrasound of the glide between fascial layers in people who report the slower changes, at the reported sites and matched sites, over days	No difference from matched sites

The first is the cheapest and decides the most: an afternoon with a pencil Doppler or a thermal camera, two blinded examiners and a statistician. The fourth is the most direct: a flood of skin flow, time-locked to a felt release and absent at a pressed sham. Thermal imaging after a mild cold challenge already finds perforators, its first hot spots lying over vessels that Doppler hears and CT angiography shows [85], and laser speckle imaging of skin flow is reproducible from week to week [86]. Because each alternative predicts a different trace at the moment of release (Table 2), one careful recording would classify releases rather than merely confirm one view. None of these studies asks anyone to seek the experiences of Section 5: each observes what already happens, at the usual pace. A one-page pilot is set out separately [87].

8. Limits

- The identification is untested. No one has yet mapped tender points against perforators blind, or recorded a release.
- The reports are informal, and partly the author's own.
- Finding tender points by touch is only moderately reliable from one examiner to another [61], so any map of knots must be made blind to the map of vessels (Prediction 1).
- A finger cannot feel a small artery shut beneath the fat. The hypothesis accounts for tenderness and the block, not for a firm nodule; where a nodule is found, it belongs to muscle, and that knot is not (only) a perforator.
- The hold is demanding in healthy vessels. That old knots hold at rest leans on adaptation and remodelling measured over hours and days in isolated small arteries, which relax incompletely rather than not at all.
- The evidence on age points both ways. The breath's effects are measured in the skin of the fingers and toes and in the nerves to muscle, not at the trunk's perforators, and which of its routes matters is unknown.
- Releases to attention alone are not explained.
- The account of the slower changes in the fascia (Section 5) is the most speculative part of the paper, and its reports are the hardest to check.
- The text was written by AI (below). Every reference was taken from its own record, and the numbers from the cited sources and the project's data, but errors of emphasis or reading are possible, and corrections are welcome.

9. Safety

Meditation and related practices can harm. Across 83 studies, adverse events were found in about 8% of those who meditated, anxiety and depression most often, sometimes in people with no previous mental-health problems [88]. In the Varieties of Contemplative Experience study, which recruited practitioners who could describe challenging or difficult experiences, symptoms lasted a median of one to three years, from a few days to more than ten; nearly three-quarters reported moderate to severe impairment in some area of life, about one in six suicidality, and as many a stay in hospital [1]. From the same kind of sample, meditators in the West reporting challenges, the energy-like somatic experiences to which the reports of Section 5 belong have been described with their trajectories and their effects on people's lives, and with the many ways practitioners, teachers, doctors and therapists read and managed them [71]. Attention to the body is not always benign: past some point, its effects can turn negative [74].

Nothing in this paper is a practice, a protocol or a recommendation, and nothing in it is a reason to seek these experiences: the hypothesis concerns what happens, not what anyone should do. Anyone in difficulty can find support from Cheetah House [89], whose team includes the researchers of the Varieties of Contemplative Experience project [90], and should also see a physician. Any study with people belongs under ethics review, with ordinary breathing, comfortable pressure and a plan for referral.

10. Conclusion

Knots are among the commonest things a hand finds in a body, and among the least explained. The perforator hypothesis assembles an explanation from parts that are each measured: fixed openings in the fascia; a wall that can hold itself shut, and remembers having been shut; a drive the breath reaches in seconds; a squeeze that widens; a nerve that fires when blood returns; a parent vessel that holds its branches shut. Put together, they describe a knot as a held state of a particular vessel at a particular place, written by the body's drive and kept by the vessel itself. It is, for now, the strongest account in a narrow sense: it explains more of what is reported, with less that is new, and it can be shown wrong quickly: the cheapest test needs an afternoon and a pencil Doppler. If tender points do not sit on perforators more often than chance, or a felt release brings no flood to the skin above it, it should be set aside. If they do,

the body keeps some of its history in the calibre of small vessels at fixed openings in its fascia: a register of held states that a hand can find, the breath can reach and an instrument can read.

Box 2. Open questions

- Do tender points sit on perforators more often than chance, and does each person's set of knots recur at the same places?
- Does the skin flood at the moment a knot lets go, and does the flood come before the tingle, and the tingle before the layers slide?
- Can a small human artery held shut for hours stay shut at resting tone?
- What carries a release to attention alone?
- What, if anything, changes in the fascia over the days after many knots let go?
- Are the classical points on perforators, or only the tender *ashi* points?

Authorship and AI use

This paper was written by AI under the author's direction. Eugene Jhong conceived the hypothesis, set out first in two essays [2, 4] and developed on the project's site, chose the question and the direction of the argument, and supplied the practitioner observations that motivate it. The text was drafted by Claude Opus 5.5 (Anthropic) from those essays, the site, the project's models and the cited literature. A draft was then refereed by a second model, Claude Fable 5.1 (Anthropic), which checked each claim against its source; its corrections were checked again against the sources before they were made. Every reference was taken from its own record, PubMed's for papers indexed there and the publisher's or Crossref's for the rest, rather than from a model's memory, and every number from the cited sources or the project's data. The author is responsible for the content.

Data and code

The models, every parameter with its source, and the code behind each number on this page are open [8]. The few numbers from the project's own calculations (the wall's switch, the hyaluronan and warming checks, the comparison with acupoints) are read from its data when the page is built.

Competing interests and funding

None.

References

1. Lindahl JR, Fisher NE, Cooper DJ et al. (2017). The varieties of contemplative experience: A mixed-methods study of meditation-related challenges in Western Buddhists. *PLoS One 12:e0176239*. doi.org/10.1371/journal.pone.0176239
2. Eugene Jhong (text by Kimi and Fable) (2026). Knots of Existence and Vasocomputation. *Substack, 14 August 2026*. ejhong.substack.com/p/knots-of-existence-and-vasocomputation
3. Michael Edward Johnson (2023). Principles of Vasocomputation: A Unification of Buddhist Phenomenology, Active Inference, and Physical Reflex (Part I). *opentheory.net, 12 July 2023 (updated 2025)*. opentheory.net/2023/07/principles-of-vasocomputation-a-unification-of-buddhist-phenomenology-active-inference-and-physical-reflex-part-i/
4. Eugene Jhong (text by Fable 5.1) (2026). Knots of Existence Hypotheses. *Substack, 8 September 2026*. ejhong.substack.com/p/knots-of-existence-hypotheses
5. Jhong E. Knots as pattern memories: a two-system model of somatic release in perforators and fascial planes. Unpublished manuscript (September 2026), available from the author. An earlier version of the account, with the collar as the hold; superseded by the present paper where they differ.
6. Nicholls M, Manjoo A, Shaw P et al. (2018). A Comparison Between Rheological Properties of Intra-articular Hyaluronic Acid Preparations and Reported Human Synovial Fluid. *Adv Ther 35:523-530*. doi.org/10.1007/s12325-018-0688-y
7. Fede C, Angelini A, Stern R et al. (2018). Quantification of hyaluronan in human fasciae: variations with function and anatomical site. *J Anat 233:552-556*. doi.org/10.1111/joa.12866
8. Knots of Existence: models, parameters and runs (2026). github.com/ejhong/knots
9. Krause WE, Bellomo EG, Colby RH (2001). Rheology of sodium hyaluronate under physiological conditions. *Biomacromolecules 2:65-9*. doi.org/10.1021/bm0055798
10. Jhong E. Knots: seven mechanisms, what each would do, and a recording that would tell them apart. Preprint (2026). ejhong.github.io/knots/research/preprint/
11. Taylor GI, Palmer JH (1987). The vascular territories (angiosomes) of the body: experimental study and clinical applications. *Br J Plast Surg 40:113-41*. doi.org/10.1016/0007-1226(87)90185-8
12. Martin AL, Bissell MB, Al-Dhamin A et al. (2013). Computed tomographic angiography for localization of the cutaneous perforators of the leg. *Plast Reconstr Surg 131:792-800*. doi.org/10.1097/PRS.0b013e3182818cb4
13. Rozen WM, Chubb D, Whitaker IS et al. (2010). Deep inferior epigastric perforators do not correlate between sides of the body: the role for preoperative imaging. *J Plast Reconstr Aesthet Surg 63:e842-3*. doi.org/10.1016/j.bjps.2010.07.030
14. Dowlatshahi S, Francalancia S, Willard FH et al. (2026). Structural and functional anatomy of the retinaculum cutis: review and new insights. *Anat Sci Int*. doi.org/10.1007/s12565-026-00936-8

15. Taylor GI, Minabe T (1992). The angiosomes of the mammals and other vertebrates. *Plast Reconstr Surg 89:181-215*. doi.org/10.1097/00006534-199202000-00001
16. Weum S, de Weerd L (2016). Perforator-Guided Drug Injection in the Treatment of Abdominal Wall Pain. *Pain Med 17:1229-32*. doi.org/10.1093/pm/pnv011
17. Boelens OB, Scheltinga MR, Houterman S et al. (2013). Randomized clinical trial of trigger point infiltration with lidocaine to diagnose anterior cutaneous nerve entrapment syndrome. *Br J Surg 100:217-21*. doi.org/10.1002/bjs.8958
18. Yoshinaga T, Koga T, Koga H (2025). Effectiveness of Ultrasound-Guided Hydrodissection for Atypical Anterior Cutaneous Nerve Entrapment Syndrome Presenting as Lower Back Pain Due to Retrograde Neuropathic Pain. *Cureus 17:e94871*. doi.org/10.7759/cureus.94871
19. Burton AC (1951). On the physical equilibrium of small blood vessels. *Am J Physiol 164:319-29*. doi.org/10.1152/ajplegacy.1951.164.2.319
20. Nichol J, Girling F, Jerrard W et al. (1951). Fundamental instability of the small blood vessels and critical closing pressures in vascular beds. *Am J Physiol 164:330-44*. doi.org/10.1152/ajplegacy.1951.164.2.330
21. Cutting C, Ballantyne D, Shaw W et al. (1982). Critical closing pressure, local perfusion pressure, and the failing skin flap. *Ann Plast Surg 8:504-9*. doi.org/10.1097/00000637-198206000-00009
22. Mulvany MJ, Warshaw DM (1979). The active tension-length curve of vascular smooth muscle related to its cellular components. *J Gen Physiol 74:85-104*. doi.org/10.1085/jgp.74.1.85
23. Miller A, Rola P, Spiegel R et al. (2026). Arteriolar Collapse and Haemodynamic Incoherence in Shock: Rethinking Critical Closing Pressure. *J Pers Med 16*. doi.org/10.3390/jpm16020078
24. Bohlen HG (1979). Arteriolar closure mediated by hyperresponsiveness to norepinephrine in hypertensive rats. *Am J Physiol 236:H157-64*. doi.org/10.1152/ajpheart.1979.236.1.H157
25. Schiffrin EL, Deng LY, Larochelle P (1995). Progressive improvement in the structure of resistance arteries of hypertensive patients after 2 years of treatment with an angiotensin I-converting enzyme inhibitor. Comparison with effects of a beta-blocker. *Am J Hypertens 8:229-36*. doi.org/10.1016/0895-7061(95)96211-2
26. Hill MA, Potocnik SJ, Martinez-Lemus LA et al. (2003). Delayed arteriolar relaxation after prolonged agonist exposure: functional remodeling involving tyrosine phosphorylation. *Am J Physiol Heart Circ Physiol 285:H849-56*. doi.org/10.1152/ajpheart.00986.2002
27. Martinez-Lemus LA, Hill MA, Bolz SS et al. (2004). Acute mechanoadaptation of vascular smooth muscle cells in response to continuous arteriolar vasoconstriction: implications for functional remodeling. *FASEB J 18:708-10*. doi.org/10.1096/fj.03-0634fje
28. Syyong H, Cheung C, Solomon D et al. (2008). Adaptive response of pulmonary arterial smooth muscle to length change. *J Appl Physiol (1985) 104:1014-20*. doi.org/10.1152/japplphysiol.01203.2007
29. Bednarek ML, Speich JE, Miner AS et al. (2011). Active tension adaptation at a shortened arterial muscle length: inhibition by cytochalasin-D. *Am J Physiol Heart Circ Physiol 300:H1166-73*. doi.org/10.1152/ajpheart.00009.2010
30. Bakker EN, van der Meulen ET, van den Berg BM et al. (2002). Inward remodeling follows chronic vasoconstriction in isolated resistance arteries. *J Vasc Res 39:12-20*. doi.org/10.1159/000048989
31. Charkoudian N (2003). Skin blood flow in adult human thermoregulation: how it works, when it does not, and why. *Mayo Clin Proc 78:603-12*. doi.org/10.4065/78.5.603
32. Fede C, Petrelli L, Pirri C et al. (2022). Innervation of human superficial fascia. *Front Neuroanat 16:981426*. doi.org/10.3389/fnana.2022.981426
33. Bolton B, Carmichael EA, Stürup G (1936). Vaso-constriction following deep inspiration. *J Physiol 86:83-94*. doi.org/10.1113/jphysiol.1936.sp003345
34. Lau YT, Liu CF, Tsai CC (1995). Cutaneous vasoconstrictor response induced by inspiratory gasp in relation to sex and age. *Clin Sci (Lond) 89:233-7*. doi.org/10.1042/cs0890233
35. Mayrovitz HN (2026). Physiological Impacts and Medical Applications of the Inspiratory Gasp Vascular Reflex. *Cureus 18:e105709*. doi.org/10.7759/cureus.105709
36. Adler TE, Coovadia Y, Cirone D et al. (2019). Device-guided slow breathing reduces blood pressure and sympathetic activity in young normotensive individuals of both sexes. *J Appl Physiol (1985) 127:1042-1049*. doi.org/10.1152/japplphysiol.00442.2019
37. Ljung B, Sivertsson R (1975). Vibration-induced inhibition of vascular smooth muscle contraction. *Blood Vessels 12:38-52*. doi.org/10.1159/000158037
38. Clifford PS, Kluess HA, Hamann JJ et al. (2006). Mechanical compression elicits vasodilatation in rat skeletal muscle feed arteries. *J Physiol 572:561-7*. doi.org/10.1113/jphysiol.2005.099507
39. Minson CT, Berry LT, Joyner MJ (2001). Nitric oxide and neurally mediated regulation of skin blood flow during local heating. *J Appl Physiol (1985) 91:1619-26*. doi.org/10.1152/jappl.2001.91.4.1619
40. Taylor GI, Corlett RJ, Dhar SC et al. (2011). The anatomical (angiosome) and clinical territories of cutaneous perforating arteries: development of the concept and designing safe flaps. *Plast Reconstr Surg 127:1447-1459*. doi.org/10.1097/PRS.0b013e318208d21b
41. Shah JP, Phillips TM, Danoff JV et al. (2005). An in vivo microanalytical technique for measuring the local biochemical milieu of human skeletal muscle. *J Appl Physiol (1985) 99:1977-84*. doi.org/10.1152/japplphysiol.00419.2005

42. Alexandrou ME, Gkaliagkousi E, Loutradis C et al. (2021). Haemodialysis and peritoneal dialysis patients have severely impaired post-occlusive skin forearm vasodilatory response assessed with laser speckle contrast imaging. *Clin Kidney J 14:1419-1427*. doi.org/10.1093/ckj/sfaa136
43. Lorenzo S, Minson CT (2007). Human cutaneous reactive hyperaemia: role of BKCa channels and sensory nerves. *J Physiol 585:295-303*. doi.org/10.1113/jphysiol.2007.143867
44. Ochoa JL, Torebjörk HE (1980). Paraesthesiae from ectopic impulse generation in human sensory nerves. *Brain 103:835-53*. doi.org/10.1093/brain/103.4.835
45. Guimberteau JC, Sentucq-Rigall J, Panconi B et al. (2005). [Introduction to the knowledge of subcutaneous sliding system in humans]. *Ann Chir Plast Esthet 50:19-34*. doi.org/10.1016/j.anplas.2004.10.012
46. Cowman MK, Schmidt TA, Raghavan P et al. (2015). Viscoelastic Properties of Hyaluronan in Physiological Conditions. *F1000Res 4:622*. doi.org/10.12688/f1000research.6885.1
47. Coussot P, Nguyen QD, Huynh HT et al. (2002). Avalanche behavior in yield stress fluids. *Phys Rev Lett 88:175501*. doi.org/10.1103/PhysRevLett.88.175501
48. Langevin HM, Fox JR, Koptiuch C et al. (2011). Reduced thoracolumbar fascia shear strain in human chronic low back pain. *BMC Musculoskelet Disord 12:203*. doi.org/10.1186/1471-2474-12-203
49. Stecco A, Meneghini A, Stern R et al. (2014). Ultrasonography in myofascial neck pain: randomized clinical trial for diagnosis and follow-up. *Surg Radiol Anat 36:243-53*. doi.org/10.1007/s00276-013-1185-2
50. Segal SS, Duling BR (1986). Flow control among microvessels coordinated by intercellular conduction. *Science 234:868-70*. doi.org/10.1126/science.3775368
51. Segal SS, Damon DN, Duling BR (1989). Propagation of vasomotor responses coordinates arteriolar resistances. *Am J Physiol 256:H832-7*. doi.org/10.1152/ajpheart.1989.256.3.H832
52. Hsieh YL, Kao MJ, Kuan TS et al. (2007). Dry needling to a key myofascial trigger point may reduce the irritability of satellite MTrPs. *Am J Phys Med Rehabil 86:397-403*. doi.org/10.1097/PHM.0b013e31804a554d
53. El Assar M, García-Rojo E, Sevilleja-Ortiz A et al. (2022). Functional Role of STIM-1 and Orai1 in Human Microvascular Aging. *Cells 11*. doi.org/10.3390/cells11223675
54. Ng AV, Callister R, Johnson DG et al. (1993). Age and gender influence muscle sympathetic nerve activity at rest in healthy humans. *Hypertension 21:498-503*. doi.org/10.1161/01.hyp.21.4.498
55. Vionnet J, Calero-Romero I, Heim A et al. (2014). No major impact of skin aging on the response of skin blood flow to a submaximal local thermal stimulus. *Microcirculation 21:730-7*. doi.org/10.1111/micc.12154
56. Holowatz LA, Kenney WL (2010). Peripheral mechanisms of thermoregulatory control of skin blood flow in aged humans. *J Appl Physiol (1985) 109:1538-44*. doi.org/10.1152/japplphysiol.00338.2010
57. Gerwin RD, Dommerholt J, Shah JP (2004). An expansion of Simons' integrated hypothesis of trigger point formation. *Curr Pain Headache Rep 8:468-75*. doi.org/10.1007/s11916-004-0069-x
58. Sikdar S, Shah JP, Gebreab T et al. (2009). Novel applications of ultrasound technology to visualize and characterize myofascial trigger points and surrounding soft tissue. *Arch Phys Med Rehabil 90:1829-38*. doi.org/10.1016/j.apmr.2009.04.015
59. Hubbard DR, Berkoff GM (1993). Myofascial trigger points show spontaneous needle EMG activity. *Spine (Phila Pa 1976) 18:1803-7*. doi.org/10.1097/00007632-199310000-00015
60. Moraska AF, Hickner RC, Kohrt WM et al. (2013). Changes in blood flow and cellular metabolism at a myofascial trigger point with trigger point release (ischemic compression): a proof-of-principle pilot study. *Arch Phys Med Rehabil 94:196-200*. doi.org/10.1016/j.apmr.2012.08.216
61. Rathbone ATL, Grosman-Rimon L, Kumbhare DA (2017). Interrater Agreement of Manual Palpation for Identification of Myofascial Trigger Points: A Systematic Review and Meta-Analysis. *Clin J Pain 33:715-729*. doi.org/10.1097/AJP.0000000000000459
62. Quintner JL, Bove GM, Cohen ML (2015). A critical evaluation of the trigger point phenomenon. *Rheumatology (Oxford) 54:392-9*. doi.org/10.1093/rheumatology/keu471
63. Stecco C, Stern R, Porzionato A et al. (2011). Hyaluronan within fascia in the etiology of myofascial pain. *Surg Radiol Anat 33:891-6*. doi.org/10.1007/s00276-011-0876-9
64. Quintner JL, Cohen ML (1994). Referred pain of peripheral nerve origin: an alternative to the "myofascial pain" construct. *Clin J Pain 10:243-51*. doi.org/10.1097/00002508-199409000-00012
65. Dilley A, Lynn B, Pang SJ (2005). Pressure and stretch mechanosensitivity of peripheral nerve fibres following local inflammation of the nerve trunk. *Pain 117:462-472*. doi.org/10.1016/j.pain.2005.08.018
66. Devor M, Jänig W, Michaelis M (1994). Modulation of activity in dorsal root ganglion neurons by sympathetic activation in nerve-injured rats. *J Neurophysiol 71:38-47*. doi.org/10.1152/jn.1994.71.1.38
67. Rembold CM (1991). Relaxation, [Ca2+]i, and the latch-bridge hypothesis in swine arterial smooth muscle. *Am J Physiol 261:C41-50*. doi.org/10.1152/ajpcell.1991.261.1.C41
68. Gorassini MA, Bennett DJ, Yang JF (1998). Self-sustained firing of human motor units. *Neurosci Lett 247:13-6*. doi.org/10.1016/s0304-3940(98)00277-8

69. Heckman CJ, Johnson M, Mottram C et al. (2008). Persistent inward currents in spinal motoneurons and their influence on human motoneuron firing patterns. *Neuroscientist 14:264-75*. doi.org/10.1177/1073858408314986
70. Woolf CJ (2011). Central sensitization: implications for the diagnosis and treatment of pain. *Pain 152:S2-S15*. doi.org/10.1016/j.pain.2010.09.030
71. Cooper DJ, Lindahl JR, Palitsky R, Britton WB (2021). "Like a Vibration Cascading through the Body": Energy-Like Somatic Experiences Reported by Western Buddhist Meditators. *Religions 12:1042*. doi.org/10.3390/rel12121042
72. Susan Standring (ed.) (2020). Gray's Anatomy: The Anatomical Basis of Clinical Practice, 42nd edition. *Elsevier*.
73. Nakajima H, Imanishi N, Minabe T et al. (2004). Anatomical study of subcutaneous adipofascial tissue: a concept of the protective adipofascial system (PAFS) and lubricant adipofascial system (LAFS). *Scand J Plast Reconstr Surg Hand Surg 38:261-6*. doi.org/10.1080/02844310410029543
74. Britton WB (2019). Can mindfulness be too much of a good thing? The value of a middle way. *Curr Opin Psychol 28:159-165*. doi.org/10.1016/j.copsyc.2018.12.011
75. Heine H (1988). Anatomical structure of acupoints. *J Tradit Chin Med 8:207-12*. pubmed.ncbi.nlm.nih.gov/3216666/
76. Bossy J, Maurel JC, Godlewski G (1975). [Macroscopic substratum of acupuncture points]. *Bull Assoc Anat (Nancy) 59:357-62*.
77. Maurer N, Nissel H, Egerbacher M et al. (2019). Anatomical Evidence of Acupuncture Meridians in the Human Extracellular Matrix: Results from a Macroscopic and Microscopic Interdisciplinary Multicentre Study on Human Corpses. *Evid Based Complement Alternat Med 2019:6976892*. doi.org/10.1155/2019/6976892
78. Langevin HM, Yandow JA (2002). Relationship of acupuncture points and meridians to connective tissue planes. *Anat Rec 269:257-65*. doi.org/10.1002/ar.10185
79. Álvarez-Prats D, Carvajal-Fernández O, Valera Garrido F et al. (2019). Acupuncture Points and Perforating Cutaneous Vessels Identified Using Infrared Thermography: A Cross-Sectional Pilot Study. *Evid Based Complement Alternat Med 2019:7126439*. doi.org/10.1155/2019/7126439
80. Meng S, Tang Z, Zhu M et al. (2024). Perforating cutaneous vessels: A key feature of acupoints - Anatomical evidence from five-Shu acupoints in the upper limbs. *Clin Anat 37:33-42*. doi.org/10.1002/ca.24077
81. Molsberger AF, Manickavasagan J, Abholz HH et al. (2012). Acupuncture points are large fields: the fuzziness of acupuncture point localization by doctors in practice. *Eur J Pain 16:1264-70*. doi.org/10.1002/j.1532-2149.2012.00145.x
82. Knots of Existence. Are the perforators where the points are? (2026). ejhong.github.io/knots/research/#points
83. Bulla A, De Luca L, Campus GV et al. (2015). The localization of the distal perforators of posterior tibial artery: a cadaveric study for the correct planning of medial adipofascial flaps. *Surg Radiol Anat 37:19-25*. doi.org/10.1007/s00276-014-1275-9
84. Saint-Cyr M, Mujadzic M, Wong C et al. (2010). The Radial Artery Pedicle Perforator Flap: Vascular Analysis and Clinical Implications. *Plast Reconstr Surg 125:1469-1478*. doi.org/10.1097/PRS.0b013e3181d511e7
85. Weum S, Mercer JB, de Weerd L (2016). Evaluation of dynamic infrared thermography as an alternative to CT angiography for perforator mapping in breast reconstruction: a clinical study. *BMC Med Imaging 16:43*. doi.org/10.1186/s12880-016-0144-x
86. Roustit M, Millet C, Blaise S et al. (2010). Excellent reproducibility of laser speckle contrast imaging to assess skin microvascular reactivity. *Microvasc Res 80:505-11*. doi.org/10.1016/j.mvr.2010.05.012
87. Jhong E. A pilot recording at the moment a knot lets go: one-page protocol (2026). ejhong.github.io/knots/research/pilot/
88. Farias M, Maraldi E, Wallenkampf KC et al. (2020). Adverse events in meditation practices and meditation-based therapies: a systematic review. *Acta Psychiatr Scand 142:374-393*. doi.org/10.1111/acps.13225
89. Cheetah House (2018). Support for meditators in difficulty. *cheetahhouse.org*. www.cheetahhouse.org
90. Willoughby Britton and colleagues (2017). The Varieties of Contemplative Experience. *Brown University*. sites.brown.edu/britton/research/the-varieties-of-contemplative-experience/